abcam

ab101531

Anti-CD44 antibody [SP37]

Lab Essentials | Recombinant | RabMAb | KO Validated | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.6 Preservative: 0.1% Sodium azide Constituents: PBS, 1% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	SP37
Purification technique	Affinity purification Protein A/G
Concentration	0.05 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →
Purification notes	Purified from TCS by protein A/G.

Reactivity data

WB

Tested

Species	Human
Dilution info	-
Notes	-

Predicted

Species	Cow, Dog
Dilution info	-
Notes	-

IHC-P

Tested

Species	Human
Dilution info	1/100
Notes	Boil tissue section in citrate buffer pH 6.0 for 10 minutes followed by cooling at room temperature for 20 minutes. Incubate with primary antibody for 30 minutes at room temperature.

Predicted

Species	Cow, Dog
Dilution info	-
Notes	-

Flow Cyt

Tested

Species	Human
Dilution info	-
Notes	-

Predicted

Species	Cow, Dog
Dilution info	-
Notes	-

Target data

See full target information CD44

Function	Cell-surface receptor that plays a role in cell-cell interactions, cell adhesion and migration, helping them to sense and respond to changes in the tissue microenvironment (PubMed:16541107, PubMed:19703720, PubMed:22726066). Participates thereby in a wide variety of cellular functions including the activation, recirculation and homing of T-lymphocytes, hematopoiesis, inflammation and response to bacterial infection (PubMed:7528188). Engages, through its ectodomain, extracellular matrix components such as hyaluronan/HA, collagen, growth factors, cytokines or proteases and serves as a platform for signal transduction by assembling, via its cytoplasmic domain, protein complexes containing receptor kinases and membrane proteases (PubMed:18757307, PubMed:23589287). Such effectors include PKN2, the RhoGTPases RAC1 and RHOA, Rho-kinases and phospholipase C that coordinate signaling pathways promoting calcium mobilization and actin-mediated cytoskeleton reorganization essential for cell migration and adhesion (PubMed:15123640). Upon interaction with LGALS9 ligand, activates downstream signaling components including LCK, ERK and MAPK to promotes NK cell activation (PubMed:37006235). (Microbial infection) Promotes foot-and-mouth disease virus internalization via macropinocytosis.

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended

We do not recommend this combination. It is not covered by our product promise.


We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

---


6 product images



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD44 antibody [SP37] (ab101531)

CD44 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-CD44 antibody


Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of Human esophageal carcinoma tissue sections labeling CD44 with Purified ab101531 at 1/100 dilution (1.03 µg/ml). Heat mediated antigen retrieval was performed Heat mediated antigen retrieval with sodium citrate buffer (pH 6.0, epitope retrieval solution 1) for 20mins. Rabbit specific IHC polymer detection kit HRP/DAB (ab209101) was used as the secondary antibody. Negative control: PBS instead of the primary antibody. Hematoxylin was used as a counterstain.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD44 antibody [SP37] (ab101531)

CD44 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-CD44 antibody


Formalin-fixed, paraffin-embedded human esophageal carcinoma tissue stained for CD44 with ab101531 at a 1/100 dilution in immunohistochemical analysis.



Flow Cytometry - Anti-CD44 antibody [SP37] (ab101531)

Flow cytometric analysis of Jurkat (human T cell leukemia T lymphocyte from peripheral blood, Left) / A549 (human lung carcinoma epithelial cell, Right) cells labelling CD44 with ab101531 at 1/50 dilution (1µg) (Red) followed by a Goat Anti-Rabbit IgG (Alexa Fluor® 488, ab150081) secondary antibody used at a 1/5000 dilution compared with a Rabbit monoclonal IgG (ab172730) (Black) isotype control and an unlabelled control (cells without incubation with primary antibody and secondary antibody).

Gated on viable cells.
Negative control: Jurkat.



Western blot - Anti-CD44 antibody [SP37] (ab101531)

CD44 Western blot staining using rabbit Anti-CD44 antibody

Lanes 1 - 4: Merged signal (red and green). Green - ab101531 observed at 80 kDa. Red - loading control, ab8245 (Mouse anti-GAPDH antibody [6C5]) observed at 37kDa.
ab101531 was shown to react with CD44 in wild-type HeLa cells in western blot. Loss of signal was observed when CD44 knockout sample was used. Wild-type HeLa and CD44 knockout cell lysates were subjected to SDS-PAGE. Membranes were blocked in 3% milk in TBS-T (0.1% Tween®) before incubation with ab101531 and ab8245 (Mouse anti-GAPDH antibody [6C5]) overnight at 4°C at a 1 in 1000 dilution and a 1 in 20000 dilution respectively. Blots were incubated with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preabsorbed (ab216773) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preabsorbed (ab216776) secondary antibodies at 1 in 20000 dilution for 1 hour at room temperature before imaging.

All lanes:
Western blot - Anti-CD44 antibody [SP37] (ab101531) at 1/1000 dilution

Lane 1:
Wild-type HeLa (Human epithelial cell line from cervix adenocarcinoma) whole cell lysate at 20 µg

Lane 2:
Western blot - Human CD44 knockout HeLa cell lysate (ab263938) at 20 µg


Lane 3:
A549 (Human lung carcinoma cell line) whole cell lysate at 20 µg

Lane 4:
LNCaP (Human prostate cancer cell line) whole cell lysate at 20 µg

Performed under reducing conditions.

Predicted band size: 81 kDa

Observed band size: 80 kDa



Western blot - Anti-CD44 antibody [SP37] (ab101531)

CD44 Western blot staining using rabbit Anti-CD44 antibody

False colour image of Western blot: Anti-CD44 antibody [SP37] staining at 1/1000 dilution, shown in green; Mouse anti-Alpha Tubulin [DM1A] (ab7291) loading control staining at 1/20000 dilution, shown in red. In Western blot, ab101531 was shown to bind specifically to CD44. A band was observed at 75-80 kDa in wild-type HeLa cell lysates with no signal observed at this size in CD44 knockout cell line ab262515 (knockout cell lysate ab263938). To generate this image, wild-type and CD44 knockout HeLa cell lysates were analysed. First, samples were run on an SDS-PAGE gel then transferred onto a nitrocellulose membrane. Membranes were blocked in 3 % milk in TBS-0.1 % Tween® 20 (TBS-T) before incubation with primary antibodies overnight at 4°C. Blots were washed four times in TBS-T, incubated with secondary antibodies for 1 h at room temperature, washed again four times then imaged. Secondary antibodies used were Goat anti-Rabbit IgG H&L (IRDye® 800CW) preabsorbed (ab216773) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preabsorbed (ab216776) at 1/20000 dilution.
All lanes:
Western blot - Anti-CD44 antibody [SP37] (ab101531) at 1/1000 dilution

Lane 1:
Wild-type HeLa cell lysate at 20 µg

Lane 2:
Western blot - Human CD44 knockout HeLa cell lysate (ab263938) at 20 µg


Lane 3:
A549 cell lysate at 20 µg

Lane 4:
LNCaP cell lysate at 20 µg

Performed under reducing conditions.

Predicted band size: 81 kDa

Observed band size: 75-80 kDa



Western blot - Anti-CD44 antibody [SP37] (ab101531)

All lanes:
Western blot - Anti-CD44 antibody [SP37] (ab101531) at 1/25 dilution

All lanes:
HeLa cell lysate

Predicted band size: 41 kDa, 81 kDa

Observed band size: 80 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.