abcam

ab101532

Anti-CDX2 antibody [SP54]

Recombinant | RabMAb | Advanced Validation | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.1% Sodium azide Constituents: PBS, 1% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	SP54
Purification technique	Affinity purification Protein A/G
Concentration	0.052 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →
Purification notes	Purified from TCS by protein A/G.

Reactivity data

IHC-P

Tested

Species	Human
Dilution info	1/100
Notes	Boil tissue section in 10mM Citrate buffer, pH 6.0 for 10-20 min followed by cooling at RT for 20 min.

	Perform heat-mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.

Predicted

Species	Mouse
Dilution info	-
Notes	-

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	1/200
Notes	ab199376 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.

Predicted

Species	Mouse
Dilution info	-
Notes	-

ICC/IF

Tested

Species	Human
Dilution info	1/50
Notes	-

Predicted

Species	Mouse
Dilution info	-
Notes	-

ChIC/CUT&RUN-seq

Tested

Species	Human
Dilution info	5 µg
Notes	-

Predicted

Species	Mouse
Dilution info	-
Notes	-

Target data

See full target information CDX2

Function Transcription factor which regulates the transcription of multiple genes expressed in the intestinal epithelium (By similarity). Binds to the promoter of the intestinal sucrase-isomaltase SI and activates SI transcription (By similarity). Binds to the DNA sequence 5'-ATAAAAACTTAT-3' in the promoter region of VDR and activates VDR transcription (By similarity). Binds to and activates transcription of LPH (By similarity). Activates transcription of CLDN2 and intestinal mucin MUC2 (By similarity). Binds to the 5'-AATTTTTTACAACACCT-3' DNA sequence in the promoter region of CA1 and activates CA1 transcription (By similarity). Important in broad range of functions from early differentiation to maintenance of the intestinal epithelial lining of both the small and large intestine. Binds preferentially to methylated DNA (PubMed:28473536).

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C

Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	CDX2 or Caudal Type Homeobox 2 is a transcription factor involved in intestinal development and differentiation. Alternative names for CDX2 include Cdx-2 and CDX-2. This protein has a molecular mass of approximately 38 kDa. CDX2 is expressed mainly in the epithelial cells of the intestine. Researchers can identify CDX2 through techniques like CDX2 stain CDX2 immunostain and CDX2 IHC which are useful for determining its expression in tissue samples.
Biological function summary	CDX2 plays an important role in controlling the expression of genes necessary for maintaining intestinal epithelial cell identity. It functions as a part of regulatory networks that guide cell fate and differentiation in the gut. CDX2 does not generally form complexes but acts as a transcriptional regulator that binds directly to DNA influencing genes involved in cell adhesion proliferation and the maintenance of tissue structure.
Pathways	CDX2 is integral to the Wnt signaling and Hedgehog pathways which are vital for intestinal development and homeostasis. Within these pathways CDX2 works closely with other proteins like beta-catenin influencing cell proliferation and differentiation. The Wnt pathway in particular modulates the expression of proliferative genes while CDX2 ensures that proliferation maintains tissue integrity by guiding cellular differentiation.
Associated diseases and disorders	CDX2 has a significant link to colorectal cancer and Barrett's esophagus. Loss of CDX2 expression often marks the progression from normal to malignant tissue in colorectal cancer where it connects with proteins like p-catenin affecting cellular growth and differentiation. Similarly abnormal expression of CDX2 correlates with the development of Barrett's esophagus a condition that may progress to esophageal adenocarcinoma. These disease associations make CDX2 a potential diagnostic marker and therapeutic target in gastrointestinal pathologies.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended

We do not recommend this combination. It is not covered by our product promise.


We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

---


6 product images



Flow Cytometry (Intracellular) - Anti-CDX2 antibody [SP54] (ab101532)

CDX2 Flow Cytometry (Intracellular) staining using rabbit Anti-CDX2 antibody

Flow cytometry analysis of Caco-2 (human colorectal adenocarcinoma epithelial cell) labeling CDX2 with purified ab101532 at 1/200 dilution (0.56 µg/ml) (red). Cells were fixed with 4% paraformaldehyde and permeabilised with 90% methanol. Goat anti rabbit IgG (Alexa Fluor® 488, ab150077) at 1/2000 dilution was used as a secondary antibody. Isotype control -Rabbit monoclonal IgG (ab172730) (black). Unlableled control -Unlabelled cells (blue).




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CDX2 antibody [SP54] (ab101532)

CDX2 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-CDX2 antibody

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of Human colon carcinoma tissue sections labeling CDX2 with ab101532 at 1/100 dilution (1.11 µg/ml). Heat mediated antigen retrieval with Citrate buffer (pH 6.0, epitope retrieval solution 1) for 20 mins. Goat Anti-Rabbit & Mouse IgG (HRP) was used as the secondary antibody. Hematoxylin was used


as a counterstain. Positive staining on the human colon carcinoma, performed on a Leica Biosystems BOND™ RX instrument.
The section was incubated with ab101532 for 30 mins at room temperature.



Immunocytochemistry/ Immunofluorescence - Anti-CDX2 antibody [SP54] (ab101532)

CDX2 Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-CDX2 antibody


Immunocytochemistry/ Immunofluorescence analysis of Caco-2 (human colorectal adenocarcinoma epithelial cell) cells labeling CDX2 with purified ab101532 at 1/50 (2.22 µg/ml). Cells were fixed in 4% paraformaldehyde and permeabilized with 0.1% Triton X-100. Cells were counterstained with ab195889 Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor® 594) 1/200 (2.5 µg/ml). Goat anti rabbit IgG (Alexa Fluor® 488, ab150077) was used as the secondary antibody at 1/1000 (2 µg/ml) dilution. DAPI was used as nuclear counterstain. PBS instead of the primary antibody was used as the secondary antibody only control.



ChIC/CUT&RUN sequencing - Anti-CDX2 antibody [SP54] (ab101532)

ChIC/CUT&RUN was performed using a pAG-MNase at a final concentration of 700 ng/mL, 2.5 x 10^5 Caco-2 (human colorectal adenocarcinoma epithelial cell) cells and 5 µg of ab101532 [SP54]. The resulting DNA was sequenced on the Illumina NovaSeq 6000 to a depth of 10 million reads. The negative IgG control ab172730 is also shown.


The University of Geneva owns patents relevant to ChIC (Chromatin Immuno-Cleavage) methods.



ChIC/CUT&RUN sequencing - Anti-CDX2 antibody [SP54] (ab101532)

ChIC/CUT&RUN was performed using a pAG-MNase at a final concentration of 700 ng/mL, 2.5 x 10^5 Caco-2 (human colorectal adenocarcinoma epithelial cell) cells and 5 µg of ab101532 [SP54]. The resulting DNA was sequenced on the Illumina NovaSeq 6000 to a depth of 10 million reads. The negative IgG control ab172730 is also shown.

The University of Geneva owns patents relevant to ChIC (Chromatin Immuno-Cleavage) methods.



ChIC/CUT&RUN sequencing - Anti-CDX2 antibody [SP54] (ab101532)

ChIC/CUT&RUN was performed using a pAG-MNase at a final concentration of 700 ng/mL, 2.5 x 10^5 Caco-2 (human colorectal adenocarcinoma epithelial cell) cells and 5 µg of ab101532 [SP54]. The resulting DNA was sequenced on the Illumina NovaSeq 6000 to a depth of 10 million reads. The negative IgG control ab172730 is also shown.

The University of Geneva owns patents relevant to ChIC (Chromatin Immuno-Cleavage) methods.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.