abcam

ab108417

Anti-Cytokeratin 14 antibody [EPR1612]

Recombinant | RabMAb | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR1612
Purification technique	Affinity purification Protein A
Concentration	0.104 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

ICC/IF

Tested

Species	Human
Dilution info	1/100 - 1/500
Notes	-

IHC-P

Tested

Species	Human
Dilution info	0.001 µg/mL
Notes	Antigen retrieval is recommended.

Flow Cyt

Tested

Species	Human
Dilution info	-
Notes	-

WB

Not recommended

Species	Human
Dilution info	-
Notes	-

IP

Not recommended

Species	Human
Dilution info	-
Notes	-

Target data

See full target information Keratin, type I cytoskeletal 14

Function	The nonhelical tail domain is involved in promoting KRT5-KRT14 filaments to self-organize into large bundles and enhances the mechanical properties involved in resilience of keratin intermediate filaments in vitro.

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.


12 product images



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Cytokeratin 14 antibody [EPR1612] (ab108417)

Cytokeratin 14 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining of A431 WT cell pellet using rabbit Anti-Cytokeratin 14 antibody


IHC image of Cytokeratin 14 staining in a section of formalin-fixed paraffin-embedded A431 WT cell pellet block performed on a Leica Biosystems BOND® RX instrument. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH6, epitope retrieval solution 1) for 20mins. The section was then incubated with ab108417, 0.001ug/ml, for 15 mins at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX. The inset secondary-only control image is taken from an identical assay without primary antibody.



Flow Cytometry - Anti-Cytokeratin 14 antibody [EPR1612] (ab108417)

Cytokeratin 14 Flow Cytometry staining of A431 (human epidermoid carcinoma) cells using rabbit Anti-Cytokeratin 14 antibody

Flow Cytometry analysis of A431 (human epidermoid carcinoma) cells labeling Cytokeratin 14 with purified ab108417 at 1:250 dilution(10ug/ml) (red). Cells were fixed with 4% paraformaldehyde and permeabilised with 90% methanol. A Goat anti rabbit IgG (Alexa Fluor® 488)(ab150077)(1:2000 dilution) was used as the secondary antibody. Rabbit monoclonal IgG (Black)(ab172730) was used as the isotype control, Cell without incubation with primary antibody and secondary antibody (Blue) were used as the unlabeled control.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Cytokeratin 14 antibody [EPR1612] (ab108417)

Cytokeratin 14 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining of Breast carcinoma tissue using rabbit Anti-Cytokeratin 14 antibody

ab108417 showing negative staining in Breast carcinoma tissue.

Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Cytokeratin 14 antibody [EPR1612] (ab108417)

Cytokeratin 14 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining of Colonic adenocarcinoma tissue using rabbit Anti-Cytokeratin 14 antibody

ab108417 showing negative staining in Colonic adenocarcinoma tissue.


Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.



Immunocytochemistry/ Immunofluorescence - Anti-Cytokeratin 14 antibody [EPR1612] (ab108417)

Cytokeratin 14 Immunocytochemistry/ Immunofluorescence staining of Wild-type A431 cells using rabbit Anti-Cytokeratin 14 antibody

ab108417 staining KRT14 in wild-type A431 cells (top panel) and KRT14 knockout A431 cells (bottom panel). The cells were fixed with 100% methanol (5 min) then permeabilized with 0.1% Triton X-100 for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween for 1h. The cells were then incubated with ab108417 at 1/100 dilution and ab7291 (Mouse monoclonal to alpha Tubulin) at 1/1000 dilution overnight at 4°C followed by a further incubation at room temperature for 1h with a goat secondary antibody to rabbit IgG (Alexa Fluor® 488) (ab150081) at 2 µg/ml (shown in green) and a goat secondary antibody to mouse IgG (Alexa Fluor® 594) (ab150120) at 2 µg/ml (shown in red). Nuclear DNA was labelled in blue with DAPI. Image was taken with a confocal microscope (Leica-Microsystems TCS SP8).




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Cytokeratin 14 antibody [EPR1612] (ab108417)

Cytokeratin 14 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining of Human squamous cell cervical carcinoma tissue using rabbit Anti-Cytokeratin 14 antibody

ab108417, at a 1/250 dilution, staining Cytokeratin 14 in paraffin embedded Human squamous cell cervical carcinoma tissue by Immunohistochemistry.

Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Cytokeratin 14 antibody [EPR1612] (ab108417)

Cytokeratin 14 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining of Normal prostate basal epithelium tissue using rabbit Anti-Cytokeratin 14 antibody

ab108417 showing positive staining in Normal prostate basal epithelium tissue.

Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Cytokeratin 14 antibody [EPR1612] (ab108417)

Cytokeratin 14 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining of Human tonsil tissue using rabbit Anti-Cytokeratin 14 antibody

ab108417, at a 1/250 dilution, staining Cytokeratin 14 in paraffin embedded Human tonsil tissue by Immunohistochemistry.


Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Cytokeratin 14 antibody [EPR1612] (ab108417)

Cytokeratin 14 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining of A431 KO cell pellet using rabbit Anti-Cytokeratin 14 antibody

Negative control image: IHC image of Cytokeratin 14 staining in a section of formalin-fixed paraffin-embedded A431 KO cell pellet block performed on a Leica Biosystems BOND® RX instrument. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH6, epitope retrieval solution 1) for 20mins. The section was then incubated with ab108417, 0.001ug/ml, for 15 mins at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX. The inset secondary-only control image is taken from an identical assay without primary antibody.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Cytokeratin 14 antibody [EPR1612] (ab108417)

Cytokeratin 14 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining of Normal kidney tissue using rabbit Anti-Cytokeratin 14 antibody

ab108417 showing negative staining in Normal kidney tissue.

Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Cytokeratin 14 antibody [EPR1612] (ab108417)

Cytokeratin 14 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining of Normal skin tissue using rabbit Anti-Cytokeratin 14 antibody

ab108417 showing positive staining in Normal skin tissue.

Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Cytokeratin 14 antibody [EPR1612] (ab108417)

Cytokeratin 14 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining of Human skin tissue using rabbit Anti-Cytokeratin 14 antibody

ab108417, at a 1/250 dilution, staining Cytokeratin 14 in paraffin embedded Human skin tissue by Immunohistochemistry.

Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.