abcam

ab108928

Anti-FANCD2 antibody [EPR2302]

Anti-FANCD2 antibody [EPR2302] (ab108928) is a rabbit monoclonal antibody detecting FANCD2 in **Western Blot, Flow Cytometry (Intra), IP, IHC-P, ICC/IF**. Suitable for **Human, Mouse, Rat**.

- KO validated for confirmed specificity
- Biophysical QC for unrivalled batch-batch consistency
- Over 40 publications

Recombinant | RabMAb | KO Validated | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR2302
Purification technique	Affinity purification Protein A
Concentration	0.182 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

IHC-P

Tested

Species	Human
Dilution info	1/50 - 1/250
Notes	Perform heat-mediated antigen retrieval before commencing with IHC staining protocol.

Not recommended

Species	Mouse, Rat
Dilution info	-
Notes	-

IP

Tested

Species	Human
Dilution info	1/10 - 1/100
Notes	-

Expected

Species	Mouse, Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

WB

Tested

Species	Mouse
Dilution info	1/1000 - 1/10000
Notes	-

Species	Rat
Dilution info	1/1000 - 1/10000
Notes	-

Species	Human
Dilution info	1/1000 - 1/10000
Notes	-

ICC/IF

Tested

Species	Human
Dilution info	1/250 - 1/500
Notes	-

Expected

Species	Mouse, Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	1/200
Notes	-

Expected

Species	Mouse, Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

Target data

See full target information FANCD2

Function	Required for maintenance of chromosomal stability (PubMed:11239453, PubMed:14517836). Promotes accurate and efficient pairing of homologs during meiosis (PubMed:14517836). Involved in the repair of DNA double-strand breaks, both by homologous recombination and single-strand annealing (PubMed:15671039, PubMed:15650050, PubMed:30335751, PubMed:36385258). The FANCI-FANCD2 complex binds and scans double-stranded DNA (dsDNA) for DNA damage; this complex stalls at DNA junctions between double-stranded DNA and single-stranded DNA (By similarity). May participate in S phase and G2 phase checkpoint activation upon DNA damage (PubMed:15377654). Plays a role in preventing breakage and loss of missegregating chromatin at the end of cell division, particularly after replication stress (PubMed:15454491, PubMed:15661754). Required for the targeting, or stabilization, of BLM to non-centromeric abnormal structures induced by replicative stress (PubMed:15661754, PubMed:19465921). Promotes BRCA2/FANCD1 loading onto damaged chromatin (PubMed:11239454, PubMed:12239151, PubMed:12086603, PubMed:15115758, PubMed:15199141, PubMed:15671039, PubMed:18212739). May also be involved in B-cell immunoglobulin isotype switching.

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

What is this antibody validated in?
Anti-FANCD2 antibody [EPR2302] (ab108928) is a rabbit recombinant monoclonal antibody and is validated for use in Western Blot (WB), Flow Cytometry (Intra), Immunoprecipitation (IP), Immunohistochemistry (IHC-P), Immunocytochemistry/immunofluorescence (ICC/IF) in Human, Mouse, Rat samples.

What is the molecular weight of FANCD2?
Anti-FANCD2 [EPR2302] (ab108928) specifically detects a band for FANCD2 (UniProt: Q9BXW9) at a molecular weight of 166kDa.

Trusted by the scientific community
Anti-FANCD2 [EPR2302] (ab108928) was first used in a scientific publication in 2011 and has been cited over 40 times in peer-reviewed journals.

Specificity confirmed
The specificity of Anti-FANCD2 antibody [EPR2302] (ab108928) has been confirmed by Western blot testing in FANCD2 Knockout HAP1 cells.

Other related products
We have a range of other formats of antibody clone [EPR2302] also available for your convenience: ab108928, Alexa Fluor® 488 - ab200759, Alexa Fluor® 647 - ab200763, Carrier free - ab221932, PE - ab307174, APC - ab307175, HRP - ab307176, Alkaline Phosphatase - ab308832, Alexa Fluor® 594 - ab310563, Alexa Fluor® 555 - ab312092, Alexa Fluor® 568 - ab312572, Alexa Fluor® 750 - ab320916

Trial sizes available!
Test your antibody or perform pre-screening before committing to a larger quantity. Sold in 20µl. Discover our selection of trial-size antibodies.

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	FANCD2 also known as Fanconi anemia group D2 functions mechanically in DNA repair processes. This protein plays a role in the cellular response to DNA damage specifically interstrand crosslinks. FANCD2 has a molecular weight of approximately 164 kDa. Expression of FANCD2 occurs broadly in various tissues but reaches higher levels in cells that undergo rapid division such as hematopoietic cells.
Biological function summary	FANCD2 collaborates with multiple proteins as part of the Fanconi anemia (FA) complex to ensure genomic stability. The FA complex operates by facilitating the repair of DNA interstrand crosslinks that impede replication. FANCD2 becomes activated through monoubiquitination which serves as a signal for recruitment to DNA damage sites. It essentially functions as a coordinator that brings together important components for repair.
Pathways	FANCD2 integrates into the Fanconi anemia pathway and the homologous recombination repair pathway. These pathways are critical for maintaining DNA integrity and preventing chromosomal instability. Within these pathways interactions with proteins such as BRCA1 and BRCA2 reinforce the repair

processes. FANCD2's connection to these proteins exemplifies its role in complex mechanisms that preserve genomic fidelity.

Associated diseases and disorders

FANCD2 links significantly to Fanconi anemia a genetic disorder that causes bone marrow failure and increased cancer susceptibility. Damage or malfunction of FANCD2 can disrupt DNA repair leading to cellular dysfunction seen in this condition. Additionally there is a connection to breast cancer whereby FANCD2 interacts with BRCA2 indicating a shared pathway involved in tumor suppressor functions. These associations underline the importance of FANCD2 in disease pathology and its potential as a therapeutic target.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

12 product images




Immunocytochemistry/ Immunofluorescence - Anti-FANCD2 antibody [EPR2302] (ab108928)

FANCD2 Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-FANCD2 antibody

Immunocytochemistry/ Immunofluorescence analysis of HeLa (Human cervix adenocarcinoma epithelial cell) cells labeling FANCD2 with purified ab108928 at 1/500 dilution (0.4 µg/ml). Cells were fixed in 4% Paraformaldehyde and permeabilized with 0.1% tritonX-100. Cells were counterstained with ab195889 Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa

Fluor® 594) at 1/200 (2.5 µg/ml) dilution. Goat anti rabbit IgG (Alexa Fluor® 488, ab150077) was used as the secondary antibody at 1/1000 (2 µg/ml) dilution. DAPI (blue) was used as nuclear counterstain. PBS instead of the primary antibody was used as the secondary antibody only control.




This image is courtesy of an anonymous customer review

Immunocytochemistry/ Immunofluorescence - Anti-FANCD2 antibody [EPR2302] (ab108928)

FANCD2 Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-FANCD2 antibody


ab108928 (unpurified) staining FANCD2 in human U2OS osteosarcoma cells by ICC/IF (Immunocytochemistry/immunofluorescence). Cells were fixed with paraformaldehyde, permeabilized with 0.5% Triton X-100 and blocked with 10% goat serum for 1 hour at 20°C. Samples were incubated with primary antibody (1/300) for 18 hours at 4°C. An Alexa Fluor® 488-conjugated goat anti-rabbit IgG monoclonal (1/1000) was used as the secondary antibody.



Immunocytochemistry/ Immunofluorescence - Anti-FANCD2 antibody [EPR2302] (ab108928)

FANCD2 Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-FANCD2 antibody


ab108928 (unpurified) staining FANCD2 in wild-type HAP1 cells (top panel) and FANCD2 knockout HAP1 cells (bottom panel). The cells were fixed with 4% formaldehyde (10 minutes), permeabilized with 0.1% Triton X-100 for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween for 1 hour. The cells were then incubated with ab108928 at 1/250 dilution and ab195889 at 1/250 dilution (shown in pseudo colour red) overnight at +4°C, followed by a further incubation at room temperature for 1 hour with a goat secondary antibody to Rabbit IgG (Alexa Fluor® 488) (ab150081) at 2 µg/ml (shown in green). Nuclear DNA was labelled in blue with DAPI.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-FANCD2 antibody [EPR2302] (ab108928)

FANCD2 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-FANCD2 antibody

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of Human tonsil tissue sections labeling FANCD2 with purified ab108928 at 1/50 dilution (3.6 µg/ml). Perform heat mediated antigen retrieval using ab93684 (Tris/EDTA buffer, pH 9.0). ImmunoHistoProbe one step HRP Polymer (ready to use) was used as the secondary antibody. Negative control: PBS instead of the primary antibody. Hematoxylin was used as a counterstain.




Flow Cytometry (Intracellular) - Anti-FANCD2 antibody [EPR2302] (ab108928)

FANCD2 Flow Cytometry (Intracellular) staining using rabbit Anti-FANCD2 antibody

Intracellular Flow Cytometry analysis of Jurkat (human acute T cell leukemia) cells labeling FANCD2 (red) with ab108928 (purified) at a 1/200 dilution. Cells were fixed with 4% paraformaldehyde and permeabilized with 90% methanol. A goat anti-rabbit IgG (Alexa Fluorr®


488) (ab150077) was used as the secondary antibody at a 1/2000 dilution. Black - Rabbit monoclonal IgG (ab172730). Blue (unlabeled control) - Cells without incubation with primary and secondary antibodies.



Immunocytochemistry/ Immunofluorescence - Anti-FANCD2 antibody [EPR2302] (ab108928)

FANCD2 Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-FANCD2 antibody


HeLa (human epithelial cell line from cervix adenocarcinoma) cells stained for FANCD2 (green) using ab108928 (unpurified) (1/250 dilution) in ICC/IF.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-FANCD2 antibody [EPR2302] (ab108928)

FANCD2 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-FANCD2 antibody


Immunohistochemical staining of paraffin-embedded human tonsil tissue using ab108928 (unpurified) at a dilution of 1/100.
Perform heat mediated antigen retrieval before commencing with IHC staining protocol.



Immunoprecipitation - Anti-FANCD2 antibody [EPR2302] (ab108928)

ab108928 (purified) at 1/20 dilution (1ug) immunoprecipitating FANCD2 in HeLa whole cell lysates.
Lane 1: HeLa (Human cervix adenocarcinoma epithelial cell) whole cell lysates 10ug
Lane 2 (+): ab108928 & HeLa whole cell lysates
Lane 3 (-): Rabbit monoclonal IgG (ab172730) instead of ab108928 in HeLa whole cell lysates
For western blotting, VeriBlot for IP Detection Reagent (HRP) (ab131366) was used at 1/1000 dilution.
Blocking and diluting buffer: 5% NFDM/TBST.
All lanes:
Immunoprecipitation - Anti-FANCD2 antibody [EPR2302] (ab108928)

Predicted band size: 166 kDa



Western blot - Anti-FANCD2 antibody [EPR2302] (ab108928)

All lanes:
Western blot - Anti-FANCD2 antibody [EPR2302] (ab108928) at 1/1000 dilution

Lane 1:
HeLa (Human cervix adenocarcinoma epithelial cell) whole cell lysates at 20 µg

Lane 2:
MCF7 (Human breast adenocarcinoma epithelial cell) whole cell lysates at 20 µg

Lane 3:
C6 (Rat glial tumor glial cell) whole cell lysates at 20 µg

Lane 4:

RAW264.7 (Mouse Abelson murine leukemia virus-induced tumor macrophage) whole cell lysates at 20 µg


Lane 5:
NIH/3T3 (Mouse embryonic fibroblast) whole cell lysates at 20 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/20000 dilution

Predicted band size: 166 kDa

Observed band size: 165 kDa




Western blot - Anti-FANCD2 antibody [EPR2302] (ab108928)

FANCD2 Western blot staining using rabbit Anti-FANCD2 antibody

Lane 1: Wild-type HAP1 cell lysate (20 µg)
Lane 2: FANCD2 knockout HAP1 cell lysate (20 µg)
Lane 3: HeLa cell lysate (20 µg)
Lane 4: HEK293 cell lysate (20 µg)
Lanes 1 - 4: Merged signal (red and green). Green - ab108928 (unpurified) observed at 165 kDa. Red - loading control, ab8245, observed at 37 kDa.
ab108928 was shown to specifically react with FANCD2 when FANCD2 knockout samples were used. Wild-type and FANCD2 knockout samples were subjected to SDS-PAGE. ab108928 and ab8245 (loading control to GAPDH) were diluted 1/1000 and 1/2000 respectively and incubated overnight at 4°C. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preadsorbed (ab216773) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preadsorbed (ab216776) secondary antibodies at 1/10 000 dilution for 1 h at room temperature before imaging.

All lanes:
Western blot - Anti-FANCD2 antibody [EPR2302] (ab108928)

Predicted band size: 166 kDa



Western blot - Anti-FANCD2 antibody [EPR2302] (ab108928)

FANCD2 Western blot staining using rabbit Anti-FANCD2 antibody

Lanes 1- 2: Merged signal (red and green). Green - ab108928 observed at 166 kDa. Red - Anti-alpha Tubulin antibody [DM1A] - Loading Control (ab7291) observed at 50 kDa.
ab108928 was shown to react with FANCD2 in wild-type HeLa cells in western blot. Loss of signal was observed when knockout cell line ab261743 (knockout cell lysate ab257173) was used. Wild-type HeLa and FANCD2 knockout HeLa cell lysates were subjected to SDS-PAGE. Membrane was blocked for 1 hour at room temperature in 0.1% TBST with 3% non-fat dried milk. ab108928 and Anti-alpha Tubulin antibody [DM1A] - Loading Control (ab7291) overnight at 4°C at a 1 in 1000 dilution and a 1 in 20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye®800CW) preadsorbed (ab216773) and Goat anti-Mouse IgG H&L (IRDye®680RD) preadsorbed (ab216776) secondary antibodies at 1 in 20000 dilution for 1 hour at room temperature before imaging.

All lanes:
Western blot - Anti-FANCD2 antibody [EPR2302] (ab108928) at 1/1000 dilution

Lane 1:

Wild-type HeLa cell lysate at 20 µg


Lane 2:
FANCD2 knockout HeLa cell lysate at 20 µg

Lane 2:
Western blot - Human FANCD2 knockout HeLa cell line (ab261743)

Performed under reducing conditions.

Predicted band size: 166 kDa

Observed band size: 166 kDa



Western blot - Anti-FANCD2 antibody [EPR2302] (ab108928)

FANCD2 Western blot staining using rabbit Anti-FANCD2 antibody

Western blot: Anti-FANCD2 antibody [EPR2302] (ab108928) staining at 1/1000 dilution, shown in green; Mouse anti-Alpha Tubulin [DM1A] (ab7291) loading control staining at 1/20000 dilution, shown in magenta. In Western blot, ab108928 was shown to bind specifically to FANCD2. A band was observed at 160 kDa in wild-type HCT 116 cell lysates with no signal observed at this size in FANCD2 knockout cell line. To generate this image, wild-type and FANCD2 knockout HCT 116 cell lysates were analysed. First, samples were run on an SDS-PAGE gel then transferred onto a nitrocellulose membrane. Membranes were blocked in 3 % milk in TBS-0.1 % Tween® 20 (TBS-T) before incubation with primary antibodies overnight at 4 °C. Blots were washed four times in TBS-T, incubated with secondary antibodies for 1 h at room temperature, washed again four times then imaged. Secondary antibodies used were Goat anti-Rabbit IgG H&L 800CW and Goat anti-Mouse IgG H&L 680RD at 1/20000 dilution.
All lanes:
Western blot - Anti-FANCD2 antibody [EPR2302] (ab108928) at 1/1000 dilution

Lane 1:
Wild-type HCT 116 cell lysate at 20 µg

Lane 2:
FANCD2 knockout HCT 116 cell lysate at 20 µg

Lane 3:
Wild-type HeLa ab255448 cell lysate at 20 µg

Lane 4:
FANCD2 knockout HeLa ab261743 cell lysate at 20 µg

Secondary

Lanes 1 - 4:
Goat anti-Rabbit IgG H&L 800CW at 1/20000 dilution

Lanes 1 - 4:
Goat anti-Mouse IgG H&L 680RD at 1/20000 dilution

Performed under reducing conditions.

Observed band size: 160 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.