abcam

ab109122

Anti-SATB1 antibody [EPR3951]

Anti-SATB1 antibody [EPR3951] (ab109122) is a rabbit monoclonal antibody detecting SATB1 in **Western Blot, Flow Cytometry, IP, IHC-P, ICC/IF**. Suitable for **Human, Mouse, Rat**.

- KO validated for confirmed specificity
- Biophysical QC for unrivalled batch-batch consistency
- Over 20 publications

Recombinant | RabMAb | KO Validated | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR3951
Purification technique	Affinity purification Protein A
Concentration	0.119 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

IHC-P

Tested

Species	Human
Dilution info	1/500
Notes	heat up to 98 degrees C, below boiling, and then let cool for 10-20 min. Perform heat-mediated antigen retrieval before commencing with IHC staining protocol.

Species	Rat
Dilution info	1/500
Notes	heat up to 98 degrees C, below boiling, and then let cool for 10-20 min. Perform heat-mediated antigen retrieval before commencing with IHC staining protocol.

Species	Mouse
Dilution info	1/500
Notes	heat up to 98 degrees C, below boiling, and then let cool for 10-20 min. Perform heat-mediated antigen retrieval before commencing with IHC staining protocol.

IP

Tested

Species	Rat
Dilution info	1/20
Notes	-

Expected

Species	Human, Mouse
Dilution info	Use at an assay dependent concentration.
Notes	-

Flow Cyt

Tested

Species	Human
Dilution info	1/20

Notes	-

Expected

Species	Rat, Mouse
Dilution info	Use at an assay dependent concentration.
Notes	-

WB

Tested

Species	Human
Dilution info	1/1000
Notes	-

Species	Mouse
Dilution info	1/1000
Notes	-

Species	Rat
Dilution info	1/1000
Notes	-

ICC/IF

Tested

Species	Human
Dilution info	1/50
Notes	-

Expected

Species	Rat, Mouse
Dilution info	Use at an assay dependent concentration.

Notes	-

Target data

See full target information SATB1

Function	Crucial silencing factor contributing to the initiation of X inactivation mediated by Xist RNA that occurs during embryogenesis and in lymphoma (By similarity). Binds to DNA at special AT-rich sequences, the consensus SATB1-binding sequence (CSBS), at nuclear matrix- or scaffold-associated regions. Thought to recognize the sugar-phosphate structure of double-stranded DNA. Transcriptional repressor controlling nuclear and viral gene expression in a phosphorylated and acetylated status-dependent manner, by binding to matrix attachment regions (MARs) of DNA and inducing a local chromatin-loop remodeling. Acts as a docking site for several chromatin remodeling enzymes (e.g. PML at the MHC-I locus) and also by recruiting corepressors (HDACs) or coactivators (HATs) directly to promoters and enhancers. Modulates genes that are essential in the maturation of the immune T-cell CD8SP from thymocytes. Required for the switching of fetal globin species, and beta- and gamma-globin genes regulation during erythroid differentiation. Plays a role in chromatin organization and nuclear architecture during apoptosis. Interacts with the unique region (UR) of cytomegalovirus (CMV). Alu-like motifs and SATB1-binding sites provide a unique chromatin context which seems preferentially targeted by the HIV-1 integration machinery. Moreover, HIV-1 Tat may overcome SATB1-mediated repression of IL2 and IL2RA (interleukin) in T-cells by binding to the same domain as HDAC1. Delineates specific epigenetic modifications at target gene loci, directly up-regulating metastasis-associated genes while down-regulating tumor-suppressor genes. Reprograms chromatin organization and the transcription profiles of breast tumors to promote growth and metastasis. Promotes neuronal differentiation of neural stem/progenitor cells in the adult subventricular zone, possibly by positively regulating the expression of NEUROD1 (By similarity).

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

What is this antibody validated in?
Anti-SATB1 antibody [EPR3951] (ab109122) is a rabbit recombinant monoclonal antibody and is validated for use in Western Blot (WB), Flow Cytometry (Flow Cyt), Immunoprecipitation (IP), Immunohistochemistry (IHC-P), Immunocytochemistry/immunofluorescence (ICC/IF) in Human, Mouse, Rat samples.

What is the molecular weight of SATB1?
Anti-SATB1 [EPR3951] (ab109122) specifically detects a band for SATB1 (UniProt: Q01826) at a molecular weight of 86kDa.

Trusted by the scientific community
Anti-SATB1 [EPR3951] (ab109122) was first used in a scientific publication in 2011 and has been cited over 20 times in peer-reviewed journals.

Specificity confirmed
The specificity of Anti-SATB1 antibody [EPR3951] (ab109122) has been confirmed by Western blot testing in SATB1 Knockout HAP1 cells.

Other related products
We have a range of other formats of antibody clone [EPR3951] also available for your convenience: ab109122, HRP - ab203964, Carrier free - ab239944

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Trial sizes available!
Test your antibody or perform pre-screening before committing to a larger quantity. Sold in 20µl. Discover our selection of trial-size antibodies.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	SATB1 or Special AT-rich Sequence Binding Protein 1 is a nuclear matrix-associated protein with a mass of approximately 85 kDa. It modifies chromatin structure by organizing the density of DNA through loop formation which regulates gene expression. SATB1 is expressed in thymocytes breast tissue and various other cell types and plays a significant role in cellular differentiation and development. It

	binds selectively to base-unpairing regions allowing it to influence expression patterns of multiple genes.
Biological function summary	SATB1 acts as a genome organizer and transcription regulator by linking chromatin remodeling complexes to specific genomic loci. It participates in chromatin loop domain organization influencing higher-order chromatin structure and gene silencing or activation. SATB1 does not usually work alone but rather associates with corepressors such as HDAC1 and coactivators like PCAF in regulating transcription. Its functions have been noted as particularly important in T cell development where it assists in the regulation of T cell receptor genes.
Pathways	SATB1 integrates into the Wnt signaling pathway and influences the Notch signaling pathway. The protein interacts with B-catenin in the Wnt pathway aiding in sustaining key developmental processes. In the Notch signaling pathway SATB1 influences cell fate decisions impacting differentiation and proliferation through its interaction with RBP-Jκ and other transcription modulators. MUTYH and LRP6 are among the proteins within these pathways influenced by SATB1's regulatory capacity.
Associated diseases and disorders	SATB1 links to various cancer types including breast cancer and colorectal cancer. Overexpression of SATB1 can drive tumor progression and metastasis by altering gene expression landscapes. It's known to affect pathways regulated by proteins like BRCA1 in breast cancer contributing to disease pathogenicity. In immune disorders such as autoimmune diseases SATB1 interacts with STAT5 impacting cytokine signaling pathways and thereby influencing immunological tolerance and efficacy.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.


9 product images



Immunocytochemistry/ Immunofluorescence - Anti-SATB1 antibody [EPR3951] (ab109122)

SATB1 Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-SATB1 antibody


Immunocytochemistry analysis of Jurkat (Human T cell leukemia T lymphocyte) cells labeling SATB1 with Purified ab109122 at 1:50 dilution (2.5 µg/ml). Cells were fixed in 4% Paraformaldehyde and permeabilized with 0.1% tritonX-100. Cells were counterstained with ab195889 Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor® 594) 1:200 (2.5 µg/ml). Goat anti rabbit IgG (Alexa Fluor® 488, ab150077) was used as the secondary antibody at 1:1000 (2 µg/ml) dilution. DAPI (blue) was used as nuclear counterstain. PBS instead of the primary antibody was used as the secondary antibody only control.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-SATB1 antibody [EPR3951] (ab109122)

SATB1 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-SATB1 antibody


Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human lung carcinoma tissue sections labeling SATB1 with Purified ab109122 at 1:500 dilution (0.246 µg/ml). Heat mediated antigen retrieval was performed using Heat mediated antigen retrieval using Bond™ Epitope Retrieval Solution 2 (pH 9.0) . Tissue was counterstained with Hematoxylin. Rabbit specific IHC polymer detection kit HRP/DAB (ab209101) secondary antibody was used. PBS instead of the primary antibody was used as the negative control.



Flow Cytometry - Anti-SATB1 antibody [EPR3951] (ab109122)

SATB1 Flow Cytometry staining using rabbit Anti-SATB1 antibody

Flow Cytometry analysis of Jurkat (Human T cell leukemia T lymphocyte) cells labelling SATB1 with Purified ab109122 at 1:20 dilution (10 µg/ml) (Red). Cells were fixed with 4% Paraformaldehyde and permeabilised with 90% Methanol. A Goat anti rabbit IgG (Alexa Fluor® 488, ab150077) secondary antibody was used at 1:2000. Isotype control - Rabbit monoclonal IgG (Black). Unlabelled control - Cell without incubation with primary antibody and secondary antibody (Blue).




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-SATB1 antibody [EPR3951] (ab109122)

SATB1 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-SATB1 antibody

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of rat liver tissue sections labeling SATB1 with Purified ab109122 at 1:500 dilution (0.246 µg/ml). Heat mediated antigen retrieval was performed using Heat mediated antigen retrieval using Bond™


Epitope Retrieval Solution 2 (pH 9.0) . Tissue was counterstained with Hematoxylin. Rabbit specific IHC polymer detection kit HRP/DAB (ab209101) secondary antibody was used. PBS instead of the primary antibody was used as the negative control.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-SATB1 antibody [EPR3951] (ab109122)

SATB1 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-SATB1 antibody


Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of mouse liver tissue sections labeling SATB1 with Purified ab109122 at 1:500 dilution (0.246 µg/ml). Heat mediated antigen retrieval was performed using Heat mediated antigen retrieval using Bond™ Epitope Retrieval Solution 2 (pH 9.0) . Tissue was counterstained with Hematoxylin. Rabbit specific IHC polymer detection kit HRP/DAB (ab209101) secondary antibody was used. PBS instead of the primary antibody was used as the negative control.



Immunoprecipitation - Anti-SATB1 antibody [EPR3951] (ab109122)

Purified ab109122 at 1:20 dilution (0.6µg) immunoprecipitating SATB1 in Rat thymus lysate.
Lane 1 (input): Rat thymus lysate 10 µg
Lane 2 (+): ab109122 + Rat thymus lysate.
Lane 3 (-): Rabbit monoclonal IgG (ab172730) instead of ab109122 in Rat thymus lysate.
VeriBlot for IP Detection Reagent (HRP)(ab131366) (1:5000 dilution) was used for Western blotting.
Blocking Buffer and concentration: 5% NFDM/TBST.
Diluting buffer and concentration: 5% NFDM/TBST.
Observed band size: kDa
All lanes:
Immunoprecipitation - Anti-SATB1 antibody [EPR3951] (ab109122)

Predicted band size: 85 kDa



Western blot - Anti-SATB1 antibody [EPR3951] (ab109122)


SATB1 Western blot staining using rabbit Anti-SATB1 antibody

Lane 1: Wild-type HAP1 whole cell lysate (20 µg)
Lane 2: SATB1 knockout HAP1 whole cell lysate (20 µg)
Lane 3: Jurkat whole cell lysate (20 µg)
Lane 4: THP1 whole cell lysate (20 µg)
Lanes 1 - 4: Merged signal (red and green). Green - ab109122 observed at 100 kDa. Red - loading control, ab9484, observed at 37 kDa.

ab109122 was shown to specifically react with SATB1 in wild-type HAP1 cells as signal was lost in SATB1 knockout cells. Wild-type and SATB1 knockout samples were subjected to SDS-PAGE. ab109122 and ab9484 (Mouse anti-GAPDH loading control) were incubated overnight at 4°C at 1/1000 dilution and 1/20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preabsorbed ab216773 and Goat anti-Mouse IgG H&L (IRDye® 680RD) preabsorbed ab216776 secondary antibodies at 1/20000 dilution for 1 hour at room temperature before imaging.

All lanes:
Western blot - Anti-SATB1 antibody [EPR3951] (ab109122)

Predicted band size: 85 kDa



Western blot - Anti-SATB1 antibody [EPR3951] (ab109122)

All lanes:
Western blot - Anti-SATB1 antibody [EPR3951] (ab109122) at 1/1000 dilution

Lane 1:
LNCaP (Human prostate carcinoma epithelial cell) whole cell lysate at 20 µg

Lane 2:
THP-1 (Human monocytic leukemia monocyte) whole cell lysate at 20 µg

Lane 3:
Mouse liver lysate at 20 µg


Lane 4:
Mouse thymus lysate at 20 µg

Lane 5:
Rat thymus lysate at 20 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/20000 dilution

Predicted band size: 85 kDa



Western blot - Anti-SATB1 antibody [EPR3951] (ab109122)

All lanes:
Western blot - Anti-SATB1 antibody [EPR3951] (ab109122) at 1/1000 dilution

Lane 1:
Jurkat (Human T cell leukemia T lymphocyte) nuclear extract lysate at 20 µg

Lane 2:
Jurkat (Human T cell leukemia T lymphocyte) without nuclear extract lysate at 20 µg

Lane 3:
Jurkat (Human T cell leukemia T lymphocyte) whole cell lysate at 20 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/20000 dilution

Predicted band size: 85 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.