abcam

ab109367

Anti-Peroxiredoxin 2/PRP antibody [EPR5154]

Anti-Peroxiredoxin 2/PRP antibody [EPR5154] (ab109367) is a rabbit monoclonal antibody detecting Peroxiredoxin 2/PRP in **Western Blot, Flow Cytometry (Intra), IHC-P, ICC/IF**. Suitable for **Human, Mouse, Rat**.

- KO validated for confirmed specificity
- Biophysical QC for unrivalled batch-batch consistency
- Over 40 publications

Recombinant | RabMAb | KO Validated | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 50% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR5154
Purification technique	Affinity purification Protein A
Concentration	0.848 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

IHC-P

Tested

Species	Human
Dilution info	1/500 - 1/2000
Notes	Perform heat-mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.

Expected

Species	Mouse, Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

Predicted

Species	Pig
Dilution info	-
Notes	-

WB

Tested

Species	Mouse
Dilution info	1/1000 - 1/10000
Notes	-

Species	Human
Dilution info	1/1000 - 1/10000
Notes	-

Species	Rat
Dilution info	1/1000 - 1/10000
Notes	-

Predicted

Species	Pig

Dilution info	-
Notes	-

ICC/IF

Tested

Species	Human
Dilution info	1/100 - 1/500
Notes	-

Expected

Species	Mouse, Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

Predicted

Species	Pig
Dilution info	-
Notes	-

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	1/200 - 1/21200
Notes	ab172730 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.

Expected

Species	Mouse, Rat
Dilution info	Use at an assay dependent concentration.

Notes	-

Predicted

Species	Pig
Dilution info	-
Notes	-

Target data

See full target information PRDX2

Function	Thiol-specific peroxidase that catalyzes the reduction of hydrogen peroxide and organic hydroperoxides to water and alcohols, respectively. Plays a role in cell protection against oxidative stress by detoxifying peroxides and as sensor of hydrogen peroxide-mediated signaling events. Might participate in the signaling cascades of growth factors and tumor necrosis factor-alpha by regulating the intracellular concentrations of H(2)O(2).

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Stable for 12 months at -20°C

Notes

What is this antibody validated in?
Anti-Peroxiredoxin 2/PRP antibody [EPR5154] (ab109367) is a rabbit recombinant monoclonal antibody and is validated for use in Western Blot (WB), Flow Cytometry (Intra), Immunohistochemistry (IHC-P), Immunocytochemistry/immunofluorescence (ICC/IF) in Human, Mouse, Rat samples.

What is the molecular weight of Peroxiredoxin 2/PRP?
Anti-Peroxiredoxin 2/PRP [EPR5154] (ab109367) specifically detects a band for Peroxiredoxin 2/PRP (UniProt: P32119) at a molecular weight of 22kDa.

Trusted by the scientific community
Anti-Peroxiredoxin 2/PRP [EPR5154] (ab109367) was first used in a scientific publication in 2011 and has been cited over 40 times in peer-reviewed journals.

Specificity confirmed
The specificity of Anti-Peroxiredoxin 2/PRP antibody [EPR5154] (ab109367) has been confirmed by Western blot testing in PRDX2 Knockout HAP1 cells.

Other related products
We have a range of other formats of antibody clone [EPR5154] also available for your convenience: ab109367, Alexa Fluor® 647 - ab197041, HRP - ab197042, Alexa Fluor® 488 - ab197536, Carrier free - ab227988

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Trial sizes available!
Test your antibody or perform pre-screening before committing to a larger quantity. Sold in 20µl. Discover our selection of trial-size antibodies.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	Peroxiredoxin 2 (also known as PRP or PRDX2) is an antioxidant enzyme that plays an important role in reducing peroxides and protecting cells from oxidative damage. The target consisting of approximately 22kDa mass is expressed heavily in erythrocytes but can also be found in other tissues like the heart and liver. PRDX2 belongs to the peroxiredoxin family and its redox activity contributes significantly to cellular homeostasis and defense against oxidative stress.
Biological function summary	Peroxiredoxins like PRDX2 function by breaking down hydrogen peroxide and organic hydroperoxides safeguarding cells from oxidative harm. PRDX2 often forms homodimers or higher-order oligomers which influence its catalytic efficiency and chaperone activity. Within the cell it not only acts independently but also associates with other cellular components reflecting its participation in important cellular processes including cell proliferation and apoptosis.

Pathways	PRDX2 engages significantly within antioxidant defense systems specifically the thioredoxin pathway. It interacts with thioredoxin reductase and thioredoxin to facilitate the reduction of peroxides maintaining the cell's redox balance. Furthermore PRDX2 intersects with cellular signaling pathways associated with inflammation and the regulation of cell death where proteins like NRF2 and KEAP1 play important roles in managing oxidant-inducible gene expression.
Associated diseases and disorders	Disruptions in PRDX2 function connect to conditions like cancer and cardiovascular diseases. In cancer altered peroxiredoxin 2 expression correlates to tumor progression and resistance to chemotherapy implicating its interaction with proteins like NF-κB in enhancing cell survival. In cardiovascular diseases oxidative stress mediated by PRDX2 imbalance can lead to myocardial infarction and heart failure linking it to proteins such as SOD2 involved in mitochondrial defense.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.


15 product images



Immunocytochemistry/ Immunofluorescence - Anti-Peroxiredoxin 2/PRP antibody [EPR5154] (ab109367)

Peroxiredoxin 2/PRP Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-Peroxiredoxin 2/PRP antibody

Immunofluorescent analysis of 100% methanol-fixed 0.1% Triton X-100 permeabilized Hap1 WT and Hap1-PRDX2 KO cells labelling Peroxiredoxin 2 with ab109367 at at 5 µg/ml concentration followed by ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed antibody at 1/1000 2 µg/ml dilution (Green). Image showing cytoplasmic staining in Hap1 WT cell line. ab7291 Anti-alpha Tubulin antibody [DM1A] was used to counterstain tubulin at 1/1000 dilution (Magenta). The nuclear counterstain was DAPI (Blue).



Flow Cytometry (Intracellular) - Anti-Peroxiredoxin 2/PRP antibody [EPR5154] (ab109367)


Peroxiredoxin 2/PRP Flow Cytometry (Intracellular) staining using rabbit Anti-Peroxiredoxin 2/PRP antibody

Flow cytometry overlay histogram showing wild-type HAP1 (green line) and PRDX2 knockout HAP1 stained with ab109367 (magenta line). The cells were fixed with 80% methanol (5 min) and then permeabilised with 0.1% PBS-Triton X-100 for 15 min. The cells were then incubated in 1x PBS containing 10% normal goat serum to block non-specific protein-protein interaction followed by the antibody (ab109367) (1x 10^6 in 100µl at 0.04 µg/ml (1/21200 dilution)) for 30min at 22°C.

The secondary antibody Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed was incubated at 1/4000 dilution for 30min at 22°C.

Isotype control antibody was Recombinant Rabbit IgG, monoclonal [EPR25A] - Isotype Control in HAP1 WT cells (black line) and HAP1-PRDX2 KO cells (grey line), used at the same concentration and conditions as the primary antibody.


Acquisition of >5000 events were collected using a 50 mW Blue laser (488nm) and 525/40 bandpass filter.



Flow Cytometry (Intracellular) - Anti-Peroxiredoxin 2/PRP antibody [EPR5154] (ab109367)

Peroxiredoxin 2/PRP Flow Cytometry (Intracellular) staining using rabbit Anti-Peroxiredoxin 2/PRP antibody


Overlay histogram showing HeLa cells fixed in 80% methanol and stained with purified ab109367 at a dilution of 1/200 (red line). The secondary antibody used was Alexa Fluorr® 488 goat anti-rabbit at a dilution of 1/500. Rabbit monoclonal IgG was used as an isotype control (black line) and cells incubated in the absence of both primary and secondary antibody were used as a negative control (blue line).



Flow Cytometry (Intracellular) - Anti-Peroxiredoxin 2/PRP antibody [EPR5154] (ab109367)

Peroxiredoxin 2/PRP Flow Cytometry (Intracellular) staining using rabbit Anti-Peroxiredoxin 2/PRP antibody


Overlay histogram showing HeLa cells stained with unpurifiedab109367 (red line). The cells were fixed with 80% methanol (5 min) and then permeabilized with 0.1% PBS-Tween for 20 min. The cells were then incubated in 1x PBS / 10% normal goat serum / 0.3M glycine to block non-specific protein-protein interactions followed by the antibody (ab109367, 1/1000 dilution) for 30 min at 22°C. The secondary antibody used was Alexa Fluor® 488 goat anti-rabbit IgG (H+L) (ab150077) at 1/2000 dilution for 30 min at 22°C. Isotype control antibody (black line) was rabbit IgG (monoclonal) (0.1µg/1x10^6 cells) used under the same conditions. Unlabelled sample (blue line) was also used as a control. Acquisition of >5,000 events were collected using a 20mW Argon ion laser (488nm) and 525/30 bandpass filter.



Immunocytochemistry/ Immunofluorescence - Anti-Peroxiredoxin 2/PRP antibody [EPR5154] (ab109367)

Peroxiredoxin 2/PRP Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-Peroxiredoxin 2/PRP antibody


Immunofluorescent staining of Peroxiredoxin 2/PRP in HeLa cells using unpurified ab109367 at 1/250 dilution.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Peroxiredoxin 2/PRP antibody [EPR5154] (ab109367)

Peroxiredoxin 2/PRP Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-Peroxiredoxin 2/PRP antibody


Immunohistochemical analysis of Peroxiredoxin 2/PRP in paraffin-embedded Human stomach tissue using unpurified ab109367 at 1/1000 dilution.
Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Peroxiredoxin 2/PRP antibody [EPR5154] (ab109367)

Peroxiredoxin 2/PRP Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-Peroxiredoxin 2/PRP antibody

Immunohistochemical analysis of Peroxiredoxin 2/PRP in paraffin-embedded Human prostatic hyperplasia tissue using unpurified ab109367 at 1/1000 dilution.
Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Peroxiredoxin 2/PRP antibody [EPR5154] (ab109367)

Peroxiredoxin 2/PRP Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-Peroxiredoxin 2/PRP antibody

Immunohistochemical staining of paraffin embedded human endometrial carcinoma with purified ab109367 at a working dilution of 1/500. The secondary antibody used is ab97051, a goat anti-rabbit IgG (H&L) at a dilution of 1/500. The sample is counter-stained with hematoxylin. Antigen

retrieval was perfomed using Tris-EDTA buffer, pH 9.0. PBS was used instead of the primary antibody as the negative control, and is shown in the inset.



Immunocytochemistry/ Immunofluorescence - Anti-Peroxiredoxin 2/PRP antibody [EPR5154] (ab109367)

Peroxiredoxin 2/PRP Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-Peroxiredoxin 2/PRP antibody

Immunofluorescence staining of HeLa cells with purified ab109367 at a working dilution of 1/100, counter-stained with DAPI. The secondary antibody was Alexa Fluor® 488 goat anti-rabbit (ab150077), used at a dilution of 1/1000. ab7291, a mouse anti-tubulin antibody (1/1000), was used to stain tubulin along with ab150120 (Alexa Fluor® 594 goat anti-mouse, 1/1000), shown in the top right hand panel. The cells were fixed in 100% methanol and permeabilized using 0.1% Triton X 100. The negative controls are shown in bottom middle and right hand panels - for negative control 1, purified ab109367 was used at a dilution of 1/500 followed by an Alexa Fluor® 594 goat anti-mouse antibody (ab150120) at a dilution of 1/500. For negative control 2, ab7291 (mouse anti-tubulin) was used at a dilution of 1/500 followed by an Alexa Fluor® 488 goat anti-rabbit antibody (ab150077) at a dilution of 1/400.





Image courtesy of a customer review submitted by Dr. Kirk McManus, Univ. of Manitoba/Cancer Care MICB, Canada

Immunocytochemistry/ Immunofluorescence - Anti-Peroxiredoxin 2/PRP antibody [EPR5154] (ab109367)

Peroxiredoxin 2/PRP Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-Peroxiredoxin 2/PRP antibody

Unpurified ab109367 (1/500) staining Peroxiredoxin 2/PRP in asynchronous HeLa cells (green). Cells were fixed in methanol and counterstained with DAPI in order to highlight the nucleus (red). For further experimental details please refer to Abreview.



Western blot - Anti-Peroxiredoxin 2/PRP antibody [EPR5154] (ab109367)

Peroxiredoxin 2/PRP Western blot staining using rabbit Anti-Peroxiredoxin 2/PRP antibody

Blocking buffer: 5% NFDM/TBST
Dilution buffer: 5% NFDM/TBST
All lanes:
Western blot - Anti-Peroxiredoxin 2/PRP antibody [EPR5154] (ab109367) at 1/10000 dilution

Lane 1:
HEK293 whole cell lysate at 10 µg


Lane 2:
LNCaP whole cell lysate at 10 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/20000 dilution

Predicted band size: 22 kDa

Observed band size: 22 kDa



Western blot - Anti-Peroxiredoxin 2/PRP antibody [EPR5154] (ab109367)

Peroxiredoxin 2/PRP Western blot staining using rabbit Anti-Peroxiredoxin 2/PRP antibody

Lanes 1- 2: Merged signal (red and green). Green - ab109367 observed at 22 kDa. Red - Anti-GAPDH antibody [6C5] - Loading Control (ab8245) observed at 37 kDa.
ab109367 was shown to react with Peroxiredoxin 2/PRP in wild-type HEK-293T cells in western blot. Loss of signal was observed when knockout cell line ab266392 (knockout cell lysate ab257041) was used. Wild-type HEK-293T and PRDX2 knockout HEK-293T cell lysates were subjected to SDS-PAGE. Membrane was blocked for 1 hour at room temperature in 0.1% TBST with 3% non-fat dried milk. ab109367 and Anti-GAPDH antibody [6C5] - Loading Control (ab8245) overnight at 4°C at a 1 in 1000 dilution and a 1 in 20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye®800CW) preadsorbed (ab216773) and Goat anti-Mouse IgG H&L (IRDye®680RD) preadsorbed (ab216776) secondary antibodies at 1 in 20000 dilution for 1 hour at room temperature before imaging.

All lanes:
Western blot - Anti-Peroxiredoxin 2/PRP antibody [EPR5154] (ab109367) at 1/1000 dilution

Lane 1:
Wild-type HEK-293T cell lysate at 20 µg


Lane 2:
PRDX2 knockout HEK-293T cell lysate at 20 µg

Lane 2:
Western blot - Human PRDX2 (Peroxiredoxin 2/PRP) knockout HEK-293T cell line (ab266392)

Performed under reducing conditions.

Predicted band size: 22 kDa

Observed band size: 22 kDa



Western blot - Anti-Peroxiredoxin 2/PRP antibody [EPR5154] (ab109367)


Peroxiredoxin 2/PRP Western blot staining using rabbit Anti-Peroxiredoxin 2/PRP antibody

Lane 1: Wild-type HAP1 cell lysate (20 µg)
Lane 2: Peroxiredoxin 2/PRP knockout HAP1 cell lysate (20 µg)
Lane 3: HeLa cell lysate (20 µg)
Lane 4: LnCaP cell lysate (20 µg)
Lanes 1 - 4: Merged signal (red and green). Green - ab109367 observed at 24 kDa. Red - loading control, ab8245, observed at 37 kDa.

ab109367 was shown to specifically react with Peroxiredoxin 2/PRP when Peroxiredoxin 2/PRP knockout samples were used. Wild-type and Peroxiredoxin 2 knockout samples were subjected to SDS-PAGE. ab109367 and ab8245 (loading control to GAPDH) were both diluted 1/1000 and incubated overnight at 4°C. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preadsorbed (ab216773) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preadsorbed (ab216776) secondary antibodies at 1/10 000 dilution for 1 h at room temperature before imaging.

All lanes:
Western blot - Anti-Peroxiredoxin 2/PRP antibody [EPR5154] (ab109367)

Predicted band size: 22 kDa



Western blot - Anti-Peroxiredoxin 2/PRP antibody [EPR5154] (ab109367)

Peroxiredoxin 2/PRP Western blot staining using rabbit Anti-Peroxiredoxin 2/PRP antibody

Blocking buffer: 5% NFDM/TBST
Dilution buffer: 5% NFDM/TBST
All lanes:
Western blot - Anti-Peroxiredoxin 2/PRP antibody [EPR5154] (ab109367) at 1/10000 dilution

Lane 1:
Mouse brain lysate at 10 µg


Lane 2:
Rat brain lysate at 10 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/20000 dilution

Predicted band size: 22 kDa

Observed band size: 22 kDa



Western blot - Anti-Peroxiredoxin 2/PRP antibody [EPR5154] (ab109367)

All lanes:
Western blot - Anti-Peroxiredoxin 2/PRP antibody [EPR5154] (ab109367) at 1/1000 dilution

Lane 1:
293T cell lysate at 10 µg

Lane 2:
HeLa cell lysate at 10 µg

Lane 3:
LnCaP cell lysate at 10 µg

Lane 4:
SH-SYSY cell lysate at 10 µg

Secondary

All lanes:
HRP labelled Goat anti-Rabbit at 1/2000 dilution

Predicted band size: 22 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.