abcam

ab11939

Anti-VEGF Receptor 2 antibody

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	Constituents: PBS
Form	Liquid
Clonality	Polyclonal
Immunogen	Recombinant Fragment Protein within Human KDR. The exact immunogen used to generate this antibody is proprietary information. Database link P35968
Carrier free	Yes
Purification technique	Affinity purification Protein A
Specificity	The polyclonal antibody will detect native and recombinant human VEGFR2/KDR in ELISA experiments and on the surface or solubilized from different human cell types.
Concentration	1 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →
Purification notes	This antibody is affinity purified from rabbit sera.

Reactivity data

Flow Cyt

Tested

Species	Human

Dilution info	1-5 µg/mL
Notes	ab171870 - Rabbit polyclonal IgG, is suitable for use as an isotype control with this antibody.

Not recommended

Species	Recombinant fragment - Human, Purified native protein - Human
Dilution info	-
Notes	-

WB

Tested

Species	Recombinant fragment - Human
Dilution info	1-5 µg/mL
Notes	-

Species	Purified native protein - Human
Dilution info	1-5 µg/mL
Notes	-

Expected

Species	Human
Dilution info	Use at an assay dependent concentration.
Notes	-

sELISA

Tested

Species	Recombinant fragment - Human
Dilution info	0.5 µg/mL
Notes	For sandwich ELISA, use this antibody as Detection at 0.5µg/ml with ab42228 as Capture.

Expected

Species	Human
Dilution info	Use at an assay dependent concentration.
Notes	-

Not recommended

Species	Purified native protein - Human
Dilution info	-
Notes	-

IHC-Fr

Tested

Species	Human
Dilution info	-
Notes	-

Not recommended

Species	Recombinant fragment - Human, Purified native protein - Human
Dilution info	-
Notes	-

ICC/IF

Tested

Species	Human
Dilution info	5 µg/mL
Notes	-

Not recommended

Species	Recombinant fragment - Human, Purified native protein - Human
Dilution info	-
Notes	-

Target data

See full target information KDR

Function

Tyrosine-protein kinase that acts as a cell-surface receptor for VEGFA, VEGFC and VEGFD. Plays an essential role in the regulation of angiogenesis, vascular development, vascular permeability, and embryonic hematopoiesis. Promotes proliferation, survival, migration and differentiation of endothelial cells. Promotes reorganization of the actin cytoskeleton. Isoforms lacking a transmembrane domain, such as isoform 2 and isoform 3, may function as decoy receptors for VEGFA, VEGFC and/or VEGFD. Isoform 2 plays an important role as negative regulator of VEGFA- and VEGFC-mediated lymphangiogenesis by limiting the amount of free VEGFA and/or VEGFC and preventing their binding to FLT4. Modulates FLT1 and FLT4 signaling by forming heterodimers. Binding of vascular growth factors to isoform 1 leads to the activation of several signaling cascades. Activation of PLCG1 leads to the production of the cellular signaling molecules diacylglycerol and inositol 1,4,5-trisphosphate and the activation of protein kinase C. Mediates activation of MAPK1/ERK2, MAPK3/ERK1 and the MAP kinase signaling pathway, as well as of the AKT1 signaling pathway. Mediates phosphorylation of PIK3R1, the regulatory subunit of phosphatidylinositol 3-kinase, reorganization of the actin cytoskeleton and activation of PTK2/FAK1. Required for VEGFA-mediated induction of NOS2 and NOS3, leading to the production of the signaling molecule nitric oxide (NO) by endothelial cells. Phosphorylates PLCG1. Promotes phosphorylation of FYN, NCK1, NOS3, PIK3R1, PTK2/FAK1 and SRC.

Storage

Shipped at conditions	Blue Ice
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	VEGF Receptor 2 also known as KDR or Flk-1 is a receptor tyrosine kinase important for mediating the effects of vascular endothelial growth factor (VEGF). This protein has a mass of approximately 150 kDa. It mostly expresses in vascular endothelial cells playing a primary role in the formation of blood vessels. The receptor can also be found in various non-endothelial tissues though in smaller amounts indicating its broad functional roles outside of the vascular system.
Biological function summary	VEGF Receptor 2 is important for angiogenesis and vascular permeability. It is part of the VEGF receptor family which also includes VEGF Receptors 1 and 3 forming a complex interaction network. Upon binding with VEGF it phosphorylates key proteins to initiate signals that regulate endothelial cell migration proliferation and survival. The receptor's function is critical for normal developmental processes and wound healing as well.
Pathways	VEGF Receptor 2 is especially important within the angiogenesis and vascular permeability pathways. It plays a central role in the VEGF signaling pathway alongside proteins like VEGF A which is the primary ligand for the receptor. The signaling through this receptor modulates various downstream proteins through phosphorylation linking it to pathways that control cellular responses essential for blood vessel formation and maintenance.
Associated diseases and disorders	VEGF Receptor 2 is significantly linked with cancer and age-related macular degeneration. In the context of cancer the overexpression or dysregulation of VEGF and its receptors including VEGF Receptor 2 often leads to enhanced tumor angiogenesis promoting tumor growth and metastasis. Additionally the receptor interacts with other proteins in pathological states such as CD309 in endothelial cell-based assays which are used for research like flow cytometry and ELISA to monitor VEGF Receptor 2 activity and its impact on diseases.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.


6 product images



Immunocytochemistry/ Immunofluorescence - Anti-VEGF Receptor 2 antibody (ab11939)

VEGF Receptor 2 Immunocytochemistry/ Immunofluorescence staining of Human dermal lymphatic endothelial cells (HDLEC) using rabbit Anti-VEGF Receptor 2 antibody


Immunofluorescence staining (green) of VEGF Receptor 2 in primary human dermal lymphatic endothelial cells (HDLEC) with ab11939 (5 µg/mL), followed by Goat anti-Mouse Alexa Fluor® 488 at 1/600 dilution. Nuclear DNA was labelled with DAPI (shown in blue).



Immunohistochemistry (Frozen sections) - Anti-VEGF Receptor 2 antibody (ab11939)

VEGF Receptor 2 Immunohistochemistry (Frozen sections) staining using rabbit Anti-VEGF Receptor 2 antibody


Immunohistochemistry (Frozen sections) analysis of human spleen tissue sections labelling VEGF Receptor 2 with ab11939.



Immunohistochemistry (Frozen sections) - Anti-VEGF Receptor 2 antibody (ab11939)

VEGF Receptor 2 Immunohistochemistry (Frozen sections) staining using rabbit Anti-VEGF Receptor 2 antibody


Immunohistochemistry (Frozen sections) analysis of human spleen tissue sections labelling VEGF Receptor 2 with ab11939.



Flow Cytometry - Anti-VEGF Receptor 2 antibody (ab11939)

VEGF Receptor 2 Flow Cytometry staining using rabbit Anti-VEGF Receptor 2 antibody


Flow Cytometry analysis of primary human dermal lymphatic endothelial cells (HDLEC) labelling VEGF Receptor 2 with ab11939.



Flow Cytometry - Anti-VEGF Receptor 2 antibody (ab11939)

VEGF Receptor 2 Flow Cytometry staining using rabbit Anti-VEGF Receptor 2 antibody

Flow Cytometry analysis of primary human umbilical vein endothelial cells (HUVEC) labelling VEGF Receptor 2 with ab11939.



Sandwich ELISA - Anti-VEGF Receptor 2 antibody (ab11939)

Standard curve for VEGF Receptor 2 (Analyte: ab54347); dilution range 1pg/ml to 1µg/ml using Capture Antibody Mouse monoclonal [mV1001.3m-h] to VEGF Receptor 2 (ab42228) at 5µg/ml and Detector Antibody Rabbit polyclonal to VEGF Receptor 2 (ab11939) at 0.5µg/ml.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.