abcam

ab119695

Anti-Cytokeratin 14 antibody [SP53]

Anti-Cytokeratin 14 antibody [SP53] (ab119695) is a rabbit monoclonal antibody detecting Cytokeratin 14 in **Western Blot, Flow Cytometry (Intra), IHC-P, IHC-Fr, ICC/IF, mIHC**. Suitable for **Human, Mouse, Rat**.

- KO validated for confirmed specificity
- Multiplex IHC validated on the Leica BOND® MAX using Opal reagents
- Biophysical QC for unrivalled batch-batch consistency
- Over 10 publications

Recombinant | RabMAb | Advanced Validation | KO Validated | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.6 Preservative: 0.1% Sodium azide Constituents: PBS, 1% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	SP53
Purification technique	Affinity purification Protein A/G
Concentration	1.17 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →
Purification notes	Purified from TCS by protein A/G.

Reactivity data

IHC-P

Tested

Species	Human
Dilution info	1/100
Notes	Perform heat-mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.

Expected

Species	Mouse, Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

Predicted

Species	Cow, Pig
Dilution info	-
Notes	-

ICC/IF

Tested

Species	Human
Dilution info	5 µg/mL
Notes	-

Expected

Species	Mouse, Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

Predicted

Species	Cow, Pig
Dilution info	-

Notes	-

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	-
Notes	ab172730 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.

Expected

Species	Mouse, Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

Predicted

Species	Cow, Pig
Dilution info	-
Notes	-

WB

Tested

Species	Human
Dilution info	1/25
Notes	-

Expected

Species	Mouse, Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

Predicted

Species	Cow, Pig
Dilution info	-
Notes	-

IHC-Fr

Tested

Species	Mouse, Rat
Dilution info	-
Notes	-

Expected

Species	Human
Dilution info	Use at an assay dependent concentration.
Notes	-

Predicted

Species	Cow, Pig
Dilution info	-
Notes	-

mIHC

Tested

Species	Human
Dilution info	-
Notes	-

Expected

Species	Mouse, Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

Predicted

Species	Cow, Pig
Dilution info	-
Notes	-

Target data

See full target information Keratin, type I cytoskeletal 14

Function	The nonhelical tail domain is involved in promoting KRT5-KRT14 filaments to self-organize into large bundles and enhances the mechanical properties involved in resilience of keratin intermediate filaments in vitro.

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

What is this antibody validated in?
Anti-Cytokeratin 14 antibody [SP53] (ab119695) is a rabbit recombinant monoclonal antibody and is validated for use in Western Blot (WB), Flow Cytometry (Intra), Immunohistochemistry (IHC-P), Immunohistochemistry (IHC-Fr), Immunocytochemistry/immunofluorescence (ICC/IF), Multiplex IHC (mIHC) in Human, Mouse, Rat samples.

What is the molecular weight of Cytokeratin 14?
Anti-Cytokeratin 14 [SP53] (ab119695) specifically detects a band for Cytokeratin 14 (UniProt: P02533) at a molecular weight of 52kDa.

Collaboration
Anti-Cytokeratin 14 [SP53] (ab119695) is a clone from the portfolio of Spring Bioscience (Roche) SP clones which have been optimised for immunohistochemistry (IHC).

Trusted by the scientific community
Anti-Cytokeratin 14 [SP53] (ab119695) was first used in a scientific publication in 2011 and has been cited over 10 times in peer-reviewed journals.

Specificity confirmed
The specificity of Anti-Cytokeratin 14 antibody [SP53] (ab119695) has been confirmed by Western blot testing in Cytokeratin 14 Knockout A431 cells.

Other related products
We have a range of other formats of antibody clone [SP53] also available for your convenience: ab119695, Carrier free - ab236439, ab279542, Biotin - ab323529

Trial sizes available!
Test your antibody or perform pre-screening before committing to a larger quantity. Sold in 20µl. Discover our selection of trial-size antibodies.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	Cytokeratin 14 also known as CK14 keratin 14 or KRT14 is a type I keratin protein with a molecular mass of approximately 50 kDa. It forms intermediate filaments in epithelial cells primarily in the basal layer of stratified epithelium and is part of the cytoskeleton. Cytokeratin 14 expression appears in tissues such as the epidermis tongue and esophagus where it contributes to the structural integrity and mechanical stability of cells. Laboratories often utilize antibodies like anti-K14 and labels such as Alexa Fluor 555 or Alexa Fluor 647 to identify cytokeratin 14 in research settings.
Biological function summary	This protein significantly contributes to the maintenance and regeneration of epithelial tissues by forming a network of filaments with keratin 5. KRT14 pairs with keratin 5 to create a keratin intermediate filament complex which plays a structural role in the resilience and elasticity of epithelial tissues. This partnership is essential for the assembly of the cytoskeleton and supports the protective barrier function of the skin and related tissues.
Pathways	Cytokeratin 14 engages in pivotal cellular processes like keratinization and wound healing. The protein is actively involved in signal transduction pathways such as the epithelial cell differentiation pathway. Within these pathways KRT14 collaborates with KRT5 and other structural proteins to regulate cell proliferation and differentiation reflecting its importance in skin health and function.
Associated diseases and disorders	Cytokeratin 14 associates with specific genetic conditions including epidermolysis bullosa simplex (EBS) and squamous cell carcinoma (SCC). In EBS mutations in the KRT14 gene lead to skin fragility. In SCC altered expression of KRT14 and related

proteins such as keratin 5 can influence tumor progression and invasion. These associations highlight the critical role of KRT14 in both normal and pathological epithelial function.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.


8 product images



Immunocytochemistry/ Immunofluorescence - Anti-Cytokeratin 14 antibody [SP53] (ab119695)

Cytokeratin 14 Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-Cytokeratin 14 antibody


ab119695 staining KRT14 in wild-type A431 cells (top panel) and KRT14 knockout A431 cells (bottom panel). The cells were fixed with 100% methanol (5 min) then permeabilized with 0.1% Triton X-100 for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween for 1h. The cells were then incubated with ab119695 at 5µg/ml concentration and ab7291 (Mouse monoclonal to alpha Tubulin) at 1/1000 dilution overnight at 4°C followed by a further incubation at room temperature for 1h with a goat secondary antibody to rabbit IgG (Alexa Fluor® 488) (ab150081) at 2 µg/ml (shown in green) and a goat secondary antibody to mouse IgG (Alexa Fluor® 594) (ab150120) at 2 µg/ml (shown in red). Nuclear DNA was labelled in blue with DAPI.
Image was taken with a confocal microscope (Leica-Microsystems TCS SP8).



Flow Cytometry (Intracellular) - Anti-Cytokeratin 14 antibody [SP53] (ab119695)

Cytokeratin 14 Flow Cytometry (Intracellular) staining using rabbit Anti-Cytokeratin 14 antibody


Flow cytometry analysis of A431 (human epidermoid carcinoma) labeling Cytokeratin 14 with purified ab119695 (red). Cells were fixed with 4% paraformaldehyde and permeabilised with 90% methanol. Goat anti rabbit IgG (Alexa Fluor® 488, ab150077) was used as a secondary antibody. Isotype control -Rabbit monoclonal IgG (ab172730) (Black). Unlableled control -Unlabelled cells (blue).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Cytokeratin 14 antibody [SP53] (ab119695)

Cytokeratin 14 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-Cytokeratin 14 antibody


Human prostate tissue stained with ab119695 at a dilution of 1/100.



Multiplex immunohistochemistry - Anti-Cytokeratin 14 antibody [SP53] (ab119695)

This data was developed using the same antibody clone in a different buffer formulation (ab236439). Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of Human breast labelling Cytokeratin 18 with ab93741 at 1/200 dilution (1.02 µg/mL) (B), Collagen VI with ab271938 at 1/500 dilution (2.084 µg/ml) (C) and Cytokeratin 14 with ab236439 at 1/2000 dilution (0.519 µg/ml) (D). Opal Polymer HRP Ms + Rb was used as a secondary antibody, and DAPI was used for a nuclear counter stain. Heat mediated antigen retrieval with Citrate buffer (pH 6.0, epitope retrieval solution 1) for 20 mins. Panel A: merged staining of anti-Cytokeratin 14 (magenta; Opal™690), anti-Cytokeratin 18 (green; Opal™520) and anti-Collagen VI (red; Opal™570) on human breast. Panel B: anti-Cytokeratin 18 stained on luminal epithelial cells. Panel C: anti-Collagen VI stained on stroma. Panel D: anti-Cytokeratin 14 stained on myoepithelial cells. The section was incubated in three rounds of staining: in the order of ab236439 for 30 mins, ab93741 for 10 mins, and ab271938 for 30 mins at room temperature. Each round was followed by a separate fluorescent tyramide signal amplification system. The immunostaining was performed on a Leica


Biosystems BOND® RX instrument with an Opal™ 4-color kit. Image acquisition was performed with Leica SP8 confocal microscope.



Immunocytochemistry/ Immunofluorescence - Anti-Cytokeratin 14 antibody [SP53] (ab119695)

Cytokeratin 14 Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-Cytokeratin 14 antibody


Immunocytochemistry/ Immunofluorescence analysis of A431(human epidermoid carcinoma epithelial cell) cells labeling Cytokeratin 14 with purified ab119695. Cells were fixed in 4% paraformaldehyde and permeabilized with 0.1% Triton X-100. Cells were counterstained with ab195889 Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor® 594)). Goat anti rabbit IgG (Alexa Fluor® 488, ab150077) was used as the secondary antibody. DAPI was used as nuclear counterstain. PBS instead of the primary antibody was used as the secondary antibody only control.



Immunohistochemistry (Frozen sections) - Anti-Cytokeratin 14 antibody [SP53] (ab119695)

Cytokeratin 14 Immunohistochemistry (Frozen sections) staining using rabbit Anti-Cytokeratin 14 antibody


Immunohistochemistry (Frozen) analysis of rat skin tissue section labeling Cytokeratin 14 with purified ab119695. Sections were fixed in 4% paraformaldehyde and permeabilized with 0.2% Triton X-100. Antigen retrieval was heat mediated antigen retrieval using sodium citrate buffer (10mM citrate pH 6.0 + 0.05% Tween-20). Goat anti rabbit IgG (Alexa Fluor® 488, ab150077) was used as the secondary antibody. DAPI was used as nuclear counterstain. PBS instead of the primary antibody was used as the secondary antibody only control.



Immunohistochemistry (Frozen sections) - Anti-Cytokeratin 14 antibody [SP53] (ab119695)

Cytokeratin 14 Immunohistochemistry (Frozen sections) staining using rabbit Anti-Cytokeratin 14 antibody

Immunohistochemistry (Frozen) analysis of mouse skin tissue section labeling Cytokeratin 14 with purified ab119695. Sections were fixed in 4% paraformaldehyde and permeabilized with 0.2% Triton X-100. Antigen retrieval was heat mediated antigen retrieval using sodium citrate buffer (10mM citrate pH 6.0 + 0.05% Tween-20). Goat anti rabbit IgG (Alexa Fluor® 488, ab150077) was used as the secondary antibody. DAPI was used as nuclear counterstain. PBS instead of the primary antibody was used as the secondary antibody only control.



Western blot - Anti-Cytokeratin 14 antibody [SP53] (ab119695)

Cytokeratin 14 Western blot staining using rabbit Anti-Cytokeratin 14 antibody

Lanes 1 - 4: Merged signal (red and green). Green - ab119695 observed at 52 kDa. Red - loading control, ab8245 (Mouse anti-GAPDH antibody [6C5]) observed at 37kDa.
ab119695 was shown to react with KRT14 in A431 wild-type cells in Western blot. Loss of signal was observed when KRT14 knockout sample was used. A431 wild-type and KRT14 knockout cell lysates were subjected to SDS-PAGE. Membranes were blocked in 3% Milk in TBS-T (0.1% Tween®) before incubation with ab119695 and ab8245 (Mouse anti-GAPDH antibody [6C5]) overnight at 4°C at a 1 in 93 dilution and a 1 in 20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preabsorbed (ab216773) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preabsorbed (ab216776) secondary antibodies at 1 in 20000 dilution for 1 hour at room temperature before imaging.

All lanes:
Western blot - Anti-Cytokeratin 14 antibody [SP53] (ab119695) at 1/93 dilution

Lane 1:
Wild-type A-431 (Human epidermoid carcinoma cell line) whole cell lysate at 20 µg

Lane 2:
Western blot - Human KRT14 (Cytokeratin 14) knockout A-431 cell lysate (ab261706) at 20 µg

Lane 3:
Human skin whole tissue lysate at 20 µg

Lane 4:
MCF7 (Human breast adenocarcinoma cell line) whole cell lysate at 20 µg

Secondary

Lanes 1 - 4:
Western blot - Goat anti-Rabbit IgG H&L (IRDye® 800CW) preadsorbed (ab216773) at 1/20000 dilution

Lanes 1 - 4:
Western blot - Goat anti-Mouse IgG H&L (IRDye® 680RD) preadsorbed (ab216776) at 1/20000 dilution

Performed under reducing conditions.

Predicted band size: 52 kDa

Observed band size: 52 kDa, 37 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.