abcam

ab123487

Biotin Anti-CD20 antibody [2H7]

Anti-CD20 antibody [2H7] conjugated to Biotin - suitable for flow cytometry, and reacts with human samples. Clone [2H7] is a mouse monoclonal antibody and one of the most commonly used clones for flow cytometry.

Design your panel with our interactive flow cytometry panel builder. Click here.

Key facts

Isotype	IgG2b
Host species	Mouse
Conjugation	Biotin
Storage buffer	pH: 7.4 Preservative: 0.1% Sodium azide Constituents: PBS
Form	Liquid
Clonality	Monoclonal
Immunogen	Cell preparation containing MS4A1 protein. The exact immunogen used to generate this antibody is proprietary information. Database link P11836
Clone number	2H7
Purification technique	Size-exclusion chromatography
Concentration	1 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →
Purification notes	The purified antibody is conjugated with Biotin-LC-NHS under optimum conditions.

Reactivity data

Flow Cyt

Tested

Species	Human
Dilution info	-
Notes	ab18418 - Mouse monoclonal IgG2b, is suitable for use as an isotype control with this antibody.

Target data

See full target information MS4A1

Function	B-lymphocyte-specific membrane protein that plays a role in the regulation of cellular calcium influx necessary for the development, differentiation, and activation of B-lymphocytes (PubMed:12920111, PubMed:3925015, PubMed:7684739). Functions as a store-operated calcium (SOC) channel component promoting calcium influx after activation by the B-cell receptor/BCR (PubMed:12920111, PubMed:18474602, PubMed:7684739).

Storage

Shipped at conditions	Blue Ice
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Storage information	Avoid freeze / thaw cycle

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	CD20 also known as MS4A1 or L26 is a membrane-spanning 4-domains subfamily A member 1 protein. It is a non-glycosylated phosphoprotein with a molecular mass of approximately 33-37 kDa. Expressed extensively on the surface of B-cells CD20 plays an important role in B-cell activation and regulation. While CD20 is absent on early pro-B cells and plasma cells its presence increases as B-cells mature. Anti-CD20 antibodies such as 2H7 are commonly used in research and treatment to deplete B-cells due to the target's consistent expression in most stages of B-cell development.

Biological function summary	CD20 contributes significantly to calcium ion transport which is essential for the activation process of B-cells. Although CD20 does not belong to a larger protein complex its role centers around forming a calcium channel that allows a sustained influx of calcium into the B-cells. This influx is important for the signaling pathways that govern B-cell activation proliferation and differentiation impacting the immune response efficacy.
Pathways	CD20 is closely involved in the B-cell receptor (BCR) signaling pathway and the Fc gamma R-mediated phagocytosis pathway. These pathways play vital roles in adaptive immunity and immune response modulation. CD20's interaction with proteins like PI3K during BCR signaling enhances receptor-mediated cellular signals subsequently influencing downstream effectors involved in cell growth and survival of B-cells.
Associated diseases and disorders	CD20 is strongly linked to certain blood cancers and autoimmune diseases including non-Hodgkin's lymphoma and rheumatoid arthritis. These conditions often exploit aberrant B-cell activity or count. CD20's expression on malignant B-cells in non-Hodgkin's lymphoma makes it an effective target for monoclonal antibody therapies such as rituximab an anti-CD20 antibody. Additionally CD20-targeted therapies can help deplete pathogenic B-cells in rheumatoid arthritis highlighting the protein's relevance in disease treatment.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

1 product image



Flow Cytometry - Biotin Anti-CD20 antibody [2H7] (ab123487)

CD20 Flow Cytometry staining using mouse Anti-CD20 antibody

Flow cytometry of human peripheral blood cells with ab123487 at 4 µg/ml

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.