abcam

ab124904

Anti-Profilin 1 antibody [EPR6304]

Lab Essentials | Recombinant | RabMAb | KO Validated

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR6304
Purification technique	Affinity purification Protein A
Dissociation constant	1.24 x 10^{-10} M
Concentration	0.14 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

IHC-P

Tested

Species	Human
Dilution info	1/500 - 1/1000
Notes	

Perform heat-mediated antigen retrieval before commencing with IHC staining protocol.

Not recommended

Species	Mouse, Rat
Dilution info	-
Notes	-

WB

Tested

Species	Human
Dilution info	1/10000 - 1/50000
Notes	-

Species	Mouse
Dilution info	1/10000 - 1/50000
Notes	-

Species	Rat
Dilution info	1/10000 - 1/50000
Notes	-

ICC/IF

Tested

Species	Human
Dilution info	1/100
Notes	**For unpurified use at 1/250 - 1/500**

Expected

Species	Mouse, Rat
Dilution info	Use at an assay dependent concentration.

Notes	-

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	1/200
Notes	ab172730 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.

Expected

Species	Mouse, Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

Target data

See full target information PFN1

Function	Binds to actin and affects the structure of the cytoskeleton. At high concentrations, profilin prevents the polymerization of actin, whereas it enhances it at low concentrations. By binding to PIP2, it inhibits the formation of IP3 and DG. Inhibits androgen receptor (AR) and HTT aggregation and binding of G-actin is essential for its inhibition of AR.

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot

Storage information	Stable for 12 months at -20°C

Notes

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	Profilin 1 also known as Profilin-1 or Profilin is a small actin-binding protein with a molecular weight of approximately 15 kDa. It is expressed across various tissues including muscle brain and epithelial cells. Profilin 1 interacts with monomeric actin (G-actin) to regulate actin polymerization by binding to actin monomers. This protein also interacts with polyphosphoinositides and proline-rich sequences playing a mechanical role in balancing actin filament assembly and disassembly.
Biological function summary	Profilin 1 acts as an important regulator of the cytoskeleton facilitating cellular processes like motility and division. It participates in the formation of actin filaments by delivering actin monomers to barbed ends promoting polymerization. Profilin 1 is part of the complex that controls actin dynamics and this ability to modulate actin structure allows it to influence cell shape and internal movement. The protein's interactions help support the actin cytoskeleton for intracellular trafficking and signal transduction.
Pathways	Profilin 1 has important roles in the regulation of the actin cytoskeleton and Rho signaling pathways. These pathways are important in cell migration and adhesion. During cellular processes Profilin 1's interactions with actin and polyphosphoinositides link it to other proteins such as Rho GTPases which are associated with cytoskeletal dynamics. Profilin 1 can influence these pathways by modulating actin dynamics and interacting with proteins involved in these signaling cascades.
Associated diseases and disorders	Disruptions in Profilin 1 function are linked to neurodegenerative diseases and cancer metastasis. For example aberrant Profilin 1 activity can contribute to amyotrophic lateral sclerosis (ALS) by affecting neuronal structure and transport. Dysregulation of its interaction with actin or polyphosphoinositides might also lead to enhanced cancer cell migration and invasion. It may interact with other

proteins implicated in these diseases such as the ALS-related protein TDP-43 or cancer-related kinases making it significant in understanding these conditions.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.


12 product images



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Profilin 1 antibody [EPR6304] (ab124904)

Profilin 1 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-Profilin 1 antibody


Unpurified ab124904, at 1/500 dilution, staining Profilin 1 in paraffin-embedded human colon tissue by Immunohistochemistry.
Heat mediated antigen retrieval was performed before commencing with IHC staining protocol.



Immunocytochemistry/ Immunofluorescence - Anti-Profilin 1 antibody [EPR6304] (ab124904)

Profilin 1 Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-Profilin 1 antibody


Immunocytochemistry/ Immunofluorescence analysis of Jurkat (Human T cell leukemia T lymphocyte) cells labeling Profilin 1 with Purified ab124904 at 1:100 dilution (1.4 µg/ml). Cells were fixed in 100% Methanol. Cells were counterstained with ab195889 Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor® 594) 1:200 (2.5 µg/ml). Goat anti rabbit IgG (Alexa Fluor® 488, ab150077) was used as the secondary antibody at 1:1000 (2 µg/ml) dilution. DAPI nuclear counterstain. PBS instead of the primary antibody was used as the secondary antibody only control.



Flow Cytometry (Intracellular) - Anti-Profilin 1 antibody [EPR6304] (ab124904)

Profilin 1 Flow Cytometry (Intracellular) staining using rabbit Anti-Profilin 1 antibody


Overlay histogram showing HeLa cells stained with unpurifiedab124904 (red line). The cells were fixed with 80% methanol (5 min) and then permeabilized with 0.1% PBS-Tween for 20 min. The cells were then incubated in 1x PBS / 10% normal goat serum / 0.3M glycine to block non-specific protein-protein interactions followed by the antibody (ab124904, 1/1000 dilution) for 30 min at 22°C. The secondary antibody used was Alexa Fluor® 488 goat anti-rabbit IgG (H&L) (ab150077) at 1/2000 dilution for 30 min at 22°C. Isotype control antibody (black line) was rabbit IgG (monoclonal) (0.1µg/1x10^6 cells) used under the same conditions. Unlabelled sample (blue line) was also used as a control. Acquisition of >5,000 events were collected using a 20mW Argon ion laser (488nm) and 525/30 bandpass filter.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Profilin 1 antibody [EPR6304] (ab124904)

Profilin 1 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-Profilin 1 antibody


Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of Human breast carcinoma tissue sections labeling Profilin 1 with Purified ab124904 at 1:800 dilution (0.18 µg/ml). Heat mediated antigen retrieval was performed using ab93684 (Tris/EDTA buffer, pH 9.0). ImmunoHistoProbe one step HRP Polymer (ready to use)was used as the secondary antibody. Negative control:PBS instead of the primary antibody. Hematoxylin was used as a counterstain.



Immunocytochemistry/ Immunofluorescence - Anti-Profilin 1 antibody [EPR6304] (ab124904)

Profilin 1 Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-Profilin 1 antibody


Unpurified ab124904, at 1/250 dilution, staining Profilin 1 in HeLa cells by Immunofluorescence.



Flow Cytometry (Intracellular) - Anti-Profilin 1 antibody [EPR6304] (ab124904)

Profilin 1 Flow Cytometry (Intracellular) staining using rabbit Anti-Profilin 1 antibody

Intracellular Flow Cytometry analysis of HeLa (Human cervix adenocarcinoma epithelial cell) cells labeling Profilin 1 with Purified ab124904 at 1/200 dilution (1 µg/ml) (red). Cells were fixed with 80% Methanol. A Goat anti rabbit IgG (Alexa Fluor® 488, ab150077) secondary antibody was used at 1/2000. Isotype control - Rabbit monoclonal IgG (Black). Unlabeled control - Cell without incubation with primary antibody and secondary antibody (Blue).



Western blot - Anti-Profilin 1 antibody [EPR6304] (ab124904)


ab124904 was shown to react with PFN1 in wild-type HAP1 cells in Western blot with loss of signal observed in a PFN1 knockout cell line. Wild-type HAP1 and PFN1 knockout cell lysates were subjected to SDS-PAGE. Membranes were blocked in 5% milk in TBST for 1 hr before incubation with ab124904 overnight at 4 °C at a 1/30000 dilution. Blots were incubated with secondary antibodies at 0.2ug/mL before imaging.

This data was provided by YCharOS Inc., an open science company with the mission of characterizing commercially available antibody reagents for all human proteins. Abcam and YCharOS are working together to help address the reproducibility crisis by enabling the life science community to better evaluate commercially available antibodies.
All lanes:
Western blot - Anti-Profilin 1 antibody [EPR6304] (ab124904) at 1/30000 dilution

Lane 1:
Wild-type HAP1 lysate at 20 µg

Lane 2:
PFN1 knock-out HAP1 lysate at 20 µg

Observed band size: 15 kDa



Western blot - Anti-Profilin 1 antibody [EPR6304] (ab124904)

Profilin 1 Western blot staining using rabbit Anti-Profilin 1 antibody

Lanes 1 - 4: Merged signal (red and green). Green - ab124904 observed at 15 kDa. Red - loading control, ab8245, observed at 37 kDa.
ab124904 was shown to specifically react with in wild-type HAP1 cells as signal was lost in PFN1 knockout cells. Wild-type and PFN1 knockout samples were subjected to SDS-PAGE. The membrane was blocked with 3% NF Milk. ab124904 and ab8245 (Mouse anti-GAPDH loading control) were incubated overnight at 4°C at 1/10000 dilution and 1/20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preabsorbed ab216773 and Goat anti-Mouse IgG H&L (IRDye® 680RD) preabsorbed ab216776 secondary antibodies at 1/20000 dilution for 1 hour at room temperature before imaging.

All lanes:
Western blot - Anti-Profilin 1 antibody [EPR6304] (ab124904) at 1/1000 dilution


Lane 1:
Wild-type HAP1 whole cell lysate at 20 µg

Lane 2:
PFN1 knockout HAP1 whole cell lysate at 20 µg

Lane 3:
HeLa whole cell lysate at 20 µg

Lane 4:
Jurkat whole cell lysate at 20 µg

Predicted band size: 15 kDa



Western blot - Anti-Profilin 1 antibody [EPR6304] (ab124904)

All lanes:
Western blot - Anti-Profilin 1 antibody [EPR6304] (ab124904) at 1/10000 dilution


All lanes:
Human fetal brain lysates at 15 µg

Secondary

All lanes:
Goat Anti-Rabbit IgG (HRP) with minimal cross-reactivity with human IgG at 1/2000 dilution

Predicted band size: 15 kDa

Observed band size: 12 kDa



Western blot - Anti-Profilin 1 antibody [EPR6304] (ab124904)

All lanes:
Western blot - Anti-Profilin 1 antibody [EPR6304] (ab124904) at 1/10000 dilution

Lane 1:
Jurkat cell lysate at 10 µg

Lane 2:
JAR cell lysate at 10 µg

Lane 3:
HepG2 cell lysate at 10 µg


Lane 4:
293T cell lysate at 10 µg

Secondary

All lanes:
HRP labelled goat anti-rabbit at 1/2000 dilution

Predicted band size: 15 kDa

Observed band size: 12 kDa



Western blot - Anti-Profilin 1 antibody [EPR6304] (ab124904)

All lanes:
Western blot - Anti-Profilin 1 antibody [EPR6304] (ab124904) at 1/10000 dilution

Lane 1:
Jurkat (Human T cell leukemia T lymphocyte) whole cell lysates at 15 µg

Lane 2:
Mouse brain lysates at 15 µg


Lane 3:
Rat brain lysates at 15 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/20000 dilution

Predicted band size: 15 kDa

Observed band size: 12 kDa



OI-RD Scanning - Anti-Profilin 1 antibody [EPR6304] (ab124904)

We have systematically measured KD (the equilibrium dissociation constant between the antibody and its antigen), of more than 840 recombinant antibodies to assess not only their individual KD values but also to see the average affinity of antibody. Based on the comparison with published literature values for mouse monoclonal antibodies, Recombinant antibodies appear to be on average 1-2 order of magnitude higher affinity.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.