abcam

ab124953

Anti-Methionine Aminopeptidase 2/p67 antibody [EPR6886(B)]

Recombinant | RabMAb

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR6886(B)
Purification technique	Affinity purification Protein A
Concentration	0.108 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

IP

Not recommended

Species	Mouse, Rat, Human
Dilution info	-
Notes	-

WB

Tested

Species	Human
Dilution info	1/1000 - 1/10000
Notes	-

Species	Mouse
Dilution info	1/1000 - 1/10000
Notes	-

Species	Rat
Dilution info	1/1000 - 1/10000
Notes	-

IHC-P

Not recommended

Species	Mouse, Rat, Human
Dilution info	-
Notes	-

ICC/IF

Not recommended

Species	Mouse, Rat, Human
Dilution info	-
Notes	-

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	1/10 - 1/100

Notes	ab172730 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.

Expected

Species	Mouse, Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

Target data

See full target information METAP2

Function	Cotranslationally removes the N-terminal methionine from nascent proteins. The N-terminal methionine is often cleaved when the second residue in the primary sequence is small and uncharged (Met-Ala-, Cys, Gly, Pro, Ser, Thr, or Val). The catalytic activity of human METAP2 toward Met-Val peptides is consistently two orders of magnitude higher than that of METAP1, suggesting that it is responsible for processing proteins containing N-terminal Met-Val and Met-Thr sequences in vivo. Protects eukaryotic initiation factor EIF2S1 from translation-inhibiting phosphorylation by inhibitory kinases such as EIF2AK2/PKR and EIF2AK1/HCR. Plays a critical role in the regulation of protein synthesis.

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Storage information	Stable for 12 months at -20°C

Notes

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility

- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.


2 product images



Flow Cytometry (Intracellular) - Anti-Methionine Aminopeptidase 2/p67 antibody [EPR6886(B)] (ab124953)

Methionine Aminopeptidase 2/p67 Flow Cytometry (Intracellular) staining using rabbit Anti-Methionine Aminopeptidase 2/p67 antibody

ab124953 at 1/10 dilution staining Methionine Aminopeptidase 2/p67 in permeabilized 293T cells by intracellular flow cytometry (red) or a Rabbit IgG (negative control) (green).



Western blot - Anti-Methionine Aminopeptidase 2/p67 antibody [EPR6886(B)] (ab124953)

Methionine Aminopeptidase 2/p67 Western blot staining using rabbit Anti-Methionine Aminopeptidase 2/p67 antibody

All lanes:
Western blot - Anti-Methionine Aminopeptidase 2/p67 antibody [EPR6886(B)] (ab124953) at 1/1000 dilution

Lane 1:
HepG2 cell lysates at 10 µg

Lane 2:
293T cell lysates at 10 µg

Lane 3:
HeLa cell lysates at 10 µg

Lane 4:
C6 cell lysates at 10 µg

Lane 5:
RAW 264.7 cell lysates at 10 µg

Lane 6:
PC12 cell lysates at 10 µg

Lane 7:
NIH 3T3 cell lysates at 10 µg

Secondary

All lanes:
Goat anti-Rabbit HRP at 1/2000 dilution

Predicted band size: 52 kDa

Observed band size: 67 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.