abcam

ab128162

Anti-CEBP Alpha/CEBPA antibody [4E10]

Key facts

Isotype	IgG1
Host species	Mouse
Storage buffer	Preservative: 0.05% Sodium azide Constituents: PBS
Form	Liquid
Clonality	Monoclonal
Immunogen	Recombinant Fragment Protein within Human CEBPA. The exact immunogen used to generate this antibody is proprietary information. Database link P49715
Clone number	4E10
Purification technique	Affinity purification Protein G
Concentration	1 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →
Purification notes	Purified from tissue culture supernatant.

Reactivity data

Flow Cyt

Tested

Species	Human
Dilution info	1/200 - 1/400
Notes	ab170190 - Mouse monoclonal IgG1, is suitable for use as an isotype control with this antibody.

Not recommended

Species	Recombinant fragment
Dilution info	-
Notes	-

ELISA

Tested

Species	Recombinant fragment
Dilution info	1/10000
Notes	-

Expected

Species	Human
Dilution info	Use at an assay dependent concentration.
Notes	-

WB

Tested

Species	Human
Dilution info	1/500 - 1/2000
Notes	-

Not recommended

Species	Recombinant fragment
Dilution info	-
Notes	-

Target data

See full target information CEBPA

Function	Transcription factor that coordinates proliferation arrest and the differentiation of myeloid progenitors, adipocytes, hepatocytes, and cells of the lung and the placenta. Binds directly to the consensus DNA sequence 5'-T[TG]NNGNAA[TG]-3' acting as an activator on distinct target genes (PubMed:11242107). During early embryogenesis, plays essential and redundant functions with CEBPB. Essential for the transition from common myeloid progenitors (CMP) to granulocyte/monocyte progenitors (GMP). Critical for the proper development of the liver and the lung (By similarity). Necessary for terminal adipocyte differentiation, is required for postnatal maintenance of systemic energy homeostasis and lipid storage (By similarity). To regulate these different processes at the proper moment and tissue, interplays with other transcription factors and modulators. Down-regulates the expression of genes that maintain cells in an undifferentiated and proliferative state through E2F1 repression, which is critical for its ability to induce adipocyte and granulocyte terminal differentiation. Reciprocally E2F1 blocks adipocyte differentiation by binding to specific promoters and repressing CEBPA binding to its target gene promoters. Proliferation arrest also depends on a functional binding to SWI/SNF complex (PubMed:14660596). In liver, regulates gluconeogenesis and lipogenesis through different mechanisms. To regulate gluconeogenesis, functionally cooperates with FOXO1 binding to IRE-controlled promoters and regulating the expression of target genes such as PCK1 or G6PC1. To modulate lipogenesis, interacts and transcriptionally synergizes with SREBF1 in promoter activation of specific lipogenic target genes such as ACAS2. In adipose tissue, seems to act as FOXO1 coactivator accessing to ADIPOQ promoter through FOXO1 binding sites (By similarity). Isoform 3. Can act as dominant-negative. Binds DNA and have transctivation activity, even if much less efficiently than isoform 2. Does not inhibit cell proliferation (PubMed:14660596). Isoform 4. Directly and specifically enhances ribosomal DNA transcription interacting with RNA polymerase I-specific cofactors and inducing histone acetylation.

Storage

Shipped at conditions	Blue Ice
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	CEBP Alpha also known as CEBPA is a transcription factor belonging to the CCAAT/enhancer-binding protein family. It is composed of a molecular mass approximately between 42 to 45 kDa and is characterized by the leucine zipper motif which contributes to its function in dimerization and DNA binding. CEBP Alpha is expressed in various tissues with notable expression in the liver adipose

	tissue and hematopoietic cells where it plays a significant role in regulating gene expression related to growth arrest and differentiation.
Biological function summary	CEBP Alpha regulates important cellular processes by acting as a transcriptional regulator. It participates in the control of cellular differentiation proliferation and metabolism. CEBP Alpha forms homo- or heterodimers with other proteins of the CEBP family to exert its functions. It induces expression of genes involved in differentiation of myeloid progenitors into granulocytes and macrophages. The protein also promotes adipogenesis by activating genes responsible for the development and function of adipocytes.
Pathways	CEBP Alpha participates actively in signaling pathways that involve cell cycle control and hematopoiesis. In the hematopoietic pathway CEBP Alpha influences the differentiation of progenitor cells into mature blood cells. Additionally it interacts with other proteins such as PU.1 an important regulator in the myeloid lineage and through the MAPK pathway influences growth arrest and differentiation responses. These interactions highlight its essential role in maintaining the balance between cell proliferation and differentiation.
Associated diseases and disorders	Alterations in the function or expression of CEBP Alpha have consequential links to acute myeloid leukemia (AML) and obesity. Mutations or disruptions in the CEBP Alpha gene can lead to impaired granulocyte differentiation contributing to the pathogenesis of AML. Additionally CEBP Alpha through its role in adipogenesis is associated with metabolic disorders such as obesity. The protein's interaction with other transcription factors like FLT3 and GATA-2 in the context of these diseases highlights its critical involvement in maintaining normal physiological functions.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.


4 product images



Flow Cytometry - Anti-CEBP Alpha/CEBPA antibody [4E10] (ab128162)

CEBP Alpha/CEBPA Flow Cytometry staining using mouse Anti-CEBP Alpha/CEBPA antibody

Flow cytometric analysis of CEBP Alpha/CEBPA expression in HeLa cells using ab128162 at 1/200 (green) and negative control (red)




Western blot - Anti-CEBP Alpha/CEBPA antibody [4E10] (ab128162)

All lanes:
Western blot - Anti-CEBP Alpha/CEBPA antibody [4E10] (ab128162) at 1/500 dilution

All lanes:
THP1 cell lysate


Predicted band size: 26 kDa, 27 kDa, 38 kDa, 47 kDa, 48 kDa

Observed band size: 47 kDa, 49 kDa



ELISA - Anti-CEBP Alpha/CEBPA antibody [4E10] (ab128162)

Example of standard curve obtained with ab128162



Western blot - Anti-CEBP Alpha/CEBPA antibody [4E10] (ab128162)

All lanes:
Western blot - Anti-CEBP Alpha/CEBPA antibody [4E10] (ab128162) at 1/500 dilution

All lanes:
Human CEBP Alpha/CEBPA (AA: 139-204) recombinant protein

Predicted band size: 38 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.