abcam

ab128939

# Anti-CLP1 antibody [EPR7181]

Recombinant | RabMAb

## Key facts

| | |
|---|---|
| **Isotype** | IgG |
| **Host species** | Rabbit |
| **Storage buffer** | pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA |
| **Form** | Liquid |
| **Clonality** | Monoclonal |
| **Immunogen** | The exact immunogen used to generate this antibody is proprietary information. |
| **Clone number** | EPR7181 |
| **Purification technique** | Affinity purification Protein A |
| **Dissociation constant** | 1.35 x $10^{-10}$ M |
| **Concentration** | 0.587 - 0.62 mg/mL The concentration of this product may be batch-dependent<br>Batch concentration finder → |

## Reactivity data

## IP

### Not recommended

| | |
|---|---|
| **Species** | Human, Mouse, Rat |
| **Dilution info** | - |
| **Notes** | - |

## WB

### Tested

| | |
|---|---|
| **Species** | Human |
| **Dilution info** | 1/1000 - 1/10000 |
| **Notes** | - |

### Predicted

| | |
|---|---|
| **Species** | Mouse, Rat |
| **Dilution info** | - |
| **Notes** | - |

## IHC-P

### Not recommended

| | |
|---|---|
| **Species** | Human, Mouse, Rat |
| **Dilution info** | - |
| **Notes** | - |

## ICC/IF

### Not recommended

| | |
|---|---|
| **Species** | Human, Mouse, Rat |
| **Dilution info** | - |
| **Notes** | - |

## Flow Cyt (Intra)

### Tested

| | |
|---|---|
| **Species** | Human |
| **Dilution info** | 1/100 - 1/500 |

| | |
|---|---|
| **Notes** | ab172730 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody. |

### Predicted

| | |
|---|---|
| **Species** | Mouse, Rat |
| **Dilution info** | - |
| **Notes** | - |

## Target data

See full target information CLP1

| | |
|---|---|
| **Function** | Polynucleotide kinase that can phosphorylate the 5'-hydroxyl groups of double-stranded RNA (dsRNA), single-stranded RNA (ssRNA), double-stranded DNA (dsDNA) and double-stranded DNA:RNA hybrids. dsRNA is phosphorylated more efficiently than dsDNA, and the RNA component of a DNA:RNA hybrid is phosphorylated more efficiently than the DNA component. Plays a key role in both tRNA splicing and mRNA 3'-end formation. Component of the tRNA splicing endonuclease complex: phosphorylates the 5'-terminus of the tRNA 3'-exon during tRNA splicing; this phosphorylation event is a prerequisite for the subsequent ligation of the two exon halves and the production of a mature tRNA (PubMed:24766809, PubMed:24766810). Its role in tRNA splicing and maturation is required for cerebellar development (PubMed:24766809, PubMed:24766810). Component of the pre-mRNA cleavage complex II (CF-II), which seems to be required for mRNA 3'-end formation. Also phosphorylates the 5'-terminus of exogenously introduced short interfering RNAs (siRNAs), which is a necessary prerequisite for their incorporation into the RNA-induced silencing complex (RISC). However, endogenous siRNAs and microRNAs (miRNAs) that are produced by the cleavage of dsRNA precursors by DICER1 already contain a 5'-phosphate group, so this protein may be dispensible for normal RNA-mediated gene silencing. |

## Storage

| | |
|---|---|
| **Shipped at conditions** | Conditional Ambient |
| **Appropriate short-term storage conditions** | +4°C |
| **Appropriate long-term storage conditions** | -20°C |
| **Storage information** | Stable for 12 months at -20°C |

## Notes

**Patented technology**
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

**What are the advantages of a recombinant monoclonal antibody?**
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

## Product promise

**Tested**

We have tested this species and application combination and it works. It is covered by our product promise.

**Expected**

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

**Predicted**

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

**Not recommended**


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.


## 3 product images




## Flow Cytometry (Intracellular) - Anti-CLP1 antibody [EPR7181] (ab128939)

CLP1 Flow Cytometry (Intracellular) staining using rabbit Anti-CLP1 antibody

ab128939 at 1/100 staining CLP1 in permeabilized Jurkat cells by intracellular flow cytometry (red), control Rabbit IgG (green).




## Western blot - Anti-CLP1 antibody [EPR7181] (ab128939)

CLP1 Western blot staining using rabbit Anti-CLP1 antibody

All lanes:
Western blot - Anti-CLP1 antibody [EPR7181] (ab128939) at 1/1000 dilution

Lane 1:
HeLa cell lysate at 10 µg

Lane 2:
HepG2 cell lysate at 10 µg

Lane 3:
293T (Human embryonic kidney epithelial cell) cell lysate at 10 µg


Lane 4:
Jurkat cell lysate at 10 µg

Secondary

All lanes:
HRP labelled goat anti-rabbit at 1/2000 dilution

Predicted band size: 48 kDa

Observed band size: 48 kDa




## OI-RD Scanning - Anti-CLP1 antibody [EPR7181] (ab128939)

We have systematically measured KD (the equilibrium dissociation constant between the antibody and its antigen), of more than 840 recombinant antibodies to assess not only their individual KD values but also to see the average affinity of antibody. Based on the comparison with published literature values for mouse monoclonal antibodies, Recombinant antibodies appear to be on average 1-2 order of magnitude higher affinity.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.