abcam

ab129206

Anti-Peroxiredoxin 3/PRDX3 antibody [EPR8114]

Recombinant | RabMAb | KO Validated | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 50% Tissue culture supernatant, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR8114
Purification technique	Affinity purification Protein A
Concentration	0.099 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

IHC-P

Tested

Species	Human
Dilution info	1/50 - 1/100
Notes	-

IP

Not recommended

Species	Human
Dilution info	-
Notes	-

WB

Tested

Species	Human
Dilution info	1/50000 - 1/200000
Notes	-

ICC/IF

Tested

Species	Human
Dilution info	1/50 - 1/100
Notes	-

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	-
Notes	-

Target data

See full target information PRDX3

Function Thiol-specific peroxidase that catalyzes the reduction of hydrogen peroxide and organic hydroperoxides to water and alcohols, respectively. Plays a role in cell protection against oxidative stress by detoxifying peroxides (PubMed:17707404, PubMed:29438714, PubMed:33889951, PubMed:7733872). Acts synergistically with

MAP3K13 to regulate the activation of NF-kappa-B in the cytosol (PubMed:12492477). Required for the maintenance of physical strength (By similarity).

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Storage information	Stable for 12 months at -20°C

Notes

Species reactivity
Mouse, Rat: We have preliminary internal testing data to indicate this antibody may not react with these species. Please contact us for more information.

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	Peroxiredoxin 3 also known as PRDX3 is an antioxidant enzyme that plays a critical role in the reduction of hydrogen peroxide to water. This protein has a mass of around 25 kDa and belongs to the thioredoxin peroxidase family. It functions within the mitochondria and is expressed in a wide variety of tissues with higher levels in the heart liver and kidney. PRDX3 acts as a protective buffer against oxidative stress by maintaining redox balance in cells.

Biological function summary	The enzyme helps to safeguard cells from oxidative damage by reducing peroxides produced during cellular metabolism. PRDX3 participates in antioxidant defense mechanisms and may form functional complexes with other proteins to enhance its peroxidase activity. Its antioxidant properties contribute to the maintenance of mitochondrial integrity and influence cellular survival and apoptosis processes.
Pathways	PRDX3 integrates into the cellular antioxidant defense system prominently in the thioredoxin pathway. In this pathway PRDX3 synergizes with thioredoxin to neutralize harmful oxidative molecules. PRDX3's activity is interconnected with proteins such as thioredoxin and glutaredoxin which also partake in counteracting oxidative stress bolstering an organism's defense against cellular damage.
Associated diseases and disorders	Alterations in PRDX3 activity associate with conditions like cancer and neurodegenerative disorders. Overexpression of PRDX3 has been observed in certain types of cancer suggesting a potential role in tumor growth through its influence on cell proliferation and survival. In neurodegenerative diseases dysfunctional PRDX3 contributes to neuronal injury. It interacts with proteins like DJ-1 which relate to pathways involved in Parkinson's disease indicating its impact on maintaining neuronal health.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.


6 product images



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Peroxiredoxin 3/PRDX3 antibody [EPR8114] (ab129206)

Peroxiredoxin 3/PRDX3 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-Peroxiredoxin 3/PRDX3 antibody


ab129206 at 1/50 dilution staining Peroxiredoxin 3/PRDX3 in paraffin-embedded Human thyroid gland carcinoma tissue by Immunohistochemistry.
Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Peroxiredoxin 3/PRDX3 antibody [EPR8114] (ab129206)

Peroxiredoxin 3/PRDX3 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-Peroxiredoxin 3/PRDX3 antibody


ab129206 at 1/50 dilution staining Peroxiredoxin 3/PRDX3 in paraffin-embedded Human liver tissue by Immunohistochemistry.
Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.



Flow Cytometry (Intracellular) - Anti-Peroxiredoxin 3/PRDX3 antibody [EPR8114] (ab129206)

Peroxiredoxin 3/PRDX3 Flow Cytometry (Intracellular) staining using rabbit Anti-Peroxiredoxin 3/PRDX3 antibody


Intracellular Flow Cytometry analysis of MCF-7 (human breast carcinoma) cells labeling Peroxiredoxin 3/PRDX3 with unpurified ab129206 at 1/150 dilution (10ug/ml) (red). Cells were fixed with 4% paraformaldehyde and permeabilised with 90% methanol. A Goat anti rabbit IgG (Alexa Fluorr® 488) (1/2000 dilution) was used as the secondary antibody. Rabbit monoclonal IgG (Black) was used as the isotype control, cells without incubation with primary antibody and secondary antibody (Blue) were used as the unlabeled control.



Immunocytochemistry/ Immunofluorescence - Anti-Peroxiredoxin 3/PRDX3 antibody [EPR8114] (ab129206)

Peroxiredoxin 3/PRDX3 Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-Peroxiredoxin 3/PRDX3 antibody

ab129206 at 1/50 dilution staining Peroxiredoxin 3/PRDX3 in MCF-7 cells by Immunofluorescence.



Western blot - Anti-Peroxiredoxin 3/PRDX3 antibody [EPR8114] (ab129206)

Peroxiredoxin 3/PRDX3 Western blot staining using rabbit Anti-Peroxiredoxin 3/PRDX3 antibody

All lanes:
Western blot - Anti-Peroxiredoxin 3/PRDX3 antibody [EPR8114] (ab129206) at 1/50000 dilution

Lane 1:
HeLa cell lysate at 10 µg

Lane 2:
293T (Human embryonic kidney epithelial cell) cell lysate at 10 µg

Lane 3:
MCF7 cell lysate at 10 µg


Lane 4:
A431 cell lysate at 10 µg

Secondary

All lanes:
Goat anti-rabbit HRP at 1/2000 dilution

Predicted band size: 28 kDa

Observed band size: 26 kDa



Western blot - Anti-Peroxiredoxin 3/PRDX3 antibody [EPR8114] (ab129206)

Peroxiredoxin 3/PRDX3 Western blot staining using rabbit Anti-Peroxiredoxin 3/PRDX3 antibody

Lane 1: Wild-type HAP1 cell lysate (20 µg)
Lane 2: Peroxiredoxin 3/PRDX3 knockout HAP1 cell lysate (20 µg)
Lane 3: MCF7 cell lysate (20 µg)
Lane 4: A431 cell lysate (20 µg)
Lanes 1 - 4: Merged signal (red and green). Green - ab129206 observed at 24 kDa. Red - loading control, ab8245, observed at 37 kDa.
ab129206 was shown to specifically react with Peroxiredoxin 3/PRDX3 when Peroxiredoxin 3/PRDX3 knockout samples were used. Wild-type and Peroxiredoxin 3/PRDX3 knockout samples were subjected to SDS-PAGE. ab129206 and ab8245 (loading control to GAPDH) were both diluted 1/50000 and 1/10000 respectively and incubated overnight at 4°C. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preadsorbed ab216773 and Goat anti-Mouse IgG H&L (IRDye® 680RD) preadsorbed ab216776 secondary antibodies at 1/10000 dilution for 1 h at room temperature before imaging.
All lanes:
Western blot - Anti-Peroxiredoxin 3/PRDX3 antibody [EPR8114] (ab129206)

Predicted band size: 28 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.