abcam

ab130751

Anti-Villin antibody [SP145]

Anti-Villin antibody [SP145] (ab130751) is a rabbit monoclonal antibody detecting Villin in **Western Blot, Flow Cytometry (Intra), IP, IHC-P, mIHC**. Suitable for **Human**.

- Multiplex IHC validated on the Leica BOND® MAX using Opal reagents
- Biophysical QC for unrivalled batch-batch consistency
- Over 40 publications

Recombinant | RabMAb | Advanced Validation | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.6 Preservative: 0.1% Sodium azide Constituents: PBS, 1% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	SP145
Purification technique	Affinity purification Protein A/G
Concentration	0.107 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →
Purification notes	Purified from TCS by protein A/G.

Reactivity data

mIHC

Tested

Species	Human
Dilution info	-
Notes	-

Predicted

Species	Mouse, Chicken, Cow, Pig
Dilution info	-
Notes	-

IHC-P

Tested

Species	Human
Dilution info	1/100
Notes	Primary incubation for 10 minutes at room temperature. Perform heat-mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.

Predicted

Species	Mouse, Chicken, Cow, Pig
Dilution info	-
Notes	-

IP

Tested

Species	Human
Dilution info	5 µg/mL
Notes	-

Predicted

Species	Mouse, Chicken, Cow, Pig

Dilution info	-
Notes	-

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	1/100
Notes	ab172730 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.

Predicted

Species	Mouse, Chicken, Cow, Pig
Dilution info	-
Notes	-

WB

Tested

Species	Human
Dilution info	1/100
Notes	Incubate for 1 hour at room temperature.

Predicted

Species	Mouse, Chicken, Cow, Pig
Dilution info	-
Notes	-

Target data

See full target information VIL1

Function	Epithelial cell-specific Ca(2+)-regulated actin-modifying protein that modulates the reorganization of microvillar actin filaments. Plays a role in the actin nucleation,

actin filament bundle assembly, actin filament capping and severing. Binds phosphatidylinositol 4,5-bisphosphate (PIP2) and lysophosphatidic acid (LPA); binds LPA with higher affinity than PIP2. Binding to LPA increases its phosphorylation by SRC and inhibits all actin-modifying activities. Binding to PIP2 inhibits actin-capping and -severing activities but enhances actin-bundling activity. Regulates the intestinal epithelial cell morphology, cell invasion, cell migration and apoptosis. Protects against apoptosis induced by dextran sodium sulfate (DSS) in the gastrointestinal epithelium. Appears to regulate cell death by maintaining mitochondrial integrity. Enhances hepatocyte growth factor (HGF)-induced epithelial cell motility, chemotaxis and wound repair. Upon S.flexneri cell infection, its actin-severing activity enhances actin-based motility of the bacteria and plays a role during the dissemination.

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	+4°C
Storage information	Do Not Freeze

Notes

What is this antibody validated in?
Anti-Villin antibody [SP145] (ab130751) is a rabbit recombinant monoclonal antibody and is validated for use in Western Blot (WB), Flow Cytometry (Intra), Immunoprecipitation (IP), Immunohistochemistry (IHC-P), Multiplex IHC (mIHC) in Human samples.

What is the molecular weight of Villin?
Anti-Villin [SP145] (ab130751) specifically detects a band for Villin (UniProt: P09327) at a molecular weight of 93kDa.

Collaboration
Anti-Villin [SP145] (ab130751) is a clone from the portfolio of Spring Bioscience (Roche) SP clones which have been optimised for immunohistochemistry (IHC).

Trusted by the scientific community
Anti-Villin [SP145] (ab130751) was first used in a scientific publication in 2012 and has been cited over 40 times in peer-reviewed journals.

Other related products
We have a range of other formats of antibody clone [SP145] also available for your convenience: ab130751, Carrier free - ab245749, Alexa Fluor® 488 - ab270921, Alexa Fluor® 647 - ab270922, ab279636, APC - ab319336, PE - ab319472, Alexa Fluor® 594 - ab319829, Alexa Fluor® 555 - ab319970, Alexa Fluor® 750 - ab321039

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Trial sizes available!
Test your antibody or perform pre-screening before committing to a larger quantity. Sold in 20µl. Discover our selection of trial-size antibodies.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended

We do not recommend this combination. It is not covered by our product promise.


We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

---


6 product images



Flow Cytometry (Intracellular) - Anti-Villin antibody [SP145] (ab130751)

Villin Flow Cytometry (Intracellular) staining using rabbit Anti-Villin antibody


Overlay histogram showing Caco 2 cells stained with ab130751 (red line). The cells were fixed with 80% methanol (5 min) and then permeabilized with 0.1% PBS-Tween for 20 min. The cells were then incubated in 1x PBS / 10% normal goat serum / 0.3M glycine to block non-specific protein-protein interactions followed by the antibody (ab130751 1/100 dilution) for 30 min at 22°C. The secondary antibody used was Alexa Fluor® 488 goat anti-rabbit IgG (H&L) (ab150077) at 1/2000 dilution for 30 min at 22°C. Isotype control antibody (black line) was rabbit IgG (monoclonal) (1µg/1x10^6 cells) used under the same conditions. Unlabelled sample (blue line) was also used as a control. Acquisition of >5000 events were collected using a 20mW Argon ion laser (488nm) and 525/30 bandpass filter.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Villin antibody [SP145] (ab130751)


Villin Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-Villin antibody

ab130751 at 1/100 dilution staining Villin in formalin-fixed paraffin-embedded Human colon tissue by Immunohistochemistry.



Multiplex immunohistochemistry - Anti-Villin antibody [SP145] (ab130751)

Fluorescence multiplex immunohistochemical analysis of the human colon (Formalin/PFA-fixed paraffin-embedded sections). Panel A: merged staining of anti-Villin (ab245749, gray; Opal™690), anti-liver FABP (ab240401, green; Opal™520) and anti-MUC2 (ab272692, red; Opal™570) on human colon. Panel B: anti-liver FABP stained on enterocytes. Panel C: anti-Villin stained on apical border. Panel D: anti-MUC2 stained on goblet cells. Rabbit specific IHC polymer detection kit HRP/DAB (ab209101) was used as a secondary antibody. The immunostaining was performed on a Leica Biosystems BOND® RX instrument with an Opal™ 4-color kit. The section was incubated in three rounds of staining: in the order of ab245749 (1/1000 dilution), ab240401 (1/8000 dilution), and ab272692 (1/5000 dilution) for 30 mins at room temperature. Each round was followed by a separate fluorescent tyramide signal amplification system. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 20 mins. DAPI (blue) was used as a nuclear counter stain. Image acquisition was performed with Leica SP8 confocal microscope.



Flow Cytometry (Intracellular) - Anti-Villin antibody [SP145] (ab130751)

Villin Flow Cytometry (Intracellular) staining using rabbit Anti-Villin antibody


Flow cytometry overlay histogram showing left HT-29 positive cells and right negative PANC-1 stained with ab130751 (red line). The cells were fixed with 80% methanol (5 min) and then permeabilised with 0.1% PBS-Triton X-100 for 15 min. The cells were then incubated in 1x PBS containing 10% normal goat serum to block non-specific protein-protein interaction followed by the antibody (ab130751) (1x 106 in 100µl at 0.2µg/ml (1/11200)) for 30min at 22°C.

The secondary antibody Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed was incubated at 1/4000 for 30min at 22°C

Isotype control antibody (black line) was Recombinant Rabbit IgG, monoclonal [EPR25A] - Isotype Control used at the same concentration and conditions as the primary antibody. Unlabelled sample (blue line) was also used as a control.

Acquisition of >5000 events were collected using a 50 mW Blue laser (488nm) and 525/40 bandpass filter.

This antibody gave a positive signal in HT-29 Fixed with 4% formaldehyde (10 min) / permeabilised with 0.1% PBS-Triton X-100 for 15 min under the same conditions.



Immunoprecipitation - Anti-Villin antibody [SP145] (ab130751)

Villin was immunoprecipitated using 0.5mg SW480 whole cell extract, 5µg of Rabbit polyclonal to Villin and 50µl of protein G magnetic beads (+). No antibody was added to the control (-).
The antibody was incubated under agitation with Protein G beads for 10min, SW480 whole cell extract lysate diluted in RIPA buffer was added to each sample and incubated for a further 10min under agitation.

Proteins were eluted by addition of 40µl SDS loading buffer and incubated for 10min at 70°C; 10µl of each sample was separated on a SDS PAGE gel, transferred to a nitrocellulose membrane, blocked with 5% BSA and probed with ab130751.

Secondary: Mouse monoclonal [SB62a] Secondary Antibody to Rabbit IgG light chain (HRP) (ab99697).

Band: 93kDa; Villin


All lanes:
Immunoprecipitation - Anti-Villin antibody [SP145] (ab130751)

Developed using the ECL technique.

Performed under reducing conditions.

Predicted band size: 93 kDa

Exposure time: 16min



Western blot - Anti-Villin antibody [SP145] (ab130751)

Villin Western blot staining of HT-29 cell lysate using rabbit Anti-Villin antibody

All lanes:
Western blot - Anti-Villin antibody [SP145] (ab130751) at 1/100 dilution

All lanes:
HT-29 cell lysate

Predicted band size: 93 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.