abcam

ab131263

Anti-IDH2 antibody [EPR7577]

Anti-IDH2 antibody [EPR7577] (ab131263) is a rabbit monoclonal antibody detecting IDH2 in **Western Blot, Flow Cytometry (Intra), IHC-P, ICC/IF**. Suitable for **Human, Mouse, Rat**.

- KO validated for confirmed specificity
- Biophysical QC for unrivalled batch-batch consistency
- Over 10 publications

Recombinant | RabMAb | KO Validated | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 50% Tissue culture supernatant, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR7577
Purification technique	Affinity purification Protein A
Dissociation constant	4.7 x 10^{-11} M
Concentration	0.202 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

IHC-P

Tested

Species	Human
Dilution info	1/100 - 1/250
Notes	Perform heat-mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.

Species	Rat
Dilution info	1/100 - 1/250
Notes	Perform heat-mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.

Expected

Species	Mouse
Dilution info	Use at an assay dependent concentration.
Notes	-

ICC/IF

Tested

Species	Human
Dilution info	1/1000
Notes	-

Expected

Species	Rat, Mouse
Dilution info	Use at an assay dependent concentration.
Notes	-

WB

Tested

Species	Human
Dilution info	1/1000 - 1/10000

Notes	-

Species	Mouse
Dilution info	1/1000 - 1/10000
Notes	-

Species	Rat
Dilution info	1/1000 - 1/10000
Notes	-

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	1/100 - 1/1000
Notes	ab172730 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.

Expected

Species	Rat, Mouse
Dilution info	Use at an assay dependent concentration.
Notes	-

Target data

See full target information IDH2

Function	Plays a role in intermediary metabolism and energy production (PubMed:19228619, PubMed:22416140). It may tightly associate or interact with the pyruvate dehydrogenase complex (PubMed:19228619, PubMed:22416140).

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage conditions	+4°C

Appropriate long-term storage conditions	-20°C
Storage information	Stable for 12 months at -20°C

Notes

What is this antibody validated in?
Anti-IDH2 antibody [EPR7577] (ab131263) is a rabbit recombinant monoclonal antibody and is validated for use in Western Blot (WB), Flow Cytometry (Intra), Immunohistochemistry (IHC-P), Immunocytochemistry/immunofluorescence (ICC/IF) in Human, Mouse, Rat samples.

What is the molecular weight of IDH2?
Anti-IDH2 [EPR7577] (ab131263) specifically detects a band for IDH2 (UniProt: P48735) at a molecular weight of 50kDa.

Trusted by the scientific community
Anti-IDH2 [EPR7577] (ab131263) was first used in a scientific publication in 2012 and has been cited over 10 times in peer-reviewed journals.

Specificity confirmed
The specificity of Anti-IDH2 antibody [EPR7577] (ab131263) has been confirmed by Western blot testing in IDH2 Knockout HAP1 cell line, ab282331.

Other related products
We have a range of other formats of antibody clone [EPR7577] also available for your convenience: ab131263, PE - ab212122, Carrier free - ab230796, APC - ab317932, Alexa Fluor® 555 - ab318055

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Trial sizes available!
Test your antibody or perform pre-screening before committing to a larger quantity. Sold in 20µl. Discover our selection of trial-size antibodies.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	Isocitrate dehydrogenase 2 (IDH2) is an enzyme that converts isocitrate to alpha-ketoglutarate in the citric acid cycle through oxidative decarboxylation. IDH2 also known as NADP-dependent isocitrate dehydrogenase has a molecular weight of about 51 kDa. This protein expresses in the mitochondria serving an important role in cellular energy production and intermediary metabolism. By facilitating this conversion IDH2 impacts cellular respiration and energy balance.
Biological function summary	The IDH2 enzyme facilitates cellular metabolism by producing NADPH critical for biosynthesis and antioxidant defense. It functions within the oxidative decarboxylation of isocitrate providing reducing equivalents to keep cellular redox balance. Although not part of a larger enzymatic complex IDH2 acts synergistically with other metabolic enzymes to maintain cellular biochemical pathways.
Pathways	The IDH2 protein participates in the citric acid cycle and redox homeostasis. IDH2 contributes to the tricarboxylic acid (TCA) pathway coupling with other TCA components such as citrate synthase and aconitase. Within the redox pathway it influences glucose metabolism via its link with enzymes like NADPH oxidase ensuring a steady supply of NADPH for biosynthetic reactions.
Associated diseases and disorders	The IDH2 protein links to certain cancers like acute myeloid leukemia (AML) due to mutations such as IDH2 R140Q and IDH2 R172K which lead to the production of oncometabolite 2-hydroxyglutarate. This oncometabolite influences epigenetic regulation and cellular differentiation. IDH2 mutations often associate with altered tumor suppressor pathways involving proteins like TP53 contributing to tumorigenesis and impaired cellular differentiation.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.


11 product images



Immunocytochemistry/ Immunofluorescence - Anti-IDH2 antibody [EPR7577] (ab131263)

IDH2 Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-IDH2 antibody


Immunocytochemistry analysis of HeLa (Human cervix adenocarcinoma epithelial cell) cells labeling IDH2 with Purified ab131263 at 1:1000 dilution (0.2 µg/ml). Cells were fixed in 4% Paraformaldehyde and permeabilized with 0.1% tritonX-100. Cells were counterstained with ab195889 Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor® 594) 1:200 (2.5 µg/ml). Goat anti rabbit IgG (Alexa Fluor® 488, ab150077) was used as the secondary antibody at 1:1000 (2 µg/ml) dilution. DAPI (blue) was used as nuclear counterstain. PBS instead of the primary antibody was used as the secondary antibody only control.



Flow Cytometry (Intracellular) - Anti-IDH2 antibody [EPR7577] (ab131263)

IDH2 Flow Cytometry (Intracellular) staining using rabbit Anti-IDH2 antibody


Flow Cytometry analysis of MCF7 (Human breast adenocarcinoma epithelial cell) cells labelling IDH2 with Purified ab131263 at 1:20 dilution (10 µg/ml) (Red). Cells were fixed with 4% Paraformaldehyde and permeabilised with 90% Methanol. A Goat anti rabbit IgG (Alexa Fluor® 488, ab150081) secondary antibody was used at 1:2000. Isotype control - Rabbit monoclonal IgG (Black). Unlabelled control - Cell without incubation with primary antibody and secondary antibody (Blue).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-IDH2 antibody [EPR7577] (ab131263)

IDH2 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-IDH2 antibody

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human liver tissue sections labeling IDH2 with Purified ab131263 at 1:100 (2.11 µg/ml). Heat mediated antigen retrieval was performed using Heat mediated antigen retrieval using Bond™ Epitope Retrieval Solution 2 (pH 9.0) . Tissue was counterstained with Hematoxylin. Rabbit specific IHC polymer detection kit HRP/DAB (ab209101) secondary antibody was used. PBS instead of the primary antibody was used as the negative control.



Flow Cytometry (Intracellular) - Anti-IDH2 antibody [EPR7577] (ab131263)

IDH2 Flow Cytometry (Intracellular) staining of Jurkat using rabbit Anti-IDH2 antibody


Flow cytometry overlay histogram showing wild-type Jurkat (green line) and IDH2 knockout Jurkat stained with ab131263 (magenta line). The cells were fixed with 80% methanol (5 min) and then permeabilised with 0.1% PBS-Triton X-100 for 15 min. The cells were then incubated in 1x PBS

containing 10% normal goat serum to block non-specific protein-protein interaction followed by the antibody (ab131263) (1x 10^6 in 100µl at 0.008 µg/ml (1/27625)) for 30min at 22°C.

The secondary antibody Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed was incubated at 1/4000 for 30min at 22°C.

Isotype control antibody was Recombinant Rabbit IgG, monoclonal [EPR25A] - Isotype Control in Jurkat WT cells (black line) and Jurkat-IDH2 KO cells (grey line), used at the same concentration and conditions as the primary antibody.


Acquisition of >5000 events were collected using a 50 mW Blue laser (488nm) and 530/30 bandpass filter.



Immunocytochemistry/ Immunofluorescence - Anti-IDH2 antibody [EPR7577] (ab131263)

IDH2 Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-IDH2 antibody


Immunofluorescent analysis of 4% Paraformaldehyde-fixed, 0.1% Triton X-100 permeabilized Jurkat IDH2 knockout cells labeling IDH2 with ab131263 at 1 µg/ml. Cells were counterstained with ab7291 Anti-alpha Tubulin antibody [DM1A] at 1 : 1000 (1 µg/ml). Goat anti rabbit IgG (Alexa Fluor® 488 ab150081, green) and goat anti mouse IgG (Alexa Fluor® 594 ab150120, magenta) were used as the secondary antibodies at 1 : 1000 (2 µg/ml) dilution. The nuclear counterstain was DAPI (Blue).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-IDH2 antibody [EPR7577] (ab131263)

IDH2 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-IDH2 antibody


Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of mouse spleen tissue sections labeling IDH2 with Purified ab131263 at 1:100 (2.11 µg/ml). Heat mediated antigen retrieval was performed using Heat mediated antigen retrieval using Bond™ Epitope Retrieval Solution 2 (pH 9.0) . Tissue was counterstained with Hematoxylin. Rabbit specific IHC polymer detection kit HRP/DAB (ab209101) secondary antibody was used. PBS instead of the primary antibody was used as the negative control.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-IDH2 antibody [EPR7577] (ab131263)

IDH2 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-IDH2 antibody

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of rat spleen tissue sections labeling IDH2 with Purified ab131263 at 1:100 (2.11 µg/ml). Heat mediated antigen retrieval was performed using Heat mediated antigen retrieval using Bond™ Epitope Retrieval Solution 2 (pH 9.0) . Tissue was counterstained with Hematoxylin. Rabbit specific IHC polymer detection kit HRP/DAB (ab209101) secondary antibody was used. PBS instead of the primary antibody was used as the negative control.



Western blot - Anti-IDH2 antibody [EPR7577] (ab131263)

IDH2 Western blot staining using rabbit Anti-IDH2 antibody

Western blot: Anti-IDH2 antibody [EPR7577] (ab131263) staining at 1/1000 dilution, shown in green; Mouse anti-GAPDH antibody [6C5] (ab8245) loading control staining at 1/20000 dilution, shown in magenta. In Western blot, ab131263 was shown to bind specifically to IDH2. A band was observed at 51 kDa in wild-type A549 cell lysates with no signal observed at this size in IDH2 knockout cell line. To generate this image, wild-type and IDH2 knockout A549 cell lysates were analysed. First, samples were run on an SDS-PAGE gel then transferred onto a nitrocellulose membrane. Membranes were blocked in 3 % milk in TBS-0.1 % Tween® 20 (TBS-T) before incubation with primary antibodies overnight at 4 °C. Blots were washed four times in TBS-T, incubated with secondary antibodies for 1 h at room temperature, washed again four times then imaged. Secondary antibodies used were Goat anti-Rabbit IgG H&L 800CW and Goat anti-Mouse IgG H&L 680RD at 1/20000 dilution.
All lanes:
Western blot - Anti-IDH2 antibody [EPR7577] (ab131263) at 1/1000 dilution

Lane 1:
Wild-type A549 cell lysate at 20 µg

Lane 2:
IDH2 knockout A549 cell lysate at 20 µg

Lane 3:
Wild-type Jurkat cell lysate at 20 µg


Lane 4:
IDH2 knockout Jurkat ab282331 cell lysate at 20 µg

Secondary

All lanes:
Goat anti-Rabbit IgG H&L 800CW and Goat anti-Mouse IgG H&L 680RD at 1/20000 dilution

Performed under reducing conditions.

Observed band size: 51 kDa



Western blot - Anti-IDH2 antibody [EPR7577] (ab131263)

IDH2 Western blot staining using rabbit Anti-IDH2 antibody

False colour image of Western blot: Anti-IDH2 antibody [EPR7577] staining at 1/1000 dilution, shown in green; Mouse anti-GAPDH antibody [6C5] (ab8245) loading control staining at 1/20000 dilution, shown in red. In Western blot, ab131263 was shown to bind specifically to IDH2. A band was observed at 48 kDa in wild-type Jurkat cell lysates with no signal observed at this size in IDH2 knockout cell line ab282331 (knockout cell lysate ab283148). To generate this image, wild-type and IDH2 knockout Jurkat cell lysates were analysed. First, samples were run on an SDS-PAGE gel then transferred onto a nitrocellulose membrane. Membranes were blocked in 3% milk in TBS-0.1 % Tween® 20 (TBS-T) before incubation with primary antibodies overnight at 4°C. Blots were washed four times in TBS-T, incubated with secondary antibodies for 1 h at room temperature, washed again four times then imaged. Secondary antibodies used were Goat anti-Rabbit IgG H&L (IRDye® 800CW) preabsorbed (ab216773) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preabsorbed (ab216776) at 1/20000 dilution.
All lanes:
Western blot - Anti-IDH2 antibody [EPR7577] (ab131263) at 1/1000 dilution

Lane 1:

Wild-type Jurkat cell lysate at 20 µg


Lane 2:
IDH2 knockout Jurkat cell lysate at 20 µg

Lane 2:
Western blot - Human IDH2 knockout Jurkat cell line (ab282331)

Performed under reducing conditions.

Predicted band size: 50 kDa

Observed band size: 48 kDa




Western blot - Anti-IDH2 antibody [EPR7577] (ab131263)

IDH2 Western blot staining using rabbit Anti-IDH2 antibody

Lane 1: Wild-type HAP1 cell lysate (20 µg)
Lane 2: IDH2 knockout HAP1 cell lysate (20 µg)
Lane 3: K562 cell lysate (20 µg)
Lane 4: HepG2 cell lysate (20 µg)
Lanes 1 - 4: Merged signal (red and green). Green - ab131263 observed at 48 kDa. Red - loading control, ab8245, observed at 37 kDa.
ab131263 was shown to recognize IDH2 when IDH2 knockout samples were used, along with additional cross-reactive bands. Wild-type and IDH2 knockout samples were subjected to SDS-PAGE. ab131263 and ab8245 (loading control to GAPDH) were both diluted 1/10 000 and incubated overnight at 4°C. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preadsorbed (ab216773) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preadsorbed (ab216776) secondary antibodies at 1/10000 dilution for 1 h at room temperature before imaging.
All lanes:
Western blot - Anti-IDH2 antibody [EPR7577] (ab131263)

Predicted band size: 50 kDa



Western blot - Anti-IDH2 antibody [EPR7577] (ab131263)

All lanes:
Western blot - Anti-IDH2 antibody [EPR7577] (ab131263) at 1/5000 dilution

Lane 1:
MOLT-4 (Human lymphoblastic leukemia T lymphoblast) whole cell lysate at 20 µg

Lane 2:
K-562 (Human chronic myelogenous leukemia lymphoblast) whole cell lysate at 20 µg

Lane 3:
U-87 MG (Human glioblastoma-astrocytoma epithelial cell) whole cell lysate at 20 µg

Lane 4:
HepG2 (Human hepatocellular carcinoma epithelial cell) whole cell lysate at 20 µg

Lane 5:
Mouse liver lysate at 20 µg

Lane 6:
Rat liver lysate at 20 µg

Lane 7:
Mouse kidney lysate at 20 µg

Lane 8:
Rat kidney lysate at 20 µg

Lane 9:
Rat stomach lysate at 20 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/20000 dilution

Predicted band size: 50 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.