abcam

ab133479

Anti-Ube1L + E1 Ubiquitin Activating Enzyme 1/UBA1 antibody [EPR4269(2)]

Recombinant | RabMAb

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR4269(2)
Purification technique	Affinity purification Protein A
Concentration	Concentration data is not available right now, please try again later Batch concentration finder →

Reactivity data

IHC-P

Tested

Species	Human
Dilution info	1/100 - 1/250

Notes	Perform heat-mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.

Predicted

Species	Mouse, Rat
Dilution info	-
Notes	-

WB

Tested

Species	Human
Dilution info	1/10000 - 1/50000
Notes	-

Predicted

Species	Mouse, Rat
Dilution info	-
Notes	-

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	1/100 - 1/1000
Notes	ab172730 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.

Predicted

Species	Mouse, Rat
Dilution info	-
Notes	-

Target data

See full target information UBA7

Function	E1-activating enzyme that catalyzes the covalent conjugation of the ubiquitin-like protein product of ISG15 to additional interferon stimulated proteins (ISGs) as well as other cellular proteins such as P53 in a process termed protein ISGylation (PubMed:27545325). Plays an essential role in antiviral immunity together with ISG15 by restricting the replication of many viruses including rabies virus, influenza virus, sindbis virus, rotavirus or human cytomegalovirus (PubMed:16254333, PubMed:19073728, PubMed:29056542, PubMed:29743376, PubMed:37722521). For example, ISG15 modification of influenza A protein NS1 disrupts the association of the NS1 with importin-alpha leading to NS1 nuclear import inhibition (PubMed:20133869). ISGylation of human cytomegalovirs protein UL26 regulates its stability and inhibits its activities to suppress NF-kappa-B signaling (PubMed:27564865).

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Storage information	Stable for 12 months at -20°C

Notes

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	Ube1L also known as E1 Ubiquitin Activating Enzyme 1 or UBA1 is a protein that plays a critical role mechanically in the ubiquitination process. This process involves the tagging of proteins for degradation by the proteasome which is essential for regulating protein turnover in cells. Ube1L has an approximate mass of 117 kDa and is expressed in various tissues across the body including the liver and muscle. The enzyme catalyzes the first step in the ubiquitination pathway by activating ubiquitin molecules in an ATP-dependent manner which is necessary for transferring ubiquitin to substrate proteins.
Biological function summary	Ube1L is essential for maintaining protein homeostasis within the cell by regulating the degradation of misfolded damaged or unneeded proteins. It operates as part of the ubiquitin-proteasome system a complex assembly of various enzymes and proteins that coordinate the target-specific degradation of proteins. This ubiquitin-proteasome system helps control protein levels to ensure cellular functions such as DNA repair cell cycle progression and apoptosis occur smoothly.
Pathways	Ube1L is integral in the ubiquitin-dependent catabolic pathway and the NF-κB signaling pathway. In the ubiquitin-dependent catabolic pathway Ube1L works alongside proteins like UbcH5B and Ubc13 to transfer ubiquitin to substrates that are marked for destruction. Then in the NF-κB signaling pathway Ube1L indirectly regulates transcription factors which influences immune response and inflammation through the conjugation and deconjugation of ubiquitin on inhibitor proteins like IκB.
Associated diseases and disorders	Disruptions in Ube1L function associate with cancer and neurodegenerative disorders such as amyotrophic lateral sclerosis (ALS). Cancer cells often exploit the dysregulation of the ubiquitin-proteasome system to avoid apoptosis and grow uncontrollably. Ube1L's interaction with proteins like p53 a tumor suppressor highlights its role in cancer pathogenesis. In ALS abnormal protein accumulations that result from inefficient ubiquitination link back to Ube1L's function and its interaction with proteins like TDP-43 has been studied to understand its contribution to disease progression.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.


4 product images



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Ube1L + E1 Ubiquitin Activating Enzyme 1/UBA1 antibody [EPR4269(2)] (ab133479)

Ube1L + E1 Ubiquitin Activating Enzyme 1/UBA1 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-Ube1L + E1 Ubiquitin Activating Enzyme 1/UBA1 antibody


Immunohistochemical analysis of paraffin-embedded human urinary bladder carcinoma tissue labelled with ab133479 at 1/100 dilution.
Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.



Flow Cytometry (Intracellular) - Anti-Ube1L + E1 Ubiquitin Activating Enzyme 1/UBA1 antibody [EPR4269(2)] (ab133479)

Ube1L + E1 Ubiquitin Activating Enzyme 1/UBA1 Flow Cytometry (Intracellular) staining using rabbit Anti-Ube1L + E1 Ubiquitin Activating Enzyme 1/UBA1 antibody

Overlay histogram showing HeLa cells stained with ab133479 (red line). The cells were fixed with 80% methanol (5 min) and then permeabilized with 0.1% PBS-Tween for 20 min. The cells were then incubated in 1x PBS / 10% normal goat serum / 0.3M glycine to block non-specific protein-protein interactions followed by the antibody (ab133479, 1/1000 dilution) for 30 min at 22°C. The secondary antibody used was Alexa Fluor® 488 goat anti-rabbit IgG (H&L) (ab150077) at 1/2000 dilution for 30 min at 22°C. Isotype control antibody (black line) was rabbit IgG (monoclonal) (0.1µg/1x10^6 cells) used under the same conditions. Unlabelled sample (blue line) was also used as a control. Acquisition of >5,000 events were collected using a 20mW Argon ion laser (488nm) and 525/30 bandpass filter.



Western blot - Anti-Ube1L + E1 Ubiquitin Activating Enzyme 1/UBA1 antibody [EPR4269(2)] (ab133479)

All lanes:
Western blot - Anti-Ube1L + E1 Ubiquitin Activating Enzyme 1/UBA1 antibody [EPR4269(2)] (ab133479) at 1/10000 dilution

Lane 1:

HL60 cell lysate at 10 µg


Lane 2:
HepG2 cell lysate at 10 µg

Lane 3:
HeLa cell lysate at 10 µg

Lane 4:
Fetal liver tissue lysate at 10 µg

Secondary

All lanes:
HRP labelled goat anti-rabbit at 1/2000 dilution



Western blot - Anti-Ube1L + E1 Ubiquitin Activating Enzyme 1/UBA1 antibody [EPR4269(2)] (ab133479)

Blocking/Diluting buffer and concentration: 5% NFDM/TBST
All lanes:
Western blot - Anti-Ube1L + E1 Ubiquitin Activating Enzyme 1/UBA1 antibody [EPR4269(2)] (ab133479) at 1/1000 dilution

Lane 1:
GST tagged Recombinant human UBA7(Ube1L) protein (Full length, 166 KDa)

Lane 2:
GST tagged Recombinant human UBA1 protein (Full length, 146 KDa)

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/20000 dilution

Observed band size: 146 kDa, 166 kDa

Exposure time: 180s

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.