abcam

ab133590

Anti-IRF4 antibody [EP5699]

Recombinant | RabMAb | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 50% Tissue culture supernatant, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EP5699
Purification technique	Affinity purification Protein A
Concentration	0.215 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

ICC/IF

Tested

Species	Human
Dilution info	1/500
Notes	-

Not recommended

Species	Mouse
Dilution info	-
Notes	-

ChIP

Not recommended

Species	Human, Mouse
Dilution info	-
Notes	-

IHC-P

Tested

Species	Human
Dilution info	1/250 - 1/500
Notes	Perform heat-mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.

Not recommended

Species	Mouse
Dilution info	-
Notes	Perform heat-mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.

IP

Tested

Species	Human
Dilution info	1/100 - 1/500
Notes	-

Not recommended

Species	Mouse
Dilution info	-
Notes	-

WB

Tested

Species	Human
Dilution info	1/10000 - 1/50000
Notes	-

Not recommended

Species	Mouse
Dilution info	-
Notes	-

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	1/100 - 1/1000
Notes	ab172730 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.

Not recommended

Species	Mouse
Dilution info	-
Notes	ab172730 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.

Target data

See full target information IRF4

Function	Transcriptional activator. Binds to the interferon-stimulated response element (ISRE) of the MHC class I promoter. Binds the immunoglobulin lambda light chain enhancer, together with PU.1. Probably plays a role in ISRE-targeted signal transduction mechanisms specific to lymphoid cells. Involved in CD8(+) dendritic cell differentiation by forming a complex with the BATF-JUNB heterodimer in immune cells, leading to recognition of AICE sequence (5'-TGAnTCA/GAAA-3'), an immune-specific regulatory element, followed by cooperative binding of BATF and IRF4 and activation of genes.

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

Species reactivity
Mouse, Rat: We have preliminary internal testing data to indicate this antibody may not react with these species. Please contact us for more information.

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	The IRF4 protein also known as Interferon Regulatory Factor 4 is a transcription factor encoded by the IRF4 gene. It weighs approximately 52 kDa. IRF4 is expressed mainly in lymphoid cells including B cells T cells and macrophages. It plays a mechanical role in regulating gene expression by binding to specific DNA sequences. Through these interactions IRF4 influences various cellular processes important for the immune response.
Biological function summary	IRF4 helps in the activation and differentiation of immune cells. It forms complexes with other transcription factors like PU.1 enhancing its regulatory functions. IRF4 also contributes to the maturation of B and T lymphocytes impacting the adaptive immune system. It acts by modulating expression of cytokines and surface markers involved in immune response.
Pathways	IRF4 functions within the immune response and cell differentiation pathways. It interacts with pathways like NF-kB and MAPK linking it to important immune regulatory processes. IRF4 associates with proteins such as c-Rel in the NF-kB pathway influencing gene expression related to immune activation and survival.
Associated diseases and disorders	IRF4 plays a role in multiple myeloma and autoimmune disorders. Aberrant expression of IRF4 can lead to dysregulated immune cell proliferation as seen in multiple myeloma. It also connects with proteins like BLIMP-1 contributing to pathogenesis of autoimmune conditions. Understanding IRF4's involvement provides insights into potential therapeutic targets for such diseases.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.


6 product images



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-IRF4 antibody [EP5699] (ab133590)

IRF4 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-IRF4 antibody


Immunohistochemical analysis of paraffin embedded Human tonsil tissue labelling IRF4, using ab133590 at a dilution of 1/250.
Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-IRF4 antibody [EP5699] (ab133590)

IRF4 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-IRF4 antibody


Immunohistochemical analysis of paraffin embedded Human lymphoma tissue labelling IRF4, using ab133590 at a dilution of 1/250.
Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.



Flow Cytometry (Intracellular) - Anti-IRF4 antibody [EP5699] (ab133590)


IRF4 Flow Cytometry (Intracellular) staining using rabbit Anti-IRF4 antibody

Overlay histogram showing Ramos cells stained with ab133590 (red line). The cells were fixed with 4% paraformaldehyde (10 min) and then permeabilized with 0.1% PBS-Tween for 20 min. The cells were then incubated in 1x PBS / 10% normal goat serum / 0.3M glycine to block non-specific protein-protein interactions followed by the antibody (ab133590, 1/1000 dilution) for 30 min at 22°C. The secondary antibody used was Alexa Fluor® 488 goat anti-rabbit IgG (H&L) (ab150077) at 1/2000 dilution for 30 min at 22°C. Isotype control antibody (black line) was rabbit IgG (monoclonal) (0.1µg/1x10^6 cells) used under the same conditions. Unlabelled sample (blue line) was also used as a control. Acquisition of >5,000 events were collected using a 20mW Argon ion laser (488nm) and 525/30 bandpass filter. This antibody gave a positive signal in Ramos cells fixed with 80% methanol (5 min)/permeabilized with 0.1% PBS-Tween for 20 min used under the same conditions.



Immunocytochemistry/ Immunofluorescence - Anti-IRF4 antibody [EP5699] (ab133590)

Immunocytochemistry/ Immunofluorescence analysis of IM-9 (human multiple myeloma B lymphoblast) labeling IRF4 with ab133590 at 1/500 dilution followed by ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed at 1/1000 dilution. Cells were fixed with 4% Paraformaldehyde and permeabilised with 0.1% TritonX-100. DAPI was used as nuclear counterstain. ab195889 Anti-alpha Tubulin mouse monoclonal antibody - Microtubule Marker (Alexa Fluor® 594) at 1/200 was used as the counterstain antibody. Confocal image showing mainly nuclear staining in IM-9 cell line. Negative control: 293T (PMID: 23804715) Image was taken with a confocal microscope(Leica-Microsystems, TCS SP8).



Immunoprecipitation - Anti-IRF4 antibody [EP5699] (ab133590)

IRF4 was immunoprecipitated from 0.35 mg IM-9 (Human multiple myeloma B Lymphoblast) whole cell lysate 10 µg with 133590 at 1/20 dilution (1µg). VeriBlot for IP Detection Reagent (HRP) (ab131366) was used at 1/5000 dilution.
Lane 1: IM-9 (Human multiple myeloma B Lymphoblast) whole cell lysate 10 µg

Lane 2: ab133590 IP in IM-9 whole cell lysate

Lane 3: Rabbit monoclonal IgG (ab172730) instead of ab133590 in IM-9 whole cell lysate


Blocking and dilution buffer and concentration: 5% NFDM/TBST.

Lower bands could be ubiquitin mediated protein degradation. (PMID: 23658517)

All lanes:
Immunoprecipitation - Anti-IRF4 antibody [EP5699] (ab133590)

Predicted band size: 52 kDa

Observed band size: 52 kDa



Western blot - Anti-IRF4 antibody [EP5699] (ab133590)

All lanes:
Western blot - Anti-IRF4 antibody [EP5699] (ab133590) at 1/10000 dilution

Lane 1:
Namalwa cell lysate at 10 µg

Lane 2:
HuT78 cell lysate at 10 µg

Lane 3:
IM9 cell lysate at 10 µg

Secondary

All lanes:
HRP labelled goat anti-rabbit at 1/2000 dilution

Predicted band size: 52 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.