abcam

ab133611

Anti-SRC3 antibody [EPR4374(3)]

Recombinant | RabMAb | KO Validated

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR4374(3)
Purification technique	Affinity purification Protein A
Concentration	1.914 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

IHC-P

Tested

Species	Human
Dilution info	1/50 - 1/100
Notes	-

ICC/IF

Tested

Species	Human
Dilution info	1/50 - 1/10000
Notes	-

IP

Not recommended

Species	Human
Dilution info	-
Notes	-

WB

Tested

Species	Human
Dilution info	1/1000 - 1/10000
Notes	-

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	1/100 - 1/500
Notes	ab172730 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.

Target data

See full target information NCOA3

Function	Nuclear receptor coactivator that directly binds nuclear receptors and stimulates the transcriptional activities in a hormone-dependent fashion. Plays a central role in creating a multisubunit coactivator complex, which probably acts via remodeling

of chromatin. Involved in the coactivation of different nuclear receptors, such as for steroids (GR and ER), retinoids (RARs and RXRs), thyroid hormone (TRs), vitamin D3 (VDR) and prostanoids (PPARs). Displays histone acetyltransferase activity. Also involved in the coactivation of the NF-kappa-B pathway via its interaction with the NFKB1 subunit.

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Storage information	Stable for 12 months at -20°C

Notes

Species reactivity
Mouse, Rat: We have preliminary internal testing data to indicate this antibody may not react with these species. Please contact us for more information.

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended

We do not recommend this combination. It is not covered by our product promise.


We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

---


5 product images



Immunocytochemistry/ Immunofluorescence - Anti-SRC3 antibody [EPR4374(3)] (ab133611)

SRC3 Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-SRC3 antibody


ab133611 staining NCOA3 in MCF-7 (positive) and MCF-7-NCOA3 KO (negative) cells (ab275827). The cells were fixed with 4% Paraformaldehyde (10 min), permeabilized with 0.1% PBS-Triton X-100 for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1%PBS-Tween for 1h. The cells were then incubated for 2 hours at room temperature with ab133611 at 0.2 µg/ml (shown in green) and ab7291, Anti-alpha Tubulin antibody [DM1A] - Loading Control (shown in pseudocolour magenta). Cells were then incubated with ab150081, Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed at 1/1000 dilution and ab150120, Goat Anti-Mouse IgG H&L (Alexa Fluor® 594) preadsorbed at 1/1000 dilution for 1 hour at room temperature. Nuclear DNA was labeled with DAPI (shown in blue). Also suitable in cells fixed with 100% Methanol (5 min). Image was acquired with a high-content analyser (Operetta CLS, Revvity) and a maximum intensity projection of confocal sections is shown.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-SRC3 antibody [EPR4374(3)] (ab133611)

SRC3 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-SRC3 antibody


Immunohistochemical analysis of KAT13B / AIB1 in paraffin embedded Human endometrium adenocarcinoma tissue using ab133611 at a dilution of 1/50.
Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.



Immunocytochemistry/ Immunofluorescence - Anti-SRC3 antibody [EPR4374(3)] (ab133611)

SRC3 Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-SRC3 antibody


Immunocytochemistry analysis of HeLa (Human cervix adenocarcinoma epithelial cell) labeling SRC3 with purified ab133611 at 1/100 dilution. Cells were fixed with 4% Paraformaldehyde and permeabilised with 0.1% tritonX-100. Goat anti rabbit IgG (Alexa Fluor® 488, ab150077) at 1/1000 (2 µg/ml) was used as the secondary antibody. ab195889 Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor® 594) 1/200 (2.29 µg/ml) was used as counterstain. Nuclei were stained blue with DAPI.
Negative control: PBS instead of the primary antibody.



Flow Cytometry (Intracellular) - Anti-SRC3 antibody [EPR4374(3)] (ab133611)

SRC3 Flow Cytometry (Intracellular) staining using rabbit Anti-SRC3 antibody

Overlay histogram showing HeLa cells stained with ab133611 (red line). The cells were fixed with 80% methanol (5 min) and then permeabilized with 0.1% PBS-Tween for 20 min. The cells were then incubated in 1x PBS / 10% normal goat serum / 0.3M glycine to block non-specific protein-protein interactions followed by the antibody (ab133611 1/100 dilution) for 30 min at 22°C. The secondary antibody used was Alexa Fluor® 488 goat anti-rabbit IgG (H&L) (ab150077) at 1/2000 dilution for 30 min at 22°C. Isotype control antibody (black line) was rabbit IgG (monoclonal) (1µg/1x10^6 cells) used under the same conditions. Unlabelled sample (blue line) was also used as a control. Acquisition of >5000 events were collected using a 20mW Argon ion laser (488nm) and 525/30 bandpass filter.



Western blot - Anti-SRC3 antibody [EPR4374(3)] (ab133611)

All lanes:
Western blot - Anti-SRC3 antibody [EPR4374(3)] (ab133611) at 1/1000 dilution

Lane 1:
Daudi cell lysate at 10 µg

Lane 2:
HeLa cell lysate at 10 µg

Lane 3:
K562 cell lysate at 10 µg

Secondary

All lanes:
HRP labelled goat anti-rabbit at 1/2000 dilution

Predicted band size: 155 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.