abcam

ab133734

Anti-PTBP1 antibody [EPR9048(B)]

Anti-PTBP1 antibody [EPR9048(B)] (ab133734) is a rabbit monoclonal antibody detecting PTBP1 in **Western Blot, Flow Cytometry (Intra), IHC-P, ICC/IF**. Suitable for **Human, Mouse**.

- KO validated for confirmed specificity
- Biophysical QC for unrivalled batch-batch consistency
- Over 20 publications

Recombinant | RabMAb | KO Validated | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 50% Tissue culture supernatant, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR9048(B)
Purification technique	Affinity purification Protein A
Concentration	0.118 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

WB

Tested

Species	Human
Dilution info	1/10000 - 1/50000
Notes	-

Expected

Species	Mouse
Dilution info	1/10000 - 1/50000
Notes	-

Predicted

Species	Rat
Dilution info	-
Notes	-

IHC-P

Tested

Species	Human
Dilution info	1/100 - 1/250
Notes	Perform antigen retrieval before commencing with IHC staining protocol.

Expected

Species	Mouse
Dilution info	1/100 - 1/250
Notes	Perform antigen retrieval before commencing with IHC staining protocol.

Predicted

Species	Rat
Dilution info	-
Notes	-

ICC/IF

Tested

Species	Human
Dilution info	1/250 - 1/500
Notes	-

Expected

Species	Mouse
Dilution info	1/250 - 1/500
Notes	-

Predicted

Species	Rat
Dilution info	-
Notes	-

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	1/2000
Notes	-

Expected

Species	Mouse
Dilution info	1/2000
Notes	-

Predicted

Species	Rat

Dilution info	-
Notes	-

IP

Not recommended

Species	Mouse, Human, Rat
Dilution info	-
Notes	-

Target data

See full target information PTBP1

Function	Plays a role in pre-mRNA splicing and in the regulation of alternative splicing events. Activates exon skipping of its own pre-mRNA during muscle cell differentiation. Binds to the polypyrimidine tract of introns. May promote RNA looping when bound to two separate polypyrimidine tracts in the same pre-mRNA. May promote the binding of U2 snRNP to pre-mRNA. Cooperates with RAVER1 to modulate switching between mutually exclusive exons during maturation of the TPM1 pre-mRNA. Represses the splicing of MAPT/Tau exon 10 (PubMed:15009664). Binds to polypyrimidine-rich controlling element (PCE) of CFTR and promotes exon skipping of CFTR exon 9, thereby antagonizing TIA1 and its role in exon inclusion of CFTR exon 9 (PubMed:14966131). Plays a role in the splicing of pyruvate kinase PKM by binding repressively to a polypyrimidine tract flanking PKM exon 9, inhibiting exon 9 inclusion and resulting in exon 10 inclusion and production of the PKM M2 isoform (PubMed:20010808). In case of infection by picornaviruses, binds to the viral internal ribosome entry site (IRES) and stimulates the IRES-mediated translation (PubMed:21518806).

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

What is this antibody validated in?
Anti-PTBP1 antibody [EPR9048(B)] (ab133734) is a rabbit recombinant monoclonal antibody and is validated for use in Western Blot (WB), Flow Cytometry (Intra), Immunohistochemistry (IHC-P), Immunocytochemistry/immunofluorescence (ICC/IF) in Human, Mouse samples.

What is the molecular weight of PTBP1?
Anti-PTBP1 [EPR9048(B)] (ab133734) specifically detects a band for PTBP1 (UniProt: P26599) at a molecular weight of 57kDa.

Trusted by the scientific community
Anti-PTBP1 [EPR9048(B)] (ab133734) was first used in a scientific publication in 2012 and has been cited over 20 times in peer-reviewed journals.

Specificity confirmed
The specificity of Anti-PTBP1 antibody [EPR9048(B)] (ab133734) has been confirmed by Western blot testing in PTBP1 Knockout HAP1 cell line, ab265155.

Other related products
We have a range of other formats of antibody clone [EPR9048(B)] also available for your convenience: ab133734, Alexa Fluor® 647 - ab201020, Carrier free - ab240079, PE - ab305838, APC - ab305839, HRP - ab305840, Alkaline Phosphatase - ab308831, Alexa Fluor® 488 - ab309768, Alexa Fluor® 594 - ab310562, Alexa Fluor® 555 - ab312091, Alexa Fluor® 568 - ab312571, Alexa Fluor® 750 - ab321066

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Trial sizes available!
Test your antibody or perform pre-screening before committing to a larger quantity. Sold in 20µl. Discover our selection of trial-size antibodies.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	PTBP1 also known as polypyrimidine tract-binding protein 1 or hnRNP I is a protein involved in RNA binding. It weighs approximately 57 kDa and is known for its expression in diverse cell types especially in the nucleus of eukaryotic cells. PTBP1

	contains four RNA recognition motifs which allow it to interact with polypyrimidine-rich sequences in pre-mRNA. This interaction plays an important role in RNA splicing and the regulation of alternative splicing events.
Biological function summary	PTBP1 influences the stability transport and translation of mRNA molecules. It is part of the heterogeneous nuclear ribonucleoprotein complex which plays a fundamental role in the processing and modification of pre-mRNA into mature mRNA. This activity impacts the expression of a wide range of genes affecting cellular function and growth. PTBP1 regulates mRNA metabolism contributing significantly to post-transcriptional gene regulation.
Pathways	The involvement of PTBP1 spans the management of alternative splicing and the modulation of gene expression. It participates importantly in the alternative splicing pathway alongside proteins like PTBP2 which perform overlapping functions in neurons. PTBP1 affects exon inclusion or exclusion influencing the genetic diversity presented in protein forms within cells. It also plays a role in the exon junction complex pathway impacting post-translational modifications and protein interactions.
Associated diseases and disorders	Variations in PTBP1 function are linked to conditions such as cancer and neurological disorders. Altered expression levels or mutations can contribute to carcinogenesis through the misregulation of splice variants specific to oncogenes and tumor suppressor genes. In neurological disorders PTBP1 interacts with proteins like FUS affecting the regulation of neuronal-specific alternative splicing. Misbalance in these interactions can have significant pathological implications contributing to diseases such as amyotrophic lateral sclerosis (ALS) and glioblastomas.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.


7 product images



Flow Cytometry (Intracellular) - Anti-PTBP1 antibody [EPR9048(B)] (ab133734)

PTBP1 Flow Cytometry (Intracellular) staining using rabbit Anti-PTBP1 antibody


Intracellular Flow Cytometry analysis of A549 (Human lung carcinoma epithelial cell) cells labeling PTBP1 (red) with purified ab133734 at a 1/2000 dilution (1ug/mL). Cells were fixed with 4% paraformaldehyde and permeabilized with 90% methanol. A goat anti rabbit IgG (Alexa Fluor® 488) (ab150077) was used as the secondary antibody at a 1/2000 dilution. Black - Rabbit monoclonal IgG (Black) (ab172730). Blue (unlabeled control) - Cell without incubation with primary antibody and secondary antibody (Blue).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-PTBP1 antibody [EPR9048(B)] (ab133734)

PTBP1 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-PTBP1 antibody


Immunohistochemical analysis of PTBP1 in paraffin embedded Human breast carcinoma tissue stained with ab133734 at a 1/100 dilution.
Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.



Immunocytochemistry/ Immunofluorescence - Anti-PTBP1 antibody [EPR9048(B)] (ab133734)

PTBP1 Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-PTBP1 antibody

Immunofluorescent staining of PTBP1 in A549 cells, using ab133734 at a 1/250 dilution.



Western blot - Anti-PTBP1 antibody [EPR9048(B)] (ab133734)

All lanes:
Western blot - Anti-PTBP1 antibody [EPR9048(B)] (ab133734) at 1/10000 dilution

Lane 1:
Human fetal liver lysate at 10 µg

Lane 2:
Jurkat cell lysate at 10 µg

Lane 3:
Daudi cell lysate at 10 µg

Lane 4:


A549 cell lysate at 10 µg

Lane 5:
BxPC3 cell lysate at 10 µg

Secondary

All lanes:
Goat anti-rabbit HRP conjugated antibody at 1/2000 dilution

Predicted band size: 57 kDa



Western blot - Anti-PTBP1 antibody [EPR9048(B)] (ab133734)


Lanes 1 - 2: Merged signal (red and green). Green - ab133734 observed at 57 kDa. Red - loading control, ab9484, observed at 37 kDa.
ab133734 was shown to recognize PTBP1 in wild-type HAP1 cells as signal was lost at the expected MW in PTBP1 knockout cells. Additional cross-reactive bands were observed in the wild-type and knockout cells. Wild-type and PTBP1 knockout samples were subjected to SDS-PAGE. ab133734 and ab9484 (Mouse anti-GAPDH loading control) were incubated overnight at 4°C at 1/10000 dilution and 1/20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preabsorbed ab216773 and Goat anti-Mouse IgG H&L (IRDye® 680RD) preabsorbed ab216776 secondary antibodies at 1/10000 dilution for 1 hour at room temperature before imaging.

All lanes:
Western blot - Anti-PTBP1 antibody [EPR9048(B)] (ab133734) at 1/1000 dilution

Lane 1:
Wild-type HAP1 whole cell lysate at 20 µg

Lane 2:
PTBP1 knockout HAP1 whole cell lysate at 20 µg

Predicted band size: 57 kDa



Western blot - Anti-PTBP1 antibody [EPR9048(B)] (ab133734)

PTBP1 Western blot staining using rabbit Anti-PTBP1 antibody

Lanes 1- 2: Merged signal (red and green). Green - ab133734 observed at 57 kDa. Red - Anti-GAPDH antibody [6C5] - Loading Control (ab8245) observed at 37 kDa.
ab133734 was shown to react with PTBP1 in wild-type HeLa cells in western blot. Loss of signal was observed when knockout cell line ab265155 (knockout cell lysate ab257614) was used. Wild-type HeLa and PTBP1 knockout HeLa cell lysates were subjected to SDS-PAGE. Membrane was blocked for 1 hour at room temperature in 0.1% TBST with 3% non-fat dried milk. ab133734 and Anti-GAPDH antibody [6C5] - Loading Control (ab8245) overnight at 4°C at a 1 in 10000 dilution and a 1 in 20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye®800CW) preadsorbed (ab216773) and Goat anti-Mouse IgG H&L (IRDye®680RD) preadsorbed (ab216776) secondary antibodies at 1 in 20000 dilution for 1 hour at room temperature before imaging.

All lanes:
Western blot - Anti-PTBP1 antibody [EPR9048(B)] (ab133734) at 1/10000 dilution

Lane 1:
Wild-type HeLa cell lysate at 20 µg

Lane 2:
Western blot - Human PTBP1 knockout HeLa cell lysate (ab257614) at 20 µg

Performed under reducing conditions.

Predicted band size: 57 kDa

Observed band size: 57 kDa




Image collected and cropped by CiteAb under a CC-BY license from the publication

Immunocytochemistry/ Immunofluorescence - Anti-PTBP1 antibody [EPR9048(B)] (ab133734)

PTBP1 Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-PTBP1 antibody

Immunocytochemistry-immunofluorescence using Anti-PTBP1 antibody [EPR9048(B)], ab133734. Publication image from Makeyev, E. V. et al., 2022, Mol Cell, 34741808. Legend direct from paper.

Hybridization-proximity labeling technology for genetically unperturbed samples(A) Top: recombinant HyPro enzyme containing the APEX2 and digoxigenin-binding domains connected by a flexible linker. Bottom: HyPro-labeling approach to identify proteins and RNAs colocalizing with an RNA target of interest.(B) SDS-PAGE analysis of purified HyPro protein peak eluted from a size exclusion chromatography column.(C) Peroxidase activity assay of purified HyPro enzyme.(D) Spot assay showing that HyPro enzyme can bind digoxigenin-labeled DNA while retaining the peroxidase activity.(E) Workflows developed to identify proteins (HyPro-MS) and RNAs (HyPro-seq) colocalized with a transcript of interest and to validate specificity of digoxigenin-labeled probes (HyPro-FISH).(F–H) HyPro-FISH analyses of (F) the 45S pre-rRNA, (G) NEAT1, and (H) the (UC)n-repeated part of PNCTR combined with immunofluorescence detection of nucleolar (No) marker FBL, paraspeckle (Ps) marker SFPQ, and PNC marker PTBP1. As expected, FBL but not 45S is detectable in Cajal bodies (Cb). Image acquisition settings were adjusted according to RNA bait abundance. All images in (F)–(H) are maximum-intensity z stacks. Scale bars, 5 µm.See also Figure S1.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.