abcam

ab134113

Anti-Niemann Pick C1 antibody [EPR5209]

Anti-Niemann Pick C1 antibody [EPR5209] (ab134113) is a rabbit monoclonal antibody detecting Niemann Pick C1 in **Western Blot, Flow Cytometry (Intra), IHC-P, ICC/IF**. Suitable for **Human, Mouse, Rat**.

- KO validated for confirmed specificity
- Biophysical QC for unrivalled batch-batch consistency
- Over 50 publications

Recombinant | RabMAb | KO Validated | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR5209
Purification technique	Affinity purification Protein A
Dissociation constant	4.9×10^{-11} M
Concentration	0.382 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

IHC-P

Tested

Species	Mouse
Dilution info	1/50 - 1/100
Notes	Perform heat-mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.

Species	Rat
Dilution info	1/50 - 1/100
Notes	Perform heat-mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.

Species	Human
Dilution info	1/50 - 1/100
Notes	Perform heat-mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.

WB

Tested

Species	Mouse
Dilution info	1/2000 - 1/10000
Notes	-

Species	Human
Dilution info	1/2000 - 1/10000
Notes	-

Expected

Species	Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

ICC/IF

Tested

Species	Mouse
Dilution info	1/70
Notes	-

Expected

Species	Rat, Human
Dilution info	Use at an assay dependent concentration.
Notes	-

Flow Cyt (Intra)

Tested

Species	Mouse
Dilution info	1/200
Notes	ab172730 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.

Expected

Species	Rat, Human
Dilution info	Use at an assay dependent concentration.
Notes	-

Target data

See full target information NPC intracellular cholesterol transporter 1

Function	Intracellular cholesterol transporter which acts in concert with NPC2 and plays an important role in the egress of cholesterol from the endosomal/lysosomal compartment (PubMed:10821832, PubMed:12554680, PubMed:18772377, PubMed:27238017, PubMed:9211849, PubMed:9927649). Unesterified cholesterol that has been released from LDLs in the lumen of the late endosomes/lysosomes is transferred by NPC2 to the cholesterol-binding pocket in the N-terminal domain of NPC1 (PubMed:18772377, PubMed:19563754, PubMed:27238017, PubMed:27378690, PubMed:28784760, PubMed:9211849, PubMed:9927649). Cholesterol binds to NPC1 with the hydroxyl group buried in the binding pocket (PubMed:19563754). Binds oxysterol with higher affinity than cholesterol. May play a role in vesicular trafficking in glia, a process that may be crucial for maintaining

the structural and functional integrity of nerve terminals (Probable). Inhibits cholesterol-mediated mTORC1 activation throught its interaction with SLC38A9 (PubMed:28336668). (Microbial infection) Acts as an endosomal entry receptor for ebolavirus.

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Storage information	Stable for 12 months at -20°C

Notes

Product Specifications
Anti-Niemann Pick C1 antibody [EPR5209] (ab134113) is a rabbit recombinant monoclonal antibody and is validated for use in Flow Cyt (Intra), ICC/IF, IHC-P, WB in human, mouse and rat samples.
Anti-Niemann Pick C1 antibody [EPR5209] (ab134113) specifically detects Niemann Pick C1 (UniProt ID: O15118; Molecular weight: 140kDa) and is sold in 100 µL and 1 mL selling sizes.

Quality and Validation
Abcam's high quality manufacturing and validation processes ensure Anti-Niemann Pick C1 antibody [EPR5209] (ab134113) has high sensitivity and specificity alongside high lot-to-lot consistency and reproducibility.
The specificity of Anti-Niemann Pick C1 antibody [EPR5209] (ab134113) has been confirmed by testing in knockout samples.
Anti-Niemann Pick C1 antibody [EPR5209] (ab134113) has been cited over 51 times in peer reviewed journals and is trusted by the scientific community.

Related Products
Conjugation-ready, carrier free format available for antibody clone EPR5209 - ab224268.
Antibody clone EPR5209 is also available pre-conjugated to a variety of labels for your convenience - APC, PE, Alexa Fluor® 488, Alexa Fluor® 594, Alexa Fluor® 555, Alexa Fluor® 647, Alexa Fluor® 750 (ab223985, ab317903, ab317983, ab318025, ab318067, ab318108, ab321075).

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply

- Animal-free batch production

For more information, read more on recombinant antibodies.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	Niemann Pick C1 also known as NPC1 or C1 protein is an integral membrane protein involved in cholesterol and lipid transport within cells. It is located in late endosomes and lysosomes and plays a role in intracellular cholesterol traffic. The NPC1 protein has a molecular weight of approximately 170 kDa. It is expressed in various tissues with high expression levels in the liver brain and spleen.
Biological function summary	This protein acts in the movement of lipids and cholesterol through cellular compartments. NPC1 functions as part of a protein complex that includes NPC2 which works to mobilize cholesterol from lysosomes to other areas in the cell where it is further processed or stored. This transport mechanism is necessary to maintain cellular lipid homeostasis and proper cellular functions.
Pathways	NPC1 integrates into the cholesterol trafficking pathway and plays an essential part in intracellular lipid regulation. It interacts with proteins involved in cholesterol metabolism such as sterol regulatory element-binding proteins (SREBPs). These interactions ensure that cholesterol distribution within the cell remains balanced and adjusts to metabolic needs. The proper function of NPC1 in these pathways is necessary for maintaining cell health and homeostasis.
Associated diseases and disorders	NPC1 mutations are directly linked to Niemann-Pick disease type C a severe lipid storage disorder. This condition is characterized by the accumulation of cholesterol and other lipids in lysosomes leading to neurological and hepatic dysfunction. NPC1's role in this disease involves disrupted cholesterol trafficking which is critical for cellular lipid homeostasis. Additionally NPC2 protein is also affected in this disorder as both work together in the cholesterol transport process.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

13 product images



Immunocytochemistry/ Immunofluorescence - Anti-Niemann Pick C1 antibody [EPR5209] (ab134113)

Niemann Pick C1 Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-Niemann Pick C1 antibody

Immunofluorescent analysis of 4% paraformaldehyde-fixed, 0.1% Triton X-100 permeabilized Neuro-2a (mouse neuroblastoma cell line) cells labeling Niemann Pick C1 with ab134113 at 1/70 followed by Goat anti-rabbit IgG (Alexa Fluor® 488) (ab150077) secondary antibody at 1/500 (green). Confocal image showing cytoplasmic staining on Neuro-2a cell line. The nuclear counterstain is DAPI (blue). Tubulin is detected with ab7291 (anti-Tubulin mouse mAb) at 1/1000 and ab150120 (Alexa Fluor® 594 Goat anti-Mouse secondary) at 1/500 (red).
The negative controls are as follows:
-ve control 1 – ab134113 at 1/70 followed by ab150120 (Alexa Fluor® 594 Goat anti-Mouse secondary) at 1/500.
-ve control 2 - ab7291 (anti-Tubulin mouse mAb) at 1/1000 followed by ab150077 (Alexa Fluor® 488 Goat Anti-Rabbit IgG H&L) at 1/500.





Image from Cermak S et al., PLoS One. 2016;11(11):e0167428. Fig 3.; doi: 10.1371/journal.pone.0167428. Reproduced under the Creative Commons license http://creativecommons.org/licenses/by/4.0/

Immunocytochemistry/ Immunofluorescence - Anti-Niemann Pick C1 antibody [EPR5209] (ab134113)


Cathepsin B/L inhibition causes NPC disease-like cholesterol accumulation in SH-SY5Y cells. Confocal microscopy of SH-SY5Y control and PADK treated cells. Cholesterol (filipin staining, white) and NPC1 (ab134113; green).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Niemann Pick C1 antibody [EPR5209] (ab134113)

Niemann Pick C1 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-Niemann Pick C1 antibody


Immunohistochemical analysis of paraffin embedded human kidney tissue labelling Niemann Pick C1 with unpurified ab134113 at 1/50.
Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Niemann Pick C1 antibody [EPR5209] (ab134113)

Niemann Pick C1 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-Niemann Pick C1 antibody


Immunohistochemical staining of paraffin embedded human liver with purified ab134113 at a working dilution of 1 in 50. The secondary antibody used is a HRP polymer for rabbit IgG. The sample is counter-stained with hematoxylin. Antigen retrieval was perfomed using Tris-EDTA buffer, pH 9.0. PBS was used instead of the primary antibody as the negative control, and is shown in the inset.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Niemann Pick C1 antibody [EPR5209] (ab134113)

Niemann Pick C1 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-Niemann Pick C1 antibody

Immunohistochemical staining of paraffin embedded mouse liver with purified ab134113 at a working dilution of 1 in 50. The secondary antibody used is a HRP polymer for rabbit IgG. The sample is counter-stained with hematoxylin. Antigen retrieval was perfomed using Tris-EDTA buffer, pH 9.0. PBS was used instead of the primary antibody as the negative control, and is shown in the inset.




Flow Cytometry (Intracellular) - Anti-Niemann Pick C1 antibody [EPR5209] (ab134113)

Niemann Pick C1 Flow Cytometry (Intracellular) staining using rabbit Anti-Niemann Pick C1 antibody

ab134113 staining Niemann Pick C1 in Neuro-2a (mouse neuroblastoma cell line) by intracellular flow cytometry. Cells were fixed with 4% paraformaldehyde, permabilised with 90% methanol and the sample was incubated with the primary antibody at a dilution of 1/200. A goat anti rabbit IgG (Alexa Fluor® 488) at a dilution of 1/2000 was used as the secondary antibody.
Isoytype control: Rabbit monoclonal IgG (Black)


Unlabeled control: Cell without incubation with primary antibody and secondary antibody (Blue)



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Niemann Pick C1 antibody [EPR5209] (ab134113)

Niemann Pick C1 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-Niemann Pick C1 antibody

Immunohistochemical staining of paraffin embedded rat cerebellum with purified ab134113 at a working dilution of 1 in 50. The secondary antibody used is a HRP polymer for rabbit IgG. The sample is counter-stained with hematoxylin. Antigen retrieval was perfomed using Tris-EDTA buffer, pH 9.0. PBS was used instead of the primary antibody as the negative control, and is shown in the inset.



Schultz et al Nat Commun. 2018; 9: 3671. Published online 2018 Sep 10. doi: 10.1038/s41467-018-06115-2. Reproduced under the Creative Commons Licence http://creativecommons.org/licenses/by/4.0/.


Immunocytochemistry/ Immunofluorescence - Anti-Niemann Pick C1 antibody [EPR5209] (ab134113)

Niemann Pick C1 Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-Niemann Pick C1 antibody

I1061T NPC1 traffics to autophagosomes
I1061T NPC1 fibroblasts were treated with vehicle or 100 nM bafilomycin A1 (Baf) for 24 h. Fixed cells were stained for LC3 (red), NPC1 (green), and DAPI (blue) then imaged by confocal microscopy. Co-localization is indicated by yellow color in the merged image. Scale bar = 25 µm.

Niemann Pick C1 (NPC1) was detected using ab134113 at 1/200 dilution.

From Figure 4b of Schultz et al.



Western blot - Anti-Niemann Pick C1 antibody [EPR5209] (ab134113)

Niemann Pick C1 Western blot staining using rabbit Anti-Niemann Pick C1 antibody

Lanes 1 - 4: Merged signal (red and green). Green - ab134113 observed at 180 kDa. Red - loading control, ab9484, observed at 37 kDa.
ab134113 was shown to specifically react with Niemann Pick C1 in wild-type HAP1 cells as signal was lost in NPC1 (Niemann Pick C1) knockout cells. Wild-type and NPC1 (Niemann Pick C1) knockout samples were subjected to SDS-PAGE. ab134113 and ab9484 (Mouse anti-GAPDH loading control) were incubated overnight at 4°C at 1/1000 dilution and 1/20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preabsorbed ab216773 and Goat anti-Mouse IgG H&L (IRDye® 680RD) preabsorbed ab216776 secondary antibodies at 1/20000 dilution for 1 hour at room temperature before imaging.

All lanes:
Western blot - Anti-Niemann Pick C1 antibody [EPR5209] (ab134113) at 1/1000 dilution


Lane 1:
Wild-type HAP1 whole cell lysate at 20 µg

Lane 2:
NPC1 (Niemann Pick C1) knockout HAP1 whole cell lysate at 20 µg

Lane 3:
HEK293 whole cell lysate at 20 µg

Lane 4:
HepG2 whole cell lysate at 20 µg

Predicted band size: 142 kDa



Western blot - Anti-Niemann Pick C1 antibody [EPR5209] (ab134113)

Niemann Pick C1 Western blot staining using rabbit Anti-Niemann Pick C1 antibody

Blocking buffer: 5% NFDM/TBST
Dilution buffer: 5% NFDM/TBST

All lanes:
Western blot - Anti-Niemann Pick C1 antibody [EPR5209] (ab134113) at 1/10000 dilution

Lane 1:
HepG2 (human liver hepatocellular carcinoma cell line) cell lysate at 20 µg

Lane 2:
THP-1 (human monocytic leukemia cell line) cell lysate at 20 µg

Lane 3:
HEK-293 (human epithelial cell line from embryonic kidney) cell lysate at 20 µg


Lane 4:
PC-3 (human prostate adenocarcinoma cell line) cell lysate at 20 µg

Secondary

All lanes:
HRP goat anti-rabbit (H+L) at 1/1000 dilution

Predicted band size: 142 kDa

Observed band size: 180 kDa



Western blot - Anti-Niemann Pick C1 antibody [EPR5209] (ab134113)

All lanes:
Western blot - Anti-Niemann Pick C1 antibody [EPR5209] (ab134113) at 1/1000 dilution

Lane 1:
3T3 L1 cell lysate at 10 µg

Lane 2:
L6 (rat skeletal muscle cell line) cell lysate at 10 µg

Lane 3:
HepG2 (human liver hepatocellular carcinoma cell line) cell lysate at 10 µg

Lane 4:
THP-1 (human monocytic leukemia cell line) cell lysate at 10 µg

Lane 5:
HEK-293T (human epithelial cell line from embryonic kidney transformed with large T antigen) cell lysate at 10 µg

Lane 6:
PC-3 (human prostate adenocarcinoma cell line) cell lysate at 10 µg

Lane 7:
Rat liver lysate at 10 µg


Lane 8:
Rat brain lysate at 10 µg

Secondary

All lanes:
Goat anti-rabbit HRP conjugated antibody at 1/2000 dilution

Predicted band size: 142 kDa

Observed band size: 180 kDa



Western blot - Anti-Niemann Pick C1 antibody [EPR5209] (ab134113)

Niemann Pick C1 Western blot staining using rabbit Anti-Niemann Pick C1 antibody

Blocking buffer: 5% NFDM/TBST
Dilution buffer: 5% NFDM/TBST

All lanes:
Western blot - Anti-Niemann Pick C1 antibody [EPR5209] (ab134113) at 1/10000 dilution

Lane 1:
3T3-L1 cell lysate at 20 µg

Lane 2:
L6 (rat skeletal muscle cell line) cell lysate at 20 µg


Lane 3:
Rat tissue lysate at 20 µg

Secondary

All lanes:
HRP goat anti-rabbit (H+L) at 1/1000 dilution

Predicted band size: 142 kDa

Observed band size: 180 kDa



OI-RD Scanning - Anti-Niemann Pick C1 antibody [EPR5209] (ab134113)

We have systematically measured KD (the equilibrium dissociation constant between the antibody and its antigen), of more than 840 recombinant antibodies to assess not only their individual KD values but also to see the average affinity of antibody. Based on the comparison with published literature values for mouse monoclonal antibodies, Recombinant antibodies appear to be on average 1-2 order of magnitude higher affinity.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.