abcam

ab134399

APC Anti-CD38 antibody [HIT2]

Anti-CD38 antibody [HIT2] conjugated to APC - suitable for flow cytometry, and reacts with human samples. Clone [HIT2] is a mouse monoclonal antibody and one of the most commonly used clones for flow cytometry.

Design your panel with our interactive flow cytometry panel builder. Click here.

Key facts

Isotype	IgG1
Host species	Mouse
Conjugation	APC
Excitation/Emission	Ex: 650nm, Em: 660nm
Storage buffer	pH: 7.4 Preservative: 0.1% Sodium azide Constituents: PBS, 0.2% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	Cell preparation containing CD38 protein. The exact immunogen used to generate this antibody is proprietary information. Database link P28907
Clone number	HIT2
Purification technique	Size-exclusion chromatography

Reactivity data

Flow Cyt

Tested

Species	Human
Dilution info	-

Notes	10 µl reagent / 100 µl of whole blood or 10^6 cells in a suspension.

Target data

See full target information CD38

Function	Multifunctional transmembrane glycoprotein able to exert enzymatic activities and also to mobilize calcium, to transduce signals, to adhere to hyaluronan and to other ligands. Synthesizes cyclic ADP-ribose (cADPR), a second messenger for glucose-induced insulin secretion (PubMed:7961800, PubMed:8253715). Synthesizes the Ca(2+) mobilizer nicotinate-adenine dinucleotide phosphate, NAADP(+), from 2'-phospho-cADPR and nicotinic acid, as well as from NADP(+) and nicotinic acid. At both pH 5.0 and pH 7.4 preferentially transforms 2'-phospho-cADPR into NAADP(+), while preferentially cleaving NADP(+) to cADPR and ADPRP rather than into NADDP(+) (PubMed:16690024). Has cADPR hydrolase activity (PubMed:7961800, PubMed:8253715). Functions also as a receptor that binds the ligand CD31 on endothelial cells, promoting lymphocyte activation, proliferation, and migration across the endothelial barrier (PubMed:9551996). Involved in the regulation of crucial dendritic cell functions acquired at the mature stage, such as CCL21-driven migration, survival, and Th1-polarizing activity (PubMed:16293598). In lamina propria T lymphocytes, CD38/CD31 cognate interactions initiate a multistep signaling pathway resulting in activation of LCK anf LAT, followed by cytokine release (PubMed:11259373).

Storage

Shipped at conditions	Blue Ice
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	+4°C

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended

We do not recommend this combination. It is not covered by our product promise.


We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

---


2 product images



Flow Cytometry - APC Anti-CD38 antibody [HIT2] (ab134399)

CD38 Flow Cytometry staining using mouse Anti-CD38 antibody

Flow cytometry analysis (surface staining) of human peripheral blood labeling CD38 with ab134399.



Flow Cytometry - APC Anti-CD38 antibody [HIT2] (ab134399)

CD38 Flow Cytometry staining using mouse Anti-CD38 antibody

Flow cytometry analysis (surface staining) of human peripheral blood labeling CD38 with ab134399.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.