abcam

ab140125

Anti-GNAI1 antibody [EPR9441(B)]

Recombinant | RabMAb

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR9441(B)
Purification technique	Affinity purification Protein A
Specificity	The immunogenic sequence of ab140125 has 92.9% homology with GNAI3 and 64% (4 continuous residues) homology with GNAI2 which might cause cross reactivity with GNAI3 and GNAI2.
Concentration	Concentration data is not available right now, please try again later Batch concentration finder →

Reactivity data

IHC-P

Tested

Species	Human
Dilution info	1/50 - 1/100

Notes	-

Predicted

Species	Mouse, Rat
Dilution info	-
Notes	-

IP

Not recommended

Species	Human, Mouse, Rat
Dilution info	-
Notes	-

WB

Tested

Species	Human
Dilution info	1/1000 - 1/10000
Notes	-

Predicted

Species	Mouse, Rat
Dilution info	-
Notes	-

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	1/999 - 1/9999

Notes	ab172730 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.

Predicted

Species	Mouse, Rat
Dilution info	-
Notes	-

Target data

See full target information GNAI1

Function	Guanine nucleotide-binding proteins (G proteins) function as transducers downstream of G protein-coupled receptors (GPCRs) in numerous signaling cascades (PubMed:18434541, PubMed:33762731, PubMed:34239069, PubMed:35610220, PubMed:37935376, PubMed:37935377, PubMed:37963465, PubMed:38552625, PubMed:8774883, PubMed:38918398, PubMed:40080544). The alpha chain contains the guanine nucleotide binding site and alternates between an active, GTP-bound state and an inactive, GDP-bound state (PubMed:18434541, PubMed:8774883). Signaling by an activated GPCR promotes GDP release and GTP binding (PubMed:18434541, PubMed:8774883). The alpha subunit has a low GTPase activity that converts bound GTP to GDP, thereby terminating the signal (PubMed:18434541, PubMed:8774883). Both GDP release and GTP hydrolysis are modulated by numerous regulatory proteins (PubMed:18434541, PubMed:8774883). Signaling is mediated via effector proteins, such as adenylate cyclase: inhibits adenylate cyclase activity of ADCY1, ADCY5 and ADCY6, leading to decreased intracellular cAMP levels (PubMed:8119955). The inactive GDP-bound form prevents the association of RGS14 with centrosomes and is required for the translocation of RGS14 from the cytoplasm to the plasma membrane. Required for normal cytokinesis during mitosis (PubMed:17635935). Required for cortical dynein-dynactin complex recruitment during metaphase (PubMed:22327364).

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C

Notes

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	The protein GNAI1 also known as G protein subunit alpha i1 is part of the G-protein family that plays an important role in signal transduction. GNAI1 has a molecular mass of approximately 40 kDa. It is mainly expressed in neuronal tissues where it interacts with receptors to regulate intracellular signaling pathways. This protein operates by inhibiting adenylate cyclase activity which reduces the production of cyclic AMP (cAMP) in response to extracellular signals.
Biological function summary	GNAI1 acts by interfering with signal transmission across membranes. It is part of the heterotrimeric G-protein complex which is essential for many signaling processes. This complex responds to activation by G-protein-coupled receptors (GPCRs) and modulates various downstream effectors essentially affecting numerous physiological processes like neurotransmission and hormone signaling.
Pathways	GNAI1 plays a significant role in the cyclic AMP (cAMP) signaling pathway and the neuroactive ligand-receptor interaction pathway. In the cAMP pathway GNAI1 associates with receptors like dopamine and serotonin receptors helping to regulate neuronal response and neurotransmitter release. This activity involves interactions with related proteins such as GNB1 and GNG2 which form the beta and gamma subunits of the heterotrimeric G-protein complex.
Associated diseases and disorders	GNAI1 relates to neuropathic conditions such as epilepsy and depression. Alterations in GNAI1 function can disrupt normal signaling contributing to these disorders. The protein interacts with other proteins in the signaling cascade such as RGS proteins which can affect its function in disease states. Researchers continue to explore how modifications or mutations in GNAI1 might offer potential therapeutic targets for these and similar disorders.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.


4 product images



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-GNAI1 antibody [EPR9441(B)] (ab140125)

GNAI1 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-GNAI1 antibody


Immunohistochemical analysis of paraffin-embedded Human thyroid carcinoma tissue labelling GNAI1 with ab140125 at 1/50 dilution.
Perform heat mediated antigen retrieval before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-GNAI1 antibody [EPR9441(B)] (ab140125)

GNAI1 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-GNAI1 antibody


Immunohistochemical analysis of paraffin-embedded Human endometrial carcinoma tissue labelling GNAI1 with ab140125 at 1/50 dilution.
Perform heat mediated antigen retrieval before commencing with IHC staining protocol.



Flow Cytometry (Intracellular) - Anti-GNAI1 antibody [EPR9441(B)] (ab140125)

GNAI1 Flow Cytometry (Intracellular) staining using rabbit Anti-GNAI1 antibody

Overlay histogram showing SHSY-5Y cells stained with ab140125 (red line). The cells were fixed with 4% paraformaldehyde (10 min) and then permeabilized with 0.1% PBS-Tween for 20 min. The cells were then incubated in 1x PBS / 10% normal goat serum / 0.3M glycine to block non-specific protein-protein interactions followed by the antibody (ab140125 1/9999 dilution) for 30 min at 22°C. The secondary antibody used was Alexa Fluor® 488 goat anti-rabbit IgG (H&L) (ab150077) at 1/2000 dilution for 30 min at 22°C. Isotype control antibody (black line) was rabbit IgG (monoclonal) (0.1µg/1x10^6 cells) used under the same conditions. Unlabelled sample (blue line) was also used as a control. Acquisition of >5000 events were collected using a 20mW Argon ion laser (488nm) and 525/30 bandpass filter. This antibody gave a positive signal in SHSY-5Y cells fixed with 80% methanol (5 min)/permeabilized with 0.1% PBS-Tween for 20 min used under the same conditions.



Western blot - Anti-GNAI1 antibody [EPR9441(B)] (ab140125)

GNAI1 Western blot staining using rabbit Anti-GNAI1 antibody

All lanes:
Western blot - Anti-GNAI1 antibody [EPR9441(B)] (ab140125) at 1/1000 dilution

Lane 1:
SH-SY5Y cell lysate at 10 µg

Lane 2:
U-87 MG cell lysate at 10 µg

Lane 3:
Human glioma tissue lysate at 10 µg

Lane 4:
Human fetal brain tissue lysate at 10 µg

Secondary

All lanes:
Goat anti-rabbit HRP conjugated at 1/2000 dilution

Predicted band size: 40 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.