abcam

ab150435

Anti-SERCA2 ATPase antibody [EPR9392]

Anti-SERCA2 ATPase antibody [EPR9392] (ab150435) is a rabbit monoclonal antibody detecting SERCA2 ATPase in **Western Blot, Flow Cytometry (Intra), IHC-P, ICC/IF**. Suitable for **Human, Mouse, Rat**.

- Biophysical QC for unrivalled batch-batch consistency
- Over 10 publications

Recombinant | RabMAb | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR9392
Purification technique	Affinity purification Protein A
Concentration	0.987 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

IHC-P

Tested

Species	Human

Dilution info	1/50 - 1/100
Notes	Perform heat-mediated antigen retrieval before commencing with IHC staining protocol.

Not recommended

Species	Mouse, Rat
Dilution info	-
Notes	-

WB

Tested

Species	Mouse
Dilution info	1/5000 - 1/10000
Notes	**For unpurified use at 1/20000 - 1/100000.**

Species	Rat
Dilution info	1/5000 - 1/10000
Notes	**For unpurified use at 1/20000 - 1/100000.**

Species	Human
Dilution info	1/5000 - 1/10000
Notes	**For unpurified use at 1/20000 - 1/100000.**

ICC/IF

Tested

Species	Human
Dilution info	1/100 - 1/250
Notes	-

Expected

Species	Mouse, Rat

Dilution info	Use at an assay dependent concentration.
Notes	-

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	1/100 - 1/1000
Notes	ab172730 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.

Expected

Species	Mouse, Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

Target data

See full target information ATP2A2

Function This magnesium-dependent enzyme catalyzes the hydrolysis of ATP coupled with the translocation of calcium from the cytosol to the sarcoplasmic reticulum lumen (PubMed:12542527, PubMed:16402920). Involved in autophagy in response to starvation. Upon interaction with VMP1 and activation, controls ER-isolation membrane contacts for autophagosome formation (PubMed:28890335). Also modulates ER contacts with lipid droplets, mitochondria and endosomes (PubMed:28890335). In coordination with FLVCR2 mediates heme-stimulated switching from mitochondrial ATP synthesis to thermogenesis (By similarity). Isoform 2. Involved in the regulation of the contraction/relaxation cycle. Acts as a regulator of TNFSF11-mediated Ca(2+) signaling pathways via its interaction with TMEM64 which is critical for the TNFSF11-induced CREB1 activation and mitochondrial ROS generation necessary for proper osteoclast generation. Association between TMEM64 and SERCA2 in the ER leads to cytosolic Ca(2+) spiking for activation of NFATC1 and production of mitochondrial ROS, thereby triggering Ca(2+) signaling cascades that promote osteoclast differentiation and activation.

Storage

Shipped at conditions	Conditional Ambient

Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

What is this antibody validated in?
Anti-SERCA2 ATPase antibody [EPR9392] (ab150435) is a rabbit recombinant monoclonal antibody and is validated for use in Western Blot (WB), Flow Cytometry (Intra), Immunohistochemistry (IHC-P), Immunocytochemistry/immunofluorescence (ICC/IF) in Human, Mouse, Rat samples.

What is the molecular weight of SERCA2 ATPase?
Anti-SERCA2 ATPase [EPR9392] (ab150435) specifically detects a band for SERCA2 ATPase (UniProt: P16615) at a molecular weight of 115kDa.

Trusted by the scientific community
Anti-SERCA2 ATPase [EPR9392] (ab150435) was first used in a scientific publication in 2012 and has been cited over 10 times in peer-reviewed journals.

Other related products
We have a range of other formats of antibody clone [EPR9392] also available for your convenience: ab150435, PE - ab237223, Carrier free - ab238426

Trial sizes available!
Test your antibody or perform pre-screening before committing to a larger quantity. Sold in 20µl. Discover our selection of trial-size antibodies.

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	SERCA2 ATPase also known as sarco/endoplasmic reticulum Cal^{2+}-ATPase 2 or ATP2A2 is an important pump responsible for the active transport of calcium ions from the cytosol into the sarcoplasmic or endoplasmic reticulum. This protein has a molecular weight of 110 kDa. It plays an important role in maintaining calcium homeostasis in muscle cells and is expressed in various tissues but particularly in cardiac and skeletal muscle. This pump uses energy from ATP hydrolysis to move calcium ions against their concentration gradient.
Biological function summary	The proper regulation of calcium ion levels by SERCA2 ATPase is essential for muscle contraction and relaxation cycles. SERCA2 exists as a part of the complex machinery in muscle cells closely interacting with phospholamban in the cardiac muscle to regulate its activity. Efficient functioning of SERCA2 directly influences calcium ion storage in the sarcoplasmic reticulum affecting muscle physiology and performance.
Pathways	The involvement of SERCA2 ATPase in the calcium signaling pathway and excitation-contraction coupling is significant. The protein's function is intimately linked with RYR2 (ryanodine receptor 2) as they both participate in the regulation of calcium ion flow in muscle cells. This interaction is necessary for the proper release and uptake of calcium ions during muscle contraction and relaxation cycles.
Associated diseases and disorders	SERCA2 ATPase mutations or dysregulation can lead to cardiac diseases such as heart failure and muscle disorders like Brody myopathy. Heart failure often results from impaired calcium handling due to disruptions in the interaction between SERCA2 and associated proteins like phospholamban. Abnormalities in SERCA2 expression or function can lead to either excessive storage or insufficient release of calcium contributing to the pathological state of these conditions.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.


9 product images



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-SERCA2 ATPase antibody [EPR9392] (ab150435)

SERCA2 ATPase Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-SERCA2 ATPase antibody


Immunohistochemical analysis of paraffin embedded Human liver tissue labelling SERCA2 ATPase with unpurified ab150435 at 1/50.
Perform heat mediated antigen retrieval before commencing with IHC staining protocol.



Immunocytochemistry/ Immunofluorescence - Anti-SERCA2 ATPase antibody [EPR9392] (ab150435)

SERCA2 ATPase Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-SERCA2 ATPase antibody


Immunocytochemistry/ Immunofluorescence analysis of HepG2 (Human hepatocellular carcinoma epithelial cell) cells labeling SERCA2 ATPase with Purified ab150435 at 1/100 dilution (10 µg/mL). Cells were fixed in 100% Methanol. Goat anti rabbit IgG (Alexa Fluor® 488, ab150077) was used as the secondary antibody at 1/1000 (2 µg/mL) dilution. DAPI (blue) was used as nuclear counterstain. PBS instead of the primary antibody was used as the secondary antibody only control.



Immunocytochemistry/ Immunofluorescence - Anti-SERCA2 ATPase antibody [EPR9392] (ab150435)

SERCA2 ATPase Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-SERCA2 ATPase antibody


Immunofluorescence analysis of HeLa cells labelling SERCA2 ATPase with unpurified ab150435 at 1/100.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-SERCA2 ATPase antibody [EPR9392] (ab150435)

SERCA2 ATPase Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-SERCA2 ATPase antibody


Immunohistochemical analysis of paraffin embedded Human kidney tissue labelling SERCA2 ATPase with unpurified ab150435 at 1/50.
Perform heat mediated antigen retrieval before commencing with IHC staining protocol.



Flow Cytometry (Intracellular) - Anti-SERCA2 ATPase antibody [EPR9392] (ab150435)

SERCA2 ATPase Flow Cytometry (Intracellular) staining using rabbit Anti-SERCA2 ATPase antibody


Overlay histogram showing HepG2 cells stained with unpurified ab150435 (red line). The cells were fixed with 80% methanol (5 min) and then permeabilized with 0.1% PBS-Tween for 20 min. The cells were then incubated in 1x PBS / 10% normal goat serum / 0.3M glycine to block non-specific protein-protein interactions followed by the antibody (ab150435, 1/1000 dilution) for 30 min at 22°C. The secondary antibody used was Alexa Fluor® 488 goat anti-rabbit IgG (H&L) (ab150077) at 1/2000 dilution for 30 min at 22°C. Isotype control antibody (black line) was rabbit IgG (monoclonal) (0.1µg/1x10^6 cells) used under the same conditions. Unlabelled sample (blue line) was also used as a control. Acquisition of >5,000 events were collected using a 20mW Argon ion laser (488nm) and 525/30 bandpass filter. This antibody gave a positive signal in HepG2 cells fixed with 4% paraformaldehyde (10 min)/permeabilized with 0.1% PBS-Tween for 20 min used under the same conditions.



Flow Cytometry (Intracellular) - Anti-SERCA2 ATPase antibody [EPR9392] (ab150435)

SERCA2 ATPase Flow Cytometry (Intracellular) staining using rabbit Anti-SERCA2 ATPase antibody


Intracellular Flow Cytometry analysis of HepG2 (Human hepatocellular carcinoma epithelial cell) cells labeling SERCA2 ATPase with Purified ab150435 at 1/1000 dilution (1 µg/mL) (Red). Cells were fixed with 80% Methanol and permeabilised with 0.1% Tween-20. A Goat anti rabbit IgG (Alexa Fluor® 488, ab150077) secondary antibody was used at 1/2000. Isotype control - Rabbit monoclonal IgG (Black). Unlabeled control - Cell without incubation with primary antibody and secondary antibody (Blue).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-SERCA2 ATPase antibody [EPR9392] (ab150435)

SERCA2 ATPase Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-SERCA2 ATPase antibody


Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of Human lung carcinoma tissue sections labeling SERCA2 ATPase with purified ab150435 at 1/50 dilution (20 µg/mL). Perform heat mediated antigen retrieval using ab93684 (Tris/EDTA buffer, pH 9.0). ImmunoHistoProbe one step HRP Polymer (ready to use) was used as the secondary antibody. Negative control: PBS instead of the primary antibody. Hematoxylin was used as a counterstain.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-SERCA2 ATPase antibody [EPR9392] (ab150435)

SERCA2 ATPase Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-SERCA2 ATPase antibody

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of Human brain tissue sections labeling SERCA2 ATPase with purified ab150435 at 1/50 dilution (20 µg/mL). Perform heat mediated antigen retrieval using ab93684 (Tris/EDTA buffer, pH 9.0). ImmunoHistoProbe one step HRP Polymer (ready to use) was used as the secondary antibody. Negative control: PBS instead of the primary antibody. Hematoxylin was used as a counterstain.



Western blot - Anti-SERCA2 ATPase antibody [EPR9392] (ab150435)

SERCA2 ATPase Western blot staining using rabbit Anti-SERCA2 ATPase antibody

Blocking/Diluting buffer: 5% NFDM/TBST
Suggest to use non-boiled samples, as boiling process could cause membrane protein aggregates (PMID: 16023741 and PMID: 8670158).

All lanes:
Western blot - Anti-SERCA2 ATPase antibody [EPR9392] (ab150435) at 1/5000 dilution

Lane 1:
HeLa (Human cervix adenocarcinoma epithelial cell) whole cell lysates boiled at 20 µg

Lane 2:
HeLa (Human cervix adenocarcinoma epithelial cell) whole cell lysates unboiled at 20 µg

Lane 3:
HepG2 (Human hepatocellular carcinoma epithelial cell) whole cell lysates boiled at 20 µg

Lane 4:
HepG2 (Human hepatocellular carcinoma epithelial cell) whole cell lysates unboiled at 20 µg

Lane 5:
Mouse brain lysates boiled at 20 µg

Lane 6:
Mouse brain lysates unboiled at 20 µg

Lane 7:
Rat brain lysates boiled at 20 µg

Lane 8:
Rat brain lysates unboiled at 20 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/20000 dilution

Predicted band size: 115 kDa

Observed band size: 115 kDa, 140 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.