abcam

ab151558

Anti-KIF5B antibody [EPR10277(B)]

Recombinant | RabMAb | KO Validated

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR10277(B)
Purification technique	Affinity purification Protein A
Concentration	0.243 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

IHC-P

Tested

Species	Human
Dilution info	1/50 - 1/100
Notes	-

WB

Tested

Species	Human
Dilution info	1/1000 - 1/10000
Notes	-

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	1/100
Notes	ab172730 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.

Target data

See full target information KIF5B

Function

Microtubule-dependent motor required for normal distribution of mitochondria and lysosomes. Can induce formation of neurite-like membrane protrusions in non-neuronal cells in a ZFYVE27-dependent manner (By similarity). Regulates centrosome and nuclear positioning during mitotic entry. During the G2 phase of the cell cycle in a BICD2-dependent manner, antagonizes dynein function and drives the separation of nuclei and centrosomes (PubMed:20386726). Required for anterograde axonal transportation of MAPK8IP3/JIP3 which is essential for MAPK8IP3/JIP3 function in axon elongation (By similarity). Through binding with PLEKHM2 and ARL8B, directs lysosome movement toward microtubule plus ends (Probable). Involved in NK cell-mediated cytotoxicity. Drives the polarization of cytolytic granules and microtubule-organizing centers (MTOCs) toward the immune synapse between effector NK lymphocytes and target cells (PubMed:24088571).

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C

Notes

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.


5 product images



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-KIF5B antibody [EPR10277(B)] (ab151558)

KIF5B Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-KIF5B antibody


Immunohistochemical analysis of paraffin-embedded Human uterus tissue labeling KIF5B with ab151558 at 1/50 dilution.
Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-KIF5B antibody [EPR10277(B)] (ab151558)

KIF5B Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-KIF5B antibody


Immunohistochemical analysis of paraffin-embedded Human colon tissue labeling KIF5B with ab151558 at 1/50 dilution.
Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.



Flow Cytometry (Intracellular) - Anti-KIF5B antibody [EPR10277(B)] (ab151558)

KIF5B Flow Cytometry (Intracellular) staining using rabbit Anti-KIF5B antibody

Overlay histogram showing HeLa cells stained with ab151558 (red line). The cells were fixed with 80% methanol (5 min) and then permeabilized with 0.1% PBS-Tween for 20 min. The cells were then incubated in 1x PBS / 10% normal goat serum / 0.3M glycine to block non-specific protein-protein interactions followed by the antibody (ab151558 1/100 dilution) for 30 min at 22°C. The secondary antibody used was Alexa Fluor® 488 goat anti-rabbit IgG (H&L) (ab150077) at 1/2000 dilution for 30 min at 22°C. Isotype control antibody (black line) was rabbit IgG (monoclonal) (1µg/1x10^6 cells) used under the same conditions. Unlabelled sample (blue line) was also used as a control. Acquisition of >5000 events were collected using a 20mW Argon ion laser (488nm) and 525/30 bandpass filter.



Western blot - Anti-KIF5B antibody [EPR10277(B)] (ab151558)

KIF5B Western blot staining using rabbit Anti-KIF5B antibody


All lanes:
Western blot - Anti-KIF5B antibody [EPR10277(B)] (ab151558) at 1/1000 dilution

Lane 1:
Jurkat cell lysate at 10 µg

Lane 2:
HeLa cell lysate at 10 µg

Lane 3:
HepG2 cell lysate at 10 µg

Predicted band size: 110 kDa



Western blot - Anti-KIF5B antibody [EPR10277(B)] (ab151558)

KIF5B Western blot staining using rabbit Anti-KIF5B antibody

Lane 1: Wild type HAP1 whole cell lysate (20 µg)
Lane 2: KIF5B knockout HAP1 whole cell lysate (20 µg)
Lane 3: Hek293 whole cell lysate (20 µg)

Lanes 1 - 3: Merged signal (red and green). Green - ab151558 observed at 110 kDa. Red - loading control, ab9484, observed at 37 kDa.

ab151558 was shown to specifically react with KIF5B in wild type cells as signal was lost in KIF5B knockout cells. Wild-type and KIF5B knockout samples were subjected to SDS-PAGE. ab151558 and ab9484 (Mouse anti GAPDH loading control) were incubated overnight at 4°C at 1000 dilution and 1/20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preabsorbed ab216773 and Goat anti-Mouse IgG H&L (IRDye® 680RD) preabsorbed ab216776 secondary antibodies at 1/20000 dilution for 1 hour at room temperature before imaging.
All lanes:
Western blot - Anti-KIF5B antibody [EPR10277(B)] (ab151558)

Predicted band size: 110 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.