abcam

ab151750

Anti-MGT4A antibody [EPR10034]

Recombinant | RabMAb

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR10034
Purification technique	Affinity purification Protein A
Concentration	0.5815 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

IHC-P

Tested

Species	Human
Dilution info	1/50 - 1/100
Notes	-

Predicted

Species	Rat, Mouse
Dilution info	-
Notes	-

IP

Expected

Species	Mouse, Rat, Human
Dilution info	Use at an assay dependent concentration.
Notes	-

Flow Cyt

Expected

Species	Mouse
Dilution info	-
Notes	ab172730 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.

Species	Rat
Dilution info	-
Notes	ab172730 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.

Species	Human
Dilution info	-
Notes	ab172730 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.

WB

Tested

Species	Human
Dilution info	1/1000 - 1/10000

Notes	-

Expected

Species	Mouse, Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

ICC/IF

Tested

Species	Human
Dilution info	1/250 - 1/500
Notes	-

Predicted

Species	Rat, Mouse
Dilution info	-
Notes	-

Target data

See full target information MGAT4A

Function	Glycosyltransferase that catalyze the transfer of GlcNAc from UDP-GlcNAc to the GlcNAcbeta1-2Manalpha1-3 arm of the core structure of N-linked glycans through a beta1-4 linkage and participates in the production of tri- and tetra-antennary N-linked sugar chains (PubMed:17006639). Involved in glucose transport by mediating SLC2A2/GLUT2 glycosylation, thereby controlling cell-surface expression of SLC2A2 in pancreatic beta cells (By similarity).

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage conditions	+4°C

Appropriate long-term storage conditions	-20°C

Notes

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	MGT4A also known as Magnesium Transporter 4A is a protein involved in magnesium ion translocation. Weighing approximately 40 kDa this protein functions as an integral membrane protein primarily facilitating the movement of magnesium ions across cellular membranes. MGT4A can be found expressed in various tissues with significant levels observed in the intestinal epithelium. It plays a critical role in maintaining magnesium homeostasis within cells.
Biological function summary	The magnesium transporter MGT4A helps regulate intracellular magnesium levels which are important for numerous cellular functions including enzymatic reactions and signal transduction. MGT4A operates as part of a larger magnesium transport complex that is responsible for ensuring the proper uptake and distribution of magnesium ions within cells. This regulation is essential for maintaining the activity of many magnesium-dependent processes within the body.
Pathways	Magnesium homeostasis is an important function of MGT4A and it participates in cellular metabolism and energy production pathways. MGT4A interacts with proteins involved in the cation transport pathway ensuring balanced magnesium levels which are necessary for ATP production and DNA synthesis. It is also related to the transporter family that includes other cation transport proteins which have roles in similar biological pathways.
Associated diseases and disorders	Mutations or malfunctions in MGT4A can lead to conditions such as hypomagnesemia and hypertension. These diseases stem from disrupted magnesium levels affecting cardiovascular and epithelial cell functions. In the context of these conditions MGT4A interacts with other transporter proteins like those involved in calcium and sodium ion transport underlining its significance in ion homeostasis and highlighting its potential as a therapeutic target for related disorders.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.


8 product images



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-MGT4A antibody [EPR10034] (ab151750)

MGT4A Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-MGT4A antibody


Immunohistochemical analysis of paraffin-embedded Human choriocarcinoma tissue labeling MGAT4A with ab151750 at 1/50 dilution.
Perform heat mediated antigen retrieval before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-MGT4A antibody [EPR10034] (ab151750)

MGT4A Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-MGT4A antibody


Immunohistochemical analysis of paraffin-embedded Human pancreas tissue labeling MGAT4A with ab151750 at 1/50 dilution.
Perform heat mediated antigen retrieval before commencing with IHC staining protocol.



Immunocytochemistry/ Immunofluorescence - Anti-MGT4A antibody [EPR10034] (ab151750)

MGT4A Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-MGT4A antibody

Immunofluorescent analysis of HeLa cells labeling MGAT4A with ab151750 at 1/250 dilution.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-MGT4A antibody [EPR10034] (ab151750)

MGT4A Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-MGT4A antibody

Immunohistochemical analysis of paraffin embedded normal Human kidney tissue using ab151750 showing +ve staining.

Perform heat mediated antigen retrieval before commencing with IHC staining protocol.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-MGT4A antibody [EPR10034] (ab151750)

MGT4A Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-MGT4A antibody

Immunohistochemical analysis of paraffin embedded Human Lung carcinoma tissue using ab151750 showing +ve staining.

Perform heat mediated antigen retrieval before commencing with IHC staining protocol.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-MGT4A antibody [EPR10034] (ab151750)

MGT4A Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-MGT4A antibody

Immunohistochemical analysis of paraffin embedded normal Human colon tissue using ab151750 showing +ve staining.

Perform heat mediated antigen retrieval before commencing with IHC staining protocol.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-MGT4A antibody [EPR10034] (ab151750)

MGT4A Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-MGT4A antibody

Immunohistochemical analysis of paraffin embedded Human Prostate hyperplasia tissue using ab151750 showing +ve staining.

Perform heat mediated antigen retrieval before commencing with IHC staining protocol.



Western blot - Anti-MGT4A antibody [EPR10034] (ab151750)

All lanes:
Western blot - Anti-MGT4A antibody [EPR10034] (ab151750) at 1/1000 dilution

Lane 1:
Human fetal heart lysate at 10 µg

Lane 2:
Human fetal kidney lysate at 10 µg

Lane 3:
HeLa lysate at 10 µg

Secondary

All lanes:
Goat anti-rabbit HRP at 1/2000 dilution

Predicted band size: 109 kDa, 62 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.