abcam

ab154837

Anti-GBGT1 antibody [EPR10467]

Recombinant | RabMAb

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR10467
Purification technique	Affinity purification Protein A
Concentration	0.276 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

IHC-P

Tested

Species	Human
Dilution info	1/100 - 1/250
Notes	-

Predicted

Species	Mouse, Rat
Dilution info	-
Notes	-

IP

Not recommended

Species	Human, Mouse, Rat
Dilution info	-
Notes	-

WB

Tested

Species	Human
Dilution info	1/1000 - 1/10000
Notes	-

Predicted

Species	Mouse, Rat
Dilution info	-
Notes	-

ICC/IF

Not recommended

Species	Human, Mouse, Rat
Dilution info	-
Notes	-

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	1/10 - 1/100
Notes	ab172730 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.

Predicted

Species	Mouse, Rat
Dilution info	-
Notes	-

Target data

See full target information GBGT1

Function	Has lost the ability to synthesize Forssman glycolipid antigen (FORS1/FG) (PubMed:10506200). Might have acquired an alternative function in glycosphingolipid metabolism, but it remains to be established. It appears to have drifted more slowly than confirmed pseudogenes in the glycosyltransferase 6 family, suggesting that it has remained under evolutionary pressure.

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C

Notes

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity

- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	The GBGT1 protein also known as Forssman glycosyltransferase functions as an enzyme involved in the biosynthesis of the Forssman antigen. It is a glycosyltransferase enzyme that adds a specific sugar residue to the terminal position on glyco-conjugates. This enzyme typically weighs around 37 kDa. GBGT1 is expressed in a variety of tissues including the kidney heart and small intestine but its expression varies significantly among different species.
Biological function summary	The Forssman glycosyltransferase enzyme is responsible for the generation of Forssman glycolipids contributing to the structure of cell membranes. These glycolipids play a role in cell-cell interactions. GBGT1 is not typically part of a larger protein complex but acts independently to modify glycolipids affecting cellular communication and recognition processes.
Pathways	GBGT1 is involved in glycosphingolipid biosynthesis pathways specifically influencing the metabolic progression from globoseries to Forssman series glycolipids. This pathway closely interacts with other enzymes like α13-galactosyltransferase which also contributes to glycolipid modification. These enzymes together maintain proper cellular roles involving signaling and interactions at the cell surface.
Associated diseases and disorders	GBGT1 engagement has been associated with immunological responses and cancer. Aberrant expression or mutation in this enzyme affects antigen presentation which contributes to various cancer types as the presence of Forssman antigen is often noted in tumor tissues. Furthermore it has connections with the α-galactosyltransferase enzyme in xenotransplantation as both are involved in the synthesis of glycolipids that can trigger immune rejection events.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended

We do not recommend this combination. It is not covered by our product promise.


We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

---


4 product images



Flow Cytometry (Intracellular) - Anti-GBGT1 antibody [EPR10467] (ab154837)

GBGT1 Flow Cytometry (Intracellular) staining using rabbit Anti-GBGT1 antibody


Intracellular Flow Cytometry analysis of permeabilized Jurkat cells labeling GBGT1 (red), using ab154837 at a 1/10 dilution, and negative control cells probed with a Rabbit IgG (green)



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-GBGT1 antibody [EPR10467] (ab154837)

GBGT1 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-GBGT1 antibody


Immunohistochemical analysis of formalin-fixed paraffin-embedded Human pancreas tissue labeling GBGT1 using ab154837 at a 1/100 dilution.
Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-GBGT1 antibody [EPR10467] (ab154837)

GBGT1 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-GBGT1 antibody

Immunohistochemical analysis of formalin-fixed paraffin-embedded Human kidney tissue labeling GBGT1 using ab154837 at a 1/100 dilution.
Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.



Western blot - Anti-GBGT1 antibody [EPR10467] (ab154837)

GBGT1 Western blot staining using rabbit Anti-GBGT1 antibody

All lanes:
Western blot - Anti-GBGT1 antibody [EPR10467] (ab154837) at 1/1000 dilution

Lane 1:
Human placenta tissue lysate at 10 µg

Lane 2:
Human ovary cancer tissue lysate at 10 µg

Lane 3:
Human fetal heart tissue lysate at 10 µg

Lane 4:
Human fetal lung tissue lysate at 10 µg

Lane 5:
HeLa cell lysate at 10 µg

Lane 6:
Jurkat cell lysate at 10 µg

Predicted band size: 40 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.