abcam

ab154867

Anti-DTYMK antibody [EPR10163]

Recombinant | RabMAb

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR10163
Purification technique	Affinity purification Protein A
Concentration	0.132 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

IHC-P

Tested

Species	Human
Dilution info	1/100 - 1/250
Notes	-

WB

Tested

Species	Human
Dilution info	1/1000 - 1/10000
Notes	-

ICC/IF

Tested

Species	Human
Dilution info	1/100 - 1/250
Notes	-

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	1/1000
Notes	-

Target data

See full target information DTYMK

Function	Catalyzes the phosphorylation of thymidine monophosphate (dTMP) to thymidine diphosphate (dTDP), the immediate precursor for the DNA building block dTTP, with ATP as the preferred phosphoryl donor in the presence of Mg(2+).

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C

Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.


7 product images



Immunocytochemistry/ Immunofluorescence - Anti-DTYMK antibody [EPR10163] (ab154867)

DTYMK Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-DTYMK antibody

Immunofluorescent analysis of HeLa cells labeling DTYMK with ab154867 at 1/100.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-DTYMK antibody [EPR10163] (ab154867)

DTYMK Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-DTYMK antibody

Immunohistochemical analysis of paraffin embedded Human normal kidney tissue using ab154867 showing +ve staining.


Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.



Flow Cytometry (Intracellular) - Anti-DTYMK antibody [EPR10163] (ab154867)

DTYMK Flow Cytometry (Intracellular) staining using rabbit Anti-DTYMK antibody


Flow Cytometry analysis of HeLa (Human cervix adenocarcinoma epithelial cell) cells labeling DTYMK with purified ab154867 at 1/1000 dilution (1 µg/mL) (Red). Cells were fixed with 4% Paraformaldehyde and permeabilised with 90% Methanol. A Goat anti rabbit IgG (Alexa Fluor® 488, ab150077) secondary antibody was used at 1/2000. Isotype control - Rabbit monoclonal IgG (Black). Unlabeled control - Cell without incubation with primary antibody and secondary antibody (Blue).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-DTYMK antibody [EPR10163] (ab154867)

Immunohistochemical analysis of paraffin embedded Human prostate hyperplasia tissue using ab154867 showing +ve staining.

Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-DTYMK antibody [EPR10163] (ab154867)

DTYMK Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-DTYMK antibody

Immunohistochemical analysis of paraffin embedded Human skeletal muscle tissue using ab154867 showing +ve staining.


Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-DTYMK antibody [EPR10163] (ab154867)

DTYMK Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-DTYMK antibody

Immunohistochemical analysis of paraffin-embedded Human thyroid gland carcinoma tissue labeling DTYMK with ab137875 at 1/100.
Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.



Western blot - Anti-DTYMK antibody [EPR10163] (ab154867)

DTYMK Western blot staining using rabbit Anti-DTYMK antibody

All lanes:
Western blot - Anti-DTYMK antibody [EPR10163] (ab154867) at 1/1000 dilution

Lane 1:
Raji cell lysate at 10 µg

Lane 2:
K562 cell lysate at 10 µg

Lane 3:
HeLa cell lysate at 10 µg

Predicted band size: 23 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.