abcam

ab155284

Anti-TPMT antibody [EPR10820(B)]

Recombinant | RabMAb | KO Validated

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR10820(B)
Purification technique	Affinity purification Protein A
Concentration	0.621 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

IHC-P

Tested

Species	Human
Dilution info	1/100 - 1/250
Notes	-

Predicted

Species	Mouse, Rat
Dilution info	-
Notes	-

WB

Tested

Species	Human
Dilution info	1/1000 - 1/10000
Notes	-

Predicted

Species	Mouse, Rat
Dilution info	-
Notes	-

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	1/100 - 1/500
Notes	ab172730 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.

Predicted

Species	Mouse, Rat
Dilution info	-
Notes	-

Target data

See full target information TPMT

Function	Catalyzes the S-methylation of thiopurine drugs such as 6-mercaptopurine (also called mercaptopurine, 6-MP or its brand name Purinethol) and 6-thioguanine (also called tioguanine or 6-TG) using S-adenosyl-L-methionine as the methyl donor (PubMed:18484748, PubMed:657528). TPMT activity modulates the cytotoxic effects of thiopurine prodrugs. A natural substrate for this enzyme has yet to be identified.

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C

Notes

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	Thiopurine S-methyltransferase often called TPMT plays a role as an enzyme that catalyzes the S-methylation of thiopurine compounds like 6-mercaptopurine and azathioprine. It exists as a monomer with a molecular mass of about 28 kDa. In humans TPMT expresses mainly in the liver and kidney. The activity of TPMT varies among individuals due to genetic polymorphisms.
Biological function summary	TPMT is important in the metabolism of therapeutic thiopurine drugs. These drugs often require precise dosing to achieve effectiveness without toxicity. TPMT does not form part of a larger complex; it functions independently to methylate thiopurine drugs. Genetic variations can lead to different TPMT activity levels affecting a person's response to these drugs significantly.

Pathways	TPMT is integral to the drug metabolism pathway specifically in the detoxification and elimination processes of thiopurine drugs. While TPMT predominantly acts on thiopurines thioredoxin reductase 1 (TXNRD1) may also have a role in modulating redox balance and affect the thiopurine metabolism indirectly through related cellular processes. Together they influence the cellular response to drugs like azathioprine.
Associated diseases and disorders	Genetic variants in TPMT significantly contribute to thiopurine drug intolerance or toxicity particularly in conditions like inflammatory bowel disease and acute lymphoblastic leukemia where these drugs are used. Patients with low TPMT activity are at risk for severe myelosuppression a life-threatening bone marrow suppression. TPMT's relationship with the enzyme inosine monophosphate dehydrogenase (IMPDH) also illustrates its connection to nucleotide synthesis further linking TPMT to therapeutic outcomes in cancers.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.


8 product images



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-TPMT antibody [EPR10820(B)] (ab155284)

TPMT Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-TPMT antibody

Immunohistochemical analysis of paraffin-embedded Human kidney tissue labeling TPMT with ab155284 at 1/100 dilution.
Perform heat mediated antigen retrieval before commencing with IHC staining protocol.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-TPMT antibody [EPR10820(B)] (ab155284)

TPMT Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-TPMT antibody

Immunohistochemical analysis of paraffin embedded Human gastric adenocarcinoma tissue using ab155284 showing +ve staining.

Perform heat mediated antigen retrieval before commencing with IHC staining protocol.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-TPMT antibody [EPR10820(B)] (ab155284)

TPMT Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-TPMT antibody

Immunohistochemical analysis of paraffin embedded Human thyroid gland carcinoma tissue using ab155284 showing +ve staining.


Perform heat mediated antigen retrieval before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-TPMT antibody [EPR10820(B)] (ab155284)

TPMT Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-TPMT antibody


Immunohistochemical analysis of paraffin-embedded Human brain tissue labeling TPMT with ab155284 at 1/100 dilution.
Perform heat mediated antigen retrieval before commencing with IHC staining protocol.



Flow Cytometry (Intracellular) - Anti-TPMT antibody [EPR10820(B)] (ab155284)

TPMT Flow Cytometry (Intracellular) staining using rabbit Anti-TPMT antibody

Intracellular flow cytometric analysis of permeabilized K562 cells labeling TPMT with ab155284 at 1/100 dilution (red) compared with a rabbit IgG (negative) (green).




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-TPMT antibody [EPR10820(B)] (ab155284)

TPMT Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-TPMT antibody

Immunohistochemical analysis of paraffin embedded Human normal uterus tissue using ab155284 showing +ve staining.

Perform heat mediated antigen retrieval before commencing with IHC staining protocol.



Western blot - Anti-TPMT antibody [EPR10820(B)] (ab155284)

All lanes:
Western blot - Anti-TPMT antibody [EPR10820(B)] (ab155284) at 1/1000 dilution

Lane 1:
K562 cell lysate at 10 µg

Lane 2:
TF1 cell lysate at 10 µg


Lane 3:
Jurkat cell lysate at 10 µg

Lane 4:
HepG2 cell lysate at 10 µg

Secondary

All lanes:
Goat anti-rabbit HRP at 1/2000 dilution

Predicted band size: 28 kDa



Western blot - Anti-TPMT antibody [EPR10820(B)] (ab155284)

TPMT Western blot staining using rabbit Anti-TPMT antibody

Lane 1: Wild-type HAP1 whole cell lysate (20 µg)
Lane 2: TPMT knockout HAP1 whole cell lysate (20 µg)
Lane 3: HepG2 whole cell lysate (20 µg)
Lane 4: Jurkat whole cell lysate (20 µg)
Lanes 1 - 4: Merged signal (red and green). Green - ab155284 observed at 28 kDa. Red - loading control, ab9484, observed at 37 kDa.

ab155284 was shown to recognize TPMT in wild-type HAP1 cells as signal was lost at the expected MW in TPMT knockout cells. Additional cross-reactive bands were observed in the wild-type and knockout cells. Wild-type and TPMT knockout samples were subjected to SDS-PAGE. ab155284 and ab9484 (Mouse anti-GAPDH loading control) were incubated overnight at 4°C at 2 µg/ml and 1/20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preabsorbed ab216773 and Goat anti-Mouse IgG H&L (IRDye® 680RD) preabsorbed ab216776 secondary antibodies at 1/10000 dilution for 1 hour at room temperature before imaging.

All lanes:
Western blot - Anti-TPMT antibody [EPR10820(B)] (ab155284)

Predicted band size: 28 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.