abcam

ab155992

Anti-TIP-1 antibody [EPR7900(2)]

Recombinant | RabMAb

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR7900(2)
Purification technique	Affinity purification Protein A
Concentration	2.048 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

IHC-P

Tested

Species	Human
Dilution info	1/1000
Notes	**For unpurified use at 1/50 - 1/100.** Perform heat-mediated antigen retrieval before commencing with IHC staining protocol.

Not recommended

Species	Mouse, Rat
Dilution info	-
Notes	-

ICC/IF

Not recommended

Species	Mouse, Rat, Human
Dilution info	-
Notes	-

WB

Tested

Species	Mouse
Dilution info	1/1000
Notes	**For unpurified use at 1/1000 - 1/10000.**

Species	Rat
Dilution info	1/1000
Notes	**For unpurified use at 1/1000 - 1/10000.**

Species	Human
Dilution info	1/1000
Notes	**For unpurified use at 1/1000 - 1/10000.**

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	1/200

Notes	For unpurified use at 1/100 - 1/250. ab172730 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.

Expected

Species	Mouse, Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

Target data

See full target information TAX1BP3

Function	May regulate a number of protein-protein interactions by competing for PDZ domain binding sites. Binds CTNNB1 and may thereby act as an inhibitor of the Wnt signaling pathway. Competes with LIN7A for KCNJ4 binding, and thereby promotes KCNJ4 internalization. May play a role in the Rho signaling pathway. May play a role in activation of CDC42 by the viral protein HPV16 E6.

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	TIP-1 also known as Tax-interacting protein-1 or glutaminase-interacting protein (GIP) is a small protein with a mass of approximately 14.5 kDa. It functions mechanically by binding to PDZ domains found in a wide range of cellular proteins. TIP-1 expression occurs in various tissues with notable presence in the brain liver and kidney. Its ability to interact with PDZ domain-containing proteins plays an important role in cellular maintenance and signaling.
Biological function summary	TIP-1 functions as a regulator of cellular signaling pathways through its interaction with PDZ-domain proteins. It does not typically form part of a larger protein complex but rather acts as an independent modulator. TIP-1 influences cellular signaling by affecting the spatial arrangement of proteins in cells thereby impacting signal transduction mechanisms. This regulation can affect various biological processes including cell proliferation and apoptosis.
Pathways	TIP-1 plays a role in the Wnt signaling pathway where it interacts with proteins such as Dishevelled which is important for transducing Wnt signals. TIP-1 also has implications in the glutamate metabolic pathway by interacting with proteins like glutaminase. These pathways are essential for cellular growth processes and energy metabolism regulation showing the interconnectivity of TIP-1 with key cellular functions.
Associated diseases and disorders	TIP-1 implicates itself in cancer and neurological disorders. Overexpression of TIP-1 has been associated with certain types of cancer where it may interact with proteins like β-catenin to impact cell proliferation and migration. In neurological disorders dysfunction in TIP-1's interactions within signaling pathways may contribute to diseases such as schizophrenia closely linked with disturbed glutamate signaling and TIP-1's PDZ domain interactions.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended

We do not recommend this combination. It is not covered by our product promise.


We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

---


3 product images



Flow Cytometry (Intracellular) - Anti-TIP-1 antibody [EPR7900(2)] (ab155992)

TIP-1 Flow Cytometry (Intracellular) staining using rabbit Anti-TIP-1 antibody


Intracellular Flow Cytometry analysis of A-673 (Human muscle Ewing's Sarcoma) cells labeling TIP-1 with purified ab155992 at 1/200 dilution (10µg/mL) (Red). Cells were fixed with 4% Paraformaldehyde and permeabilised with 90% Methanol. A Goat anti rabbit IgG (Alexa Fluor® 488, ab150077) secondary antibody was used at 1/2000. Isotype control - Rabbit monoclonal IgG (Black). Unlabeled control - Cell without incubation with primary antibody and secondary antibody (Blue).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-TIP-1 antibody [EPR7900(2)] (ab155992)

TIP-1 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-TIP-1 antibody

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human thyroid carcinoma tissue sections labeling TIP-1 with purified ab155992 at 1/1000 dilution (2.05 µg/mL). Perform heat mediated antigen retrieval using ab93684 (Tris/EDTA buffer, pH 9.0). ImmunoHistoProbe one step HRP Polymer (ready to use) was used as the secondary antibody. Negative control: PBS instead of the primary antibody. Hematoxylin was used as a counterstain.



Western blot - Anti-TIP-1 antibody [EPR7900(2)] (ab155992)

TIP-1 Western blot staining using rabbit Anti-TIP-1 antibody

Blocking/Diluting buffer: 5% NFDM/TBST
All lanes:
Western blot - Anti-TIP-1 antibody [EPR7900(2)] (ab155992) at 1/1000 dilution

Lane 1:
A-673 (Human muscle Ewing's Sarcoma) whole cell lysate at 20 µg

Lane 2:
NIH/3T3 (Mouse embryonic fibroblast) whole cell lysate at 20 µg

Lane 3:
LLC1 (Mus musculus lung rounded-loosely attached or floating) whole cell lysate at 20 µg

Lane 4:
Mouse lung lysate at 20 µg

Lane 5:
Mouse heart lysate at 20 µg

Lane 6:
C6 (Rat glial tumor glial cell) whole cell lysate at 20 µg

Lane 7:
Rat lung lysate at 20 µg

Lane 8:
Rat heart tissue lysate at 20 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/20000 dilution

Predicted band size: 14 kDa

Observed band size: 14 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.