abcam

ab157333

PE Anti-p75 NGF Receptor antibody [NGFR5]

Anti-p75 NGF Receptor antibody [NGFR5] conjugated to PE - suitable for flow cytometry, and reacts with human, rabbit samples. Clone [NGFR5] is a mouse monoclonal antibody cited in 12 publications.

Design your panel with our interactive flow cytometry panel builder. Click here.

Key facts

Isotype	IgG1
Host species	Mouse
Conjugation	PE
Excitation/Emission	Ex: 480;565nm, Em: 578nm
Storage buffer	pH: 7.4 Preservative: 0.1% Sodium azide Constituents: PBS, 0.2% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	Native Full Length Protein corresponding to Human NGFR. Database link P08138
Clone number	NGFR5
Purification technique	Size-exclusion chromatography
Epitope	The epitope is localized within amino acids 1-160.

Reactivity data

Flow Cyt

Expected

Species	Primates

Dilution info	10 µL for 10^6 Cells
Notes	10ul per 100ul whole blood. ab91357 - Mouse monoclonal IgG1, is suitable for use as an isotype control with this antibody.

Species	Ferret
Dilution info	10 µL for 10^6 Cells
Notes	10ul per 100ul whole blood. ab91357 - Mouse monoclonal IgG1, is suitable for use as an isotype control with this antibody.

Species	Cat
Dilution info	10 µL for 10^6 Cells
Notes	10ul per 100ul whole blood. ab91357 - Mouse monoclonal IgG1, is suitable for use as an isotype control with this antibody.

Species	Rabbit
Dilution info	10 µL for 10^6 Cells
Notes	10ul per 100ul whole blood. ab91357 - Mouse monoclonal IgG1, is suitable for use as an isotype control with this antibody.

Not recommended

Species	Mouse
Dilution info	-
Notes	10ul per 100ul whole blood. ab91357 - Mouse monoclonal IgG1, is suitable for use as an isotype control with this antibody.

Species	Rat
Dilution info	-
Notes	10ul per 100ul whole blood. ab91357 - Mouse monoclonal IgG1, is suitable for use as an isotype control with this antibody.

Species	Human
Dilution info	-
Notes	-

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	10 µL for 10^6 Cells
Notes	Use 10 µl reagent / 100 µl of whole blood or 10^6 cells in a suspension. ab91357 - Mouse monoclonal IgG1, is suitable for use as an isotype control with this antibody.

Predicted

Species	Rabbit, Cat, Ferret, Primates
Dilution info	-
Notes	-

Not recommended

Species	Mouse, Rat
Dilution info	-
Notes	-

Target data

See full target information NGFR

Function

Low affinity receptor which can bind to NGF, BDNF, NTF3, and NTF4. Forms a heterodimeric receptor with SORCS2 that binds the precursor forms of NGF, BDNF and NTF3 with high affinity, and has much lower affinity for mature NGF and BDNF (PubMed:24908487). Plays an important role in differentiation and survival of specific neuronal populations during development (By similarity). Can mediate cell survival as well as cell death of neural cells. Plays a role in the inactivation of RHOA (PubMed:26646181). Plays a role in the regulation of the translocation of GLUT4 to the cell surface in adipocytes and skeletal muscle cells in response to insulin, probably by regulating RAB31 activity, and thereby contributes to the regulation of insulin-dependent glucose uptake (By similarity). Necessary for the circadian oscillation of the clock genes BMAL1, PER1, PER2 and NR1D1 in the suprachiasmatic nucleus (SCmgetaN) of the brain and in liver and of the genes involved in glucose and lipid metabolism in the liver (PubMed:23785138). Together with BFAR negatively regulates NF-kappa-B and JNK-related signaling pathways (PubMed:22566094).

Storage

Shipped at conditions	Blue Ice

Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	+4°C

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	The p75 NGF receptor also known as p75 NGFR or anti-NTR is a transmembrane protein with a molecular mass of approximately 75 kDa. It is a member of the tumor necrosis factor receptor superfamily. This receptor is widely expressed in the nervous system particularly in neurons glial cells and Schwann cells. The p75 receptor primarily binds to neurotrophins such as nerve growth factor (NGF) influencing neuronal survival apoptosis and growth.
Biological function summary	The p75 receptor functions as an important part of a signaling complex that mediates diverse cellular responses. It often forms complexes with other receptors like TrkA but can also signal independently. In particular the p75 protein can form a biological chimera with different co-receptors influencing its downstream effects. This makes it significant in processes like apoptosis and neuronal differentiation as well as influencing neurotrophin-mediated signaling.
Pathways	The p75 receptor plays an important role in pathways involving neuronal survival and apoptosis. One key pathway involves the nerve growth factor (NGF) signaling pathway where p75 collaborates with Trk receptors enhancing or inhibiting NGF signaling based on cellular context. Another pathway is the JNK signaling pathway where the p75 receptor interacts with proteins like sortilin contributing to apoptosis and stress responses in neurons. These interactions determine the outcome of neurotrophic signaling highlighting its role in cell fate decisions.
Associated diseases and disorders	P75 receptor is associated with several neurodegenerative conditions and cancers. Its involvement in Alzheimer's disease is significant because altered p75 expression and function can contribute to neurodegeneration through apoptosis pathways. This receptor is also linked to glioblastoma where it might promote tumor progression in coordination with proteins such as sortilin. These links make p75 receptor a target of interest for therapeutic intervention in these diseases.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended

We do not recommend this combination. It is not covered by our product promise.


We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

---

1 product image



Flow Cytometry (Intracellular) - PE Anti-p75 NGF Receptor antibody [NGFR5] (ab157333)

P75 NGF Receptor Flow Cytometry (Intracellular) staining of REH cells using mouse Anti-p75 NGF Receptor antibody

Separation of REH cells stained using ab157333 (concentration in sample 0,33 µg/ml, red) from unstained REH cells (grey) in flow cytometry analysis (intracellular staining).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.