abcam

ab166628

Anti-Proteasome subunit beta type 2/PSMB2 antibody [EPR8826(2)(B)]

Recombinant | RabMAb

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR8826(2)(B)
Purification technique	Affinity purification Protein A
Concentration	0.631 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

IHC-P

Tested

Species	Human
Dilution info	1/100 - 1/250
Notes	-

Predicted

Species	Mouse, Rat
Dilution info	-
Notes	-

WB

Tested

Species	Human
Dilution info	1/1000 - 1/10000
Notes	-

Predicted

Species	Mouse, Rat
Dilution info	-
Notes	-

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	1/100 - 1/500
Notes	ab172730 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.

Predicted

Species	Mouse, Rat
Dilution info	-
Notes	-

Target data

See full target information PSMB2

Function	Non-catalytic component of the 20S core proteasome complex involved in the proteolytic degradation of most intracellular proteins. This complex plays numerous essential roles within the cell by associating with different regulatory particles. Associated with two 19S regulatory particles, forms the 26S proteasome and thus participates in the ATP-dependent degradation of ubiquitinated proteins. The 26S proteasome plays a key role in the maintenance of protein homeostasis by removing misfolded or damaged proteins that could impair cellular functions, and by removing proteins whose functions are no longer required. Associated with the PA200 or PA28, the 20S proteasome mediates ubiquitin-independent protein degradation. This type of proteolysis is required in several pathways including spermatogenesis (20S-PA200 complex) or generation of a subset of MHC class I-presented antigenic peptides (20S-PA28 complex).

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	Proteasome subunit beta type 2 also known as PSMB2 is a part of the proteasome complex which plays an important role in protein degradation. PSMB2 has a mass of about 22 kDa and functions within the multi-catalytic proteinase complex known as the 20S proteasome core particle. This protein is expressed in various tissues with significant activities found in cells that demand high protein turnover. PSMB2 contributes to the proteasome's chymotrypsin-like activity aiding in the breakdown of damaged or unneeded proteins into peptides.
Biological function summary	Proteasome subunit beta type 2 is essential in maintaining protein homeostasis. It functions as a component of the 20S core of the 26S proteasome where it participates in degrading ubiquitin-tagged proteins. This complex also regulates multiple cellular processes including cell cycle progression and signal transduction. Proper function of PSMB2 is necessary for removing misfolded or damaged proteins thereby preventing their accumulation which could be toxic to cells.
Pathways	Proteasome subunit beta type 2 has a role in the ubiquitin-proteasome pathway which is important for protein degradation and turnover. This pathway contributes to important cellular processes such as DNA repair and antigen processing. PSMB2 in this context is linked to other proteins like ubiquitin-ligase that mediate the tagging of proteins for degradation. Another pathway involving PSMB2 is the NF-kB signaling pathway which controls inflammatory responses and immune system regulation.
Associated diseases and disorders	Proteasome subunit beta type 2 has been implicated in neurodegenerative diseases like Parkinson’s. The dysregulation or malfunctioning of the proteasome complex might result in protein aggregation seen in such disorders. Additionally PSMB2 is associated with certain cancers due to its role in regulating apoptosis and cellular proliferative activities. Proteins like p53 which is important for tumor suppression are related to PSMB2 through these disease mechanisms highlighting its significance in maintaining normal cellular function and preventing disease progression.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.


5 product images



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Proteasome subunit beta type 2/PSMB2 antibody [EPR8826(2)(B)] (ab166628)

Proteasome subunit beta type 2/PSMB2 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-Proteasome subunit beta type 2/PSMB2 antibody


Immunohistochemical analysis of paraffin embedded Human gastric carcinoma tissue labeling Proteasome subunit beta type 2/PSMB2 with ab166628 at a 1/100 dilution.
Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.



Flow Cytometry (Intracellular) - Anti-Proteasome subunit beta type 2/PSMB2 antibody [EPR8826(2)(B)] (ab166628)

Proteasome subunit beta type 2/PSMB2 Flow Cytometry (Intracellular) staining using rabbit Anti-Proteasome subunit beta type 2/PSMB2 antibody


Intracellular flow cytometric analysis of permeabilized Jurkat cells using ab166628 at a 1/100 dilution (red) or a Rabbit IgG (negative) (green).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Proteasome subunit beta type 2/PSMB2 antibody [EPR8826(2)(B)] (ab166628)

Proteasome subunit beta type 2/PSMB2 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-Proteasome subunit beta type 2/PSMB2 antibody

Immunohistochemical analysis of paraffin embedded Human liver tissue labeling Proteasome subunit beta type 2/PSMB2 with ab166628 at a 1/100 dilution.
Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.



Western blot - Anti-Proteasome subunit beta type 2/PSMB2 antibody [EPR8826(2)(B)] (ab166628)


All lanes:
Western blot - Anti-Proteasome subunit beta type 2/PSMB2 antibody [EPR8826(2)(B)] (ab166628) at 1/1000 dilution

All lanes:
immunoprecipitation pellet from HeLa (Human cervix adenocarcinoma epithelial cell) whole cell lysate at 10 µg

Secondary

All lanes:
HRP-conjugated anti-Rabbit IgG preferentially detecting the non-reduced form of Rabbit IgG

Predicted band size: 23 kDa



Western blot - Anti-Proteasome subunit beta type 2/PSMB2 antibody [EPR8826(2)(B)] (ab166628)

All lanes:
Western blot - Anti-Proteasome subunit beta type 2/PSMB2 antibody [EPR8826(2)(B)] (ab166628) at 1/1000 dilution

Lane 1:
Jurkat cell lysate at 10 µg

Lane 2:
HepG2 cell lysate at 10 µg

Lane 3:
HeLa cell lysate at 10 µg

Lane 4:
LoVo cell lysate at 10 µg

Predicted band size: 23 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.