abcam

ab170952

Anti-ASS1 antibody [EPR12398]

Anti-ASS1 antibody [EPR12398] (ab170952) is a rabbit monoclonal antibody detecting ASS1 in **Western Blot, Flow Cytometry (Intra), IP, IHC-P, ICC/IF**. Suitable for **Human, Mouse, Rat**.

- KO validated for confirmed specificity
- Biophysical QC for unrivalled batch-batch consistency

Recombinant | RabMAb | KO Validated | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR12398
Purification technique	Affinity purification Protein A
Concentration	0.962 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

IHC-P

Tested

Species	Mouse

Dilution info	1/4000
Notes	**For unpurified use at 1/250 - 1/500.** Perform heat-mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.

Species	Human
Dilution info	1/4000
Notes	**For unpurified use at 1/250 - 1/500.** Perform heat-mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.

Not recommended

Species	Rat, Cow
Dilution info	-
Notes	-

IP

Tested

Species	Human
Dilution info	1/10 - 1/100
Notes	-

Expected

Species	Mouse, Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

Predicted

Species	Cow
Dilution info	-
Notes	-

WB

Tested

Species	Mouse
Dilution info	1/20000
Notes	**For unpurified use at 1/1000 - 1/10000.**

Species	Rat
Dilution info	1/20000
Notes	**For unpurified use at 1/1000 - 1/10000.**

Species	Human
Dilution info	1/20000
Notes	**For unpurified use at 1/1000 - 1/10000.**

Predicted

Species	Cow
Dilution info	-
Notes	-

ICC/IF

Tested

Species	Human
Dilution info	1/50 - 1/100
Notes	-

Expected

Species	Mouse, Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

Predicted

Species	Cow
Dilution info	-
Notes	-

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	1/100
Notes	For unpurified use at 1/500 - 1/1000. ab172730 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.

Expected

Species	Mouse, Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

Predicted

Species	Cow
Dilution info	-
Notes	-

Target data

See full target information ASS1

Function	One of the enzymes of the urea cycle, the metabolic pathway transforming neurotoxic amonia produced by protein catabolism into inocuous urea in the liver of ureotelic animals. Catalyzes the formation of arginosuccinate from aspartate, citrulline and ATP and together with ASL it is responsible for the biosynthesis of arginine in most body tissues.

Storage

Shipped at conditions	Conditional Ambient

Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

What is this antibody validated in?
Anti-ASS1 antibody [EPR12398] (ab170952) is a rabbit recombinant monoclonal antibody and is validated for use in Western Blot (WB), Flow Cytometry (Intra), Immunoprecipitation (IP), Immunohistochemistry (IHC-P), Immunocytochemistry/immunofluorescence (ICC/IF) in Human, Mouse, Rat samples.

What is the molecular weight of ASS1?
Anti-ASS1 [EPR12398] (ab170952) specifically detects a band for ASS1 (UniProt: P00966) at a molecular weight of 47kDa.

Trusted by the scientific community
Anti-ASS1 [EPR12398] (ab170952) was first used in a scientific publication in 2013 and has been cited over 10 times in peer-reviewed journals.

Specificity confirmed
The specificity of Anti-ASS1 antibody [EPR12398] (ab170952) has been confirmed by Western blot testing in ASS1 Knockout HAP1 cell line, ab264989.

Other related products
We have a range of other formats of antibody clone [EPR12398] also available for your convenience: ab170952, Alexa Fluor® 488 - ab208412, HRP - ab209018, PE - ab210451, Carrier free - ab231684, APC - ab310832, Alexa Fluor® 647 - ab311083, Alexa Fluor® 594 - ab311677, Alexa Fluor® 555 - ab313161

Trial sizes available!
Test your antibody or perform pre-screening before committing to a larger quantity. Sold in 20µl. Discover our selection of trial-size antibodies.

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility

- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	ASS1 or argininosuccinate synthase 1 is an enzyme that plays an important role in the urea cycle. It catalyzes the conversion of citrulline and aspartate into argininosuccinate using ATP in the process. The ASS1 protein is commonly expressed in the liver where it actively participates in the detoxification of ammonia. It has a molecular mass of approximately 46 kDa. Some alternate names for this protein include ASS and ASS-1.
Biological function summary	Argininosuccinate synthase 1 contributes significantly to the synthesis of arginine an essential amino acid through its conversion activities. The enzyme operates as a homotrimer a complex consisting of three identical subunits. This enzymatic activity is indispensable in maintaining the balance of nitrogen and ammonia within the body particularly important for liver and kidney functions.
Pathways	ASS1 is a critical component of both the urea cycle and the nitric oxide metabolism pathway. The urea cycle helps in detoxifying ammonia by converting it into urea which is excreted in urine. In nitric oxide metabolism ASS1 links with nitric oxide synthase to facilitate the use of arginine in producing nitric oxide a vital signaling molecule. The pathway engagement partners include proteins like citrin which supplies aspartate for the reaction.
Associated diseases and disorders	ASS1 has implications in citrullinemia and hepatocellular carcinoma. Citrullinemia results from mutations in the ASS1 gene leading to defective urea cycle function and ammonia accumulation. In cancer particularly hepatocellular carcinoma low expression of ASS1 correlates with poor prognosis as cancer cells often exhibit arginine auxotrophy. Connections through these conditions involve interactions with enzymes such as arginase which also influences arginine availability.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended

We do not recommend this combination. It is not covered by our product promise.


We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

---


20 product images



Flow Cytometry (Intracellular) - Anti-ASS1 antibody [EPR12398] (ab170952)

ASS1 Flow Cytometry (Intracellular) staining using rabbit Anti-ASS1 antibody


Intracellular Flow Cytometry analysis of HeLa (Human epithelial cell line from cervix adenocarcinoma) cells labeling ASS1 with purified ab170952 at 1/100 dilution (10 ug/ml) (red). Cells were fixed with 4% Paraformaldehyde. A Goat anti rabbit IgG (Alexa Fluor® 488) secondary antibody was used at 1/2000 dilution. Isotype control - Rabbit monoclonal IgG (Black). Unlabeled control - Cell without incubation with primary antibody and secondary antibody (Blue).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-ASS1 antibody [EPR12398] (ab170952)

ASS1 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-ASS1 antibody

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of Human kidney tissue sections labeling ASS1 with Purified ab170952 at 1:4000 dilution (0.25 µg/ml). Heat mediated antigen retrieval was performed using citrate Buffer, PH6. Tissue was counterstained with Hematoxylin. ab97051 Goat Anti-Rabbit IgG H&L (HRP) secondary antibody was used at 1:500 dilution. PBS instead of the primary antibody was used as the negative control.




Immunocytochemistry/ Immunofluorescence - Anti-ASS1 antibody [EPR12398] (ab170952)

ASS1 Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-ASS1 antibody

ab170952 staining ASS1 in wild-type HeLa cells (top panel) and ASS1 knockout HeLa cells (ab264989) (bottom panel). The cells were fixed with 100% methanol (5 min) then permeabilized with 0.1% Triton X-100 for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween for 1h. The cells were then incubated with ab170952 at 1/100 dilution and ab7291 (Mouse monoclonal to alpha Tubulin) at 1/1000 dilution overnight at 4°C followed by a


further incubation at room temperature for 1h with a goat secondary antibody to rabbit IgG (Alexa Fluor® 488) (ab150081) at 2 µg/ml (shown in green) and a goat secondary antibody to mouse IgG (Alexa Fluor® 594) (ab150120) at 2 µg/ml (shown in red). Nuclear DNA was labelled in blue with DAPI.
Image was taken with a confocal microscope (Leica-Microsystems TCS SP8).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-ASS1 antibody [EPR12398] (ab170952)

ASS1 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-ASS1 antibody


ab170952 showing -ve staining in Human cervical carcinoma tissue.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-ASS1 antibody [EPR12398] (ab170952)

ASS1 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-ASS1 antibody


ab170952 showing +ve staining in Human normal ureter tissue.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-ASS1 antibody [EPR12398] (ab170952)

ASS1 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-ASS1 antibody


Immunohistochemical analysis of paraffin-embedded Human kidney tissue labeling ASS1 with ab170952 at 1/250 dilution.



Flow Cytometry (Intracellular) - Anti-ASS1 antibody [EPR12398] (ab170952)

ASS1 Flow Cytometry (Intracellular) staining using rabbit Anti-ASS1 antibody


Intracellular flow cytometric analysis of permeabilized Hela cells labeling ASS1 using ab170952 at 1/500 dilution (red) or a rabbit IgG negative (green).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-ASS1 antibody [EPR12398] (ab170952)

ASS1 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-ASS1 antibody


ab170952 showing -ve staining in Human normal uterus tissue.



Immunocytochemistry/ Immunofluorescence - Anti-ASS1 antibody [EPR12398] (ab170952)

ASS1 Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-ASS1 antibody


Immunofluorescent analysis of HeLa cells labeling ASS1 using ab170952 at 1/50 dilution (red). DAPI nuclear staining (blue).



Immunocytochemistry/ Immunofluorescence - Anti-ASS1 antibody [EPR12398] (ab170952)

ASS1 Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-ASS1 antibody


Immunocytochemistry/ Immunofluorescence analysis of MCF7 (Human breast adenocarcinoma cell line) cells labeling ASS1 with Purified ab170952 at 1:100 dilution (10.2µg/ml). Cells were fixed in 4% Paraformaldehyde and permeabilized with 0.1% tritonX-100. Cells were counterstained with ab195889 Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor® 594) 1:200 (2.5 µg/ml). ab150077 Goat anti rabbit IgG(Alexa Fluor® 488) was used as the secondary antibody at 1:1000 dilution. DAPI nuclear counterstain. PBS instead of the primary antibody was used as the secondary antibody only control.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-ASS1 antibody [EPR12398] (ab170952)

ASS1 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-ASS1 antibody

ab170952 showing -ve staining in Human normal pancreas tissue.




Immunoprecipitation - Anti-ASS1 antibody [EPR12398] (ab170952)

Secondary antibody used is HRP-conjugated anti-rabbit IgG preferentially detecting the non-reduced form of rabbit IgG.
All lanes:
Immunoprecipitation - Anti-ASS1 antibody [EPR12398] (ab170952) at 1/10 dilution

Lane 1:
Human fetal liver tissue lysate at 10 µg

Lane 2:
PBS


Predicted band size: 47 kDa



Immunoprecipitation - Anti-ASS1 antibody [EPR12398] (ab170952)

ab170952 (purified) at 1:60 dilution (5ug) immunoprecipitating ASS1 in HeLa (Human cervix adenocarcinoma epithelial cell) whole cell lysate.
Lane 1 (input): HeLa (Human cervix adenocarcinoma epithelial cell) whole cell lysate 10ug
Lane 2 (+): ab170952 & HeLa (Human cervix adenocarcinoma epithelial cell) whole cell lysate
Lane 3 (-): Rabbit monoclonal IgG (ab172730) instead of ab170952 in HeLa (Human cervix adenocarcinoma epithelial cell) whole cell lysate
For western blotting, VeriBlot for IP Detection Reagent (HRP) (ab131366) was used for detection at 1:1000 dilution.
Blocking and diluting buffer: 5% NFDM/TBST.
All lanes:
Immunoprecipitation - Anti-ASS1 antibody [EPR12398] (ab170952)


Predicted band size: 47 kDa



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-ASS1 antibody [EPR12398] (ab170952)

ASS1 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-ASS1 antibody


Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of Mouse kidney tissue sections labeling ASS1 with Purified ab170952 at 1:4000 dilution (0.25 µg/ml). Heat mediated antigen retrieval was performed using citrate Buffer, PH6. Tissue was counterstained with Hematoxylin. ab97051 Goat Anti-Rabbit IgG H&L (HRP)
secondary antibody was used at 1:500 dilution. PBS instead of the primary antibody was used as the negative control.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-ASS1 antibody [EPR12398] (ab170952)

ASS1 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-ASS1 antibody

ab170952 showing +ve staining in Mouse kidney tissue.



Western blot - Anti-ASS1 antibody [EPR12398] (ab170952)

All lanes:
Western blot - Anti-ASS1 antibody [EPR12398] (ab170952) at 1/1000 dilution

Lane 1:
HeLa cell lysate at 10 µg

Lane 2:
Fetal liver tissue lysate at 10 µg


Lane 3:
Fetal kidney tissue lysate at 10 µg

Secondary

All lanes:

Goat anti-rabbit HRP at 1/2000 dilution

Predicted band size: 47 kDa



Western blot - Anti-ASS1 antibody [EPR12398] (ab170952)

ASS1 Western blot staining using rabbit Anti-ASS1 antibody

Blocking and diluting buffer: 5% NFDM/TBST
All lanes:
Western blot - Anti-ASS1 antibody [EPR12398] (ab170952) at 0.05 µg/mL

Lane 1:
HeLa (Human cervix adenocarcinoma epithelial cell) whole cell lysate at 20 µg

Lane 2:
Human fetal liver lysate at 20 µg

Lane 3:
Mouse liver lysate at 20 µg

Lane 4:
Rat liver lysate at 20 µg


Lane 5:
Mouse kidney lysate at 20 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/20000 dilution

Predicted band size: 47 kDa

Observed band size: 47 kDa



Western blot - Anti-ASS1 antibody [EPR12398] (ab170952)

ASS1 Western blot staining using rabbit Anti-ASS1 antibody

Lanes 1 - 4: Merged signal (red and green). Green - ab170952 observed at 47 kDa. Red - loading control, ab9484, observed at 37 kDa.
ab170952 was shown to recognize ASS1 in wild-type HAP1 cells as signal was lost at the expected MW in ASS1 knockout cells. Additional cross-reactive bands were observed in the wild-type and knockout cells. Wild-type and ASS1 knockout samples were subjected to SDS-PAGE. ab170952 and ab9484 (Mouse anti-GAPDH loading control) were incubated overnight at 4°C at 1/20000 dilution and 1/10000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preabsorbed ab216773 and Goat anti-Mouse IgG H&L (IRDye® 680RD) preabsorbed ab216776 secondary antibodies at 1/10000 dilution for 1 hour at room temperature before imaging.

All lanes:
Western blot - Anti-ASS1 antibody [EPR12398] (ab170952) at 1/20000 dilution

Lane 1:
Wild-type HAP1 whole cell lysate at 20 µg

Lane 2:


ASS1 knockout HAP1 whole cell lysate at 20 µg

Lane 3:
HepG2 whole cell lysate at 20 µg

Lane 4:
HeLa whole cell lysate at 20 µg

Predicted band size: 47 kDa

Observed band size: 47 kDa



Western blot - Anti-ASS1 antibody [EPR12398] (ab170952)

Lanes 1- 2: Merged signal (red and green). Green - ab170952 observed at 47 kDa. Red - Anti-GAPDH antibody [6C5] - Loading Control (ab8245) observed at 37 kDa.
ab170952 was shown to react with ASS1 in wild-type HeLa cells in western blot. Loss of signal was observed when knockout cell line ab264989 (knockout cell lysate ab257143) was used. Wild-type HeLa and ASS1 knockout HeLa cell lysates were subjected to SDS-PAGE. Membrane was blocked for 1 hour at room temperature in 0.1% TBST with 3% non-fat dried milk. ab170952 and Anti-GAPDH antibody [6C5] - Loading Control (ab8245) overnight at 4°C at a 1 in 1000 dilution and a 1 in 20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye®800CW) preadsorbed (ab216773) and Goat anti-Mouse IgG H&L (IRDye®680RD) preadsorbed (ab216776) secondary antibodies at 1 in 20000 dilution for 1 hour at room temperature before imaging.

All lanes:
Western blot - Anti-ASS1 antibody [EPR12398] (ab170952) at 1/1000 dilution


Lane 1:
Wild-type HeLa cell lysate at 20 µg

Lane 2:
ASS1 knockout HeLa cell lysate at 20 µg

Performed under reducing conditions.

Predicted band size: 47 kDa

Observed band size: 47 kDa



Western blot - Anti-ASS1 antibody [EPR12398] (ab170952)

ASS1 Western blot staining using rabbit Anti-ASS1 antibody

Blocking and diluting buffer and concentration: 5% NFDM/TBST.
In Western blot, Anti-GAPDH antibody [EPR16891] - Loading Control (ab181602).

Low/negative samples: Caco-2 (PMID: 27689335), PANC-1 (PMID: 31903153), HT-29 (PMID: 27689335).

All lanes:
Western blot - Anti-ASS1 antibody [EPR12398] (ab170952)

Lane 1:
SW480 (human colorectal adenocarcinoma epithelial cell) whole cell lysate at 20 µg

Lane 2:
T-47D (human ductal breast epithelial tumor epithelial cell) whole cell lysate at 20 µg

Lane 3:
HCT116 (human colorectal carcinoma epithelial cell) whole cell lysate at 20 µg

Lane 4:
Caco-2 (human colorectal adenocarcinoma epithelial cell) whole cell lysate at 20 µg

Lane 5:
PANC-1 (human pancreatic epithelioid carcinoma epithelial cell) whole cell lysate at 20 µg

Lane 6:
HT-29 (human colorectal adenocarcinoma epithelial cell) whole cell lysate at 20 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/20000 dilution

Observed band size: 47 kDa, 37 kDa

Exposure time: 60s

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.