abcam

ab173300

Anti-TUFM antibody [EPR12797(B)]

Recombinant | RabMAb | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 50% Tissue culture supernatant, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR12797(B)
Purification technique	Affinity purification Protein A
Concentration	0.089 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

IHC-P

Tested

Species	Human
Dilution info	1/100 - 1/250
Notes	-

Predicted

Species	Mouse, Rat
Dilution info	-
Notes	-

IP

Tested

Species	Human
Dilution info	1/10 - 1/100
Notes	-

Predicted

Species	Mouse, Rat
Dilution info	-
Notes	-

WB

Tested

Species	Human
Dilution info	1/10000 - 1/50000
Notes	-

Predicted

Species	Mouse, Rat
Dilution info	-
Notes	-

ICC/IF

Tested

Species	Human
Dilution info	1/100 - 1/250
Notes	-

Predicted

Species	Mouse, Rat
Dilution info	-
Notes	-

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	1/10 - 1/100
Notes	ab172730 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.

Predicted

Species	Mouse, Rat
Dilution info	-
Notes	-

Target data

See full target information TUFM

Function GTP hydrolase that promotes the GTP-dependent binding of aminoacyl-tRNA to the A-site of ribosomes during protein biosynthesis. Participates in mitochondrial translation (By similarity). Also plays a role in the regulation of autophagy and innate immunity (PubMed:22749352, PubMed:28407488). Recruits ATG5-ATG12 and NLRX1 at mitochondria and serves as a checkpoint of the RIGI-MAVS pathway (PubMed:28407488). In turn, inhibits RLR-mediated type I interferon while promoting autophagy (PubMed:22749352).

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	TUFM also known as elongation factor Tu mitochondrial (EF-Tu) is a protein involved in the protein synthesis process within mitochondria. It has a molecular mass of approximately 50 kDa. TUFM plays an essential role in the translation step of protein synthesis by binding aminoacyl-tRNA and positioning it on the ribosome. Researchers have identified the protein as highly expressed in tissues with high energy demand such as the heart brain and skeletal muscle.
Biological function summary	TUFM assists in mitochondrial protein synthesis which is necessary for the proper function and maintenance of the respiratory chain complexes. It forms part of the mitochondrial ribosome complex operating alongside other elongation factors like EF-G. TUFM's activity is essential for the production of proteins encoded by mitochondrial DNA which are they key components of oxidative phosphorylation machinery.

Pathways	TUFM integrates into the mitochondrial translation pathway which supports the production of proteins critical for the electron transport chain. This pathway is important for ATP production through oxidative phosphorylation. TUFM interacts with other components such as MRPS18 and MRPL3 to mediate the elongation cycle during mitochondrial protein synthesis. These relationships are essential for coordinating mitochondrial function and bioenergetics.
Associated diseases and disorders	TUFM mutations or disruptions have connections to mitochondrial disorders such as MELAS syndrome (Mitochondrial Encephalomyopathy Lactic Acidosis and Stroke-like episodes) which affects energy production. The protein's dysfunction can also relate to Parkinson's Disease where impaired mitochondrial activity plays a role. TUFM's interaction with other mitochondrial proteins like NDUFS1 part of complex I of the electron transport chain underlines its potential impact in these disorders.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.


6 product images



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-TUFM antibody [EPR12797(B)] (ab173300)

TUFM Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-TUFM antibody


Immunohistochemical analysis of paraffin embedded Human breast tissue labeling TUFM with ab173300 at 1/100.
Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.



Flow Cytometry (Intracellular) - Anti-TUFM antibody [EPR12797(B)] (ab173300)

TUFM Flow Cytometry (Intracellular) staining using rabbit Anti-TUFM antibody


Intracellular flow cytometrical analysis of permeabilized Jurkat cells with ab173300 antibody at a dilution of 1/10 (red) or a rabbit IgG (negative) (green).



Immunocytochemistry/ Immunofluorescence - Anti-TUFM antibody [EPR12797(B)] (ab173300)

TUFM Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-TUFM antibody


Immunofluorescent analysis of A431 cells labeling TUFM with ab173300 at 1/100.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-TUFM antibody [EPR12797(B)] (ab173300)

TUFM Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-TUFM antibody

Immunohistochemical analysis of paraffin embedded Human colon tissue labeling TUFM with ab173300 at 1/100.
Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.




Immunoprecipitation - Anti-TUFM antibody [EPR12797(B)] (ab173300)

Immunoprecipitation. ab173300 at 1/10000 staining TUFM immunoprecipitated from HeLa lysate using ab173300 at 1/10.
Lane 2: 1X PBS (negative control).

All lanes:
Immunoprecipitation - Anti-TUFM antibody [EPR12797(B)] (ab173300)

Predicted band size: 50 kDa

Observed band size: 43 kDa



Western blot - Anti-TUFM antibody [EPR12797(B)] (ab173300)

All lanes:
Western blot - Anti-TUFM antibody [EPR12797(B)] (ab173300) at 1/10000 dilution

Lane 1:
HepG2 cell lysate at 10 µg

Lane 2:
A431 cell lysate at 10 µg

Lane 3:
HeLa cell lysate at 10 µg

Lane 4:
Jurkat cell lysate at 10 µg

Predicted band size: 50 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.