abcam

ab174318

Anti-RPL10A antibody [EPR12344]

Recombinant | RabMAb

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR12344
Purification technique	Affinity purification Protein A
Concentration	0.214 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

IHC-P

Tested

Species	Human
Dilution info	1/1000
Notes	For unpurified use at 1/100 - 1/250 dilution. Perform heat-mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.

Species	Mouse
Dilution info	1/1000
Notes	For unpurified use at 1/100 - 1/250 dilution. Perform heat-mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.

Species	Rat
Dilution info	1/1000
Notes	For unpurified use at 1/100 - 1/250 dilution. Perform heat-mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.

IP

Not recommended

Species	Mouse, Rat, Human
Dilution info	-
Notes	-

WB

Tested

Species	Human
Dilution info	1/10000
Notes	For unpurified use at 1/10000 - 1/10000 dilution.

Species	Mouse
Dilution info	1/10000
Notes	For unpurified use at 1/10000 - 1/10000 dilution.

Species	Rat
Dilution info	1/10000
Notes	For unpurified use at 1/10000 - 1/10000 dilution.

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	1/20
Notes	ab172730 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody. For unpurified use at 1/10 - 1/100 dilution.

Expected

Species	Mouse, Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

ICC/IF

Tested

Species	Human
Dilution info	1/500
Notes	For unpurified use at 1/100 dilution.

Expected

Species	Mouse, Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

Target data

See full target information RPL10A

Function	Component of the large ribosomal subunit (PubMed:12962325, PubMed:23636399, PubMed:32669547). The ribosome is a large ribonucleoprotein complex responsible for the synthesis of proteins in the cell (PubMed:12962325, PubMed:23636399, PubMed:32669547).

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.


6 product images



Immunocytochemistry/ Immunofluorescence - Anti-RPL10A antibody [EPR12344] (ab174318)

RPL10A Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-RPL10A antibody


Immunocytochemistry analysis of HeLa (Human cervix adenocarcinoma epithelial cell) cells labeling RPL10A with Purified ab174318 at 1:500 dilution (0.4 µg/ml). Cells were fixed in 4% Paraformaldehyde and permeabilized with 0.1% tritonX-100. Cells were counterstained with ab195889 Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor® 594) 1:200 (2.5 µg/ml). Goat anti rabbit IgG (Alexa Fluor® 488, ab150077) was used as the secondary antibody at 1:1000 (2 µg/ml) dilution. DAPI (blue) was used as nuclear counterstain. PBS instead of the primary antibody was used as the secondary antibody only control.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-RPL10A antibody [EPR12344] (ab174318)

RPL10A Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-RPL10A antibody


Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human colon tissue sections labeling RPL10A with Purified ab174318 at 1:1000 dilution (0.207 µg/ml). Heat mediated antigen retrieval was performed using Perform heat mediated antigen retrieval using ab93684 (Tris/EDTA buffer, pH 9.0). Tissue was counterstained with Hematoxylin. ImmunoHistoProbe one step HRP Polymer (ready to use) secondary antibody was used. PBS instead of the primary antibody was used as the negative control.



Flow Cytometry (Intracellular) - Anti-RPL10A antibody [EPR12344] (ab174318)

RPL10A Flow Cytometry (Intracellular) staining using rabbit Anti-RPL10A antibody


Flow Cytometry analysis of Jurkat (Human T cell leukemia T lymphocyte) cells labelling RPL10A with Purified ab174318 at 1:20 dilution (10 µg/ml) (Red). Cells were fixed with 4% Paraformaldehyde and permeabilised with 90% Methanol. A Goat anti rabbit IgG (Alexa Fluor® 488, ab150077) secondary antibody was used at 1:2000. Isotype control - Rabbit monoclonal IgG (Black). Unlabelled control - Cell without incubation with primary antibody and secondary antibody (Blue).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-RPL10A antibody [EPR12344] (ab174318)

RPL10A Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-RPL10A antibody


Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of mouse liver tissue sections labeling RPL10A with Purified ab174318 at 1:1000 dilution (0.207 µg/ml). Heat mediated antigen retrieval was performed using Perform heat mediated antigen retrieval using ab93684 (Tris/EDTA buffer, pH 9.0). Tissue was counterstained with Hematoxylin. ImmunoHistoProbe one step HRP Polymer (ready to use) secondary antibody was used. PBS instead of the primary antibody was used as the negative control.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-RPL10A antibody [EPR12344] (ab174318)

RPL10A Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-RPL10A antibody

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of rat liver tissue sections labeling RPL10A with Purified ab174318 at 1:1000 dilution (0.207 µg/ml). Heat mediated antigen retrieval was performed using Perform heat mediated antigen retrieval using ab93684 (Tris/EDTA buffer, pH 9.0). Tissue was counterstained with Hematoxylin. ImmunoHistoProbe one step HRP Polymer (ready to use) secondary antibody was used. PBS instead of the primary antibody was used as the negative control.



Western blot - Anti-RPL10A antibody [EPR12344] (ab174318)

RPL10A Western blot staining using rabbit Anti-RPL10A antibody

All lanes:
Western blot - Anti-RPL10A antibody [EPR12344] (ab174318) at 1/10000 dilution

Lane 1:
Jurkat (Human T cell leukemia T lymphocyte) whole cell lysate at 20 µg

Lane 2:
HeLa (Human cervix adenocarcinoma epithelial cell) whole cell lysate at 20 µg

Lane 3:
HepG2 (Human hepatocellular carcinoma epithelial cell) whole cell lysate at 20 µg

Lane 4:
Mouse brain lysate at 20 µg

Lane 5:
Rat brain lysate at 20 µg

Lane 6:
Mouse spleen lysate at 20 µg

Lane 7:
Rat spleen lysate at 20 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/20000 dilution

Predicted band size: 25 kDa

Observed band size: 25 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.