abcam

ab174589

Anti-CD3 epsilon antibody [145-2C11] - Low endotoxin, Azide free

Key facts

Isotype	IgG
Host species	Armenian hamster
Storage buffer	pH: 7.2 Constituents: 0.87% Sodium chloride, 0.16% Sodium phosphate
Form	Liquid
Clonality	Monoclonal
Carrier free	Yes
Clone number	145-2C11
Purification technique	Affinity purification
Concentration	2 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

IP

Expected

Species	Mouse
Dilution info	Use at an assay dependent concentration.
Notes	-

Flow Cyt

Expected

Species	Mouse
Dilution info	Use at an assay dependent concentration.
Notes	-

WB

Expected

Species	Mouse
Dilution info	Use at an assay dependent concentration.
Notes	-

IF

Expected

Species	Mouse
Dilution info	Use at an assay dependent concentration.
Notes	-

FuncS (Act)

Expected

Species	Mouse
Dilution info	Use at an assay dependent concentration.
Notes	-

IHC-Fr

Expected

Species	Mouse
Dilution info	Use at an assay dependent concentration.
Notes	-

Target data

See full target information Cd3e

Function	Part of the TCR-CD3 complex present on T-lymphocyte cell surface that plays an essential role in adaptive immune response (PubMed:9843989, PubMed:10952314, PubMed:19956738, PubMed:24470497, PubMed:24899501, PubMed:40592325). When antigen presenting cells (APCs) activate T-cell receptor (TCR), TCR-mediated signals are transmitted across the cell membrane by the CD3 chains CD3D, CD3E, CD3G and CD247/CD3Z (PubMed:10952314, PubMed:19956738, PubMed:24899501, PubMed:40592325). All CD3 chains contain immunoreceptor tyrosine-based activation motifs (ITAMs) in their cytoplasmic domain (PubMed:10952314, PubMed:19956738, PubMed:24899501, PubMed:40592325). Upon TCR engagement, these motifs become phosphorylated by Src family protein tyrosine kinases LCK and FYN, resulting in the activation of downstream signaling pathways (PubMed:10952314, PubMed:19956738, PubMed:24899501, PubMed:40592325). CD3E ITAM phosphorylation creates docking sites for the protein kinase ZAP70 leading to ZAP70 phosphorylation and its conversion into a catalytically active enzyme (PubMed:10952314). In addition of this role of signal transduction in T-cell activation, CD3E plays an essential role in correct T-cell development (PubMed:10952314, PubMed:19956738, PubMed:24899501). Also participates in internalization and cell surface down-regulation of TCR-CD3 complexes via endocytosis sequences present in CD3E cytosolic region (By similarity). In addition to its role as a TCR coreceptor, it serves as a receptor for ITPRIPL1 (By similarity). Ligand recognition inhibits T-cell activation by promoting interaction with NCK1, which prevents CD3E-ZAP70 interaction and blocks the ERK-NFkB signaling cascade and calcium influx (By similarity).

Storage

Shipped at conditions	Blue Ice
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended

We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.