abcam

ab175932

Anti-NRF1 antibody [EPR5554(N)] - ChIP Grade

Anti-NRF1 antibody [EPR5554(N)] - ChIP Grade (ab175932) is a rabbit monoclonal antibody detecting NRF1 in **Western Blot, Flow Cytometry (Intra), IP, IHC-P, ICC/IF, ChIP**. Suitable for **Human, Mouse, Rat**.

- Biophysical QC for unrivalled batch-batch consistency
- Over 70 publications

Recombinant | RabMAb | Advanced Validation | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR5554(N)
Purification technique	Affinity purification Protein A
Concentration	0.761 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

ChIC/CUT&RUN-seq

Tested

Species	Human

Dilution info	5 µg
Notes	-

Expected

Species	Mouse
Dilution info	5 µg
Notes	-

Species	Rat
Dilution info	5 µg
Notes	-

IHC-P

Tested

Species	Human
Dilution info	1/50 - 1/100
Notes	-

Expected

Species	Mouse, Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

IP

Tested

Species	Human
Dilution info	1/10 - 1/100
Notes	-

Expected

Species	Mouse, Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

ChIP

Tested

Species	Mouse, Human
Dilution info	-
Notes	-

Expected

Species	Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

WB

Tested

Species	Mouse
Dilution info	1/1000 - 1/10000
Notes	-

Species	Rat
Dilution info	1/1000 - 1/10000
Notes	-

Species	Human
Dilution info	1/1000 - 1/10000
Notes	-

ICC/IF

Tested

Species	Human
Dilution info	1/50 - 1/100
Notes	-

Expected

Species	Mouse, Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

ChIP-seq

Tested

Species	Human
Dilution info	8 µg for 10^7 Cells
Notes	-

Expected

Species	Mouse, Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	1/10 - 1/150
Notes	ab172730 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.

Expected

Species	Mouse, Rat

Dilution info	Use at an assay dependent concentration.
Notes	-

Target data

See full target information NRF1

Function	Transcription factor that activates the expression of the EIF2S1 (EIF2-alpha) gene. Links the transcriptional modulation of key metabolic genes to cellular growth and development. Implicated in the control of nuclear genes required for respiration, heme biosynthesis, and mitochondrial DNA transcription and replication.

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

What is this antibody validated in?
Anti-NRF1 antibody [EPR5554(N)] - ChIP Grade (ab175932) is a rabbit recombinant monoclonal antibody and is validated for use in Western Blot (WB), Flow Cytometry (Intra), Immunoprecipitation (IP), Immunohistochemistry (IHC-P), Immunocytochemistry/immunofluorescence (ICC/IF), ChIP in Human, Mouse, Rat samples.

What is the molecular weight of NRF1?
Anti-NRF1 [EPR5554(N)] - ChIP Grade (ab175932) specifically detects a band for NRF1 (UniProt: Q16656) at a molecular weight of 54kDa.

Trusted by the scientific community
Anti-NRF1 [EPR5554(N)] - ChIP Grade (ab175932) was first used in a scientific publication in 2013 and has been cited over 70 times in peer-reviewed journals.

Other related products
We have a range of other formats of antibody clone [EPR5554(N)] also available for your convenience: ab175932,

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Trial sizes available!
Test your antibody or perform pre-screening before committing to a larger quantity. Sold in 20µl. Discover our selection of trial-size antibodies.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	The nuclear respiratory factor 1 (NRF1) also known as Alpha-Pal is a transcription factor with an approximate molecular weight of 68 kDa. It plays an important role in the regulation of genes required for mitochondrial DNA transcription and replication. Researchers find NRF1 expression in various tissues including muscle brain and liver. This protein functions mechanistically by binding to specific promoter regions of target genes to control their expression making it essential for maintaining cellular energy metabolism.
Biological function summary	NRF1 influences the expression of genes involved in the mitochondrial function and is part of the transcriptional complex. This complex includes the NRF2 protein where NRF1 partners with co-activators to modulate gene transcription. NRF1 promotes the expression of components essential for mitochondrial biogenesis and cellular respiration which are vital processes for energy production in cells. Therefore NRF1 contributes to maintaining cellular energy homeostasis and overall cellular health.
Pathways	NRF1 integrates into critical regulatory pathways for mitochondrial function and metabolism. NRF1 links closely to the PGC-1α pathway which is instrumental in the regulation of energy metabolism and mitochondrial biogenesis. Additionally NRF1 works in conjunction with NRF2 to regulate oxidative stress responses. These interactions ensure a coordinated cellular response to energy demands and help protect the cell from oxidative damage.
Associated diseases and disorders	NRF1 associates with neurodegenerative diseases and metabolic syndromes. Altered NRF1 regulation can impact diseases such as Alzheimer's disease where it may contribute to neurodegeneration through impaired mitochondrial function. Moreover the dysfunction of NRF1 might relate to metabolic disorders like diabetes as energy metabolism and mitochondrial health are disrupted. NRF1

connects with proteins like SIRT1 in these contexts further implicating NRF1 in these pathological states through its role in energy homeostasis and stress response.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.


18 product images



Immunocytochemistry/ Immunofluorescence - Anti-NRF1 antibody [EPR5554(N)] - ChIP Grade (ab175932)

NRF1 Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-NRF1 antibody


Immunocytochemistry/Immunofluorescence analysis of HeLa cells labeling NRF1 with unpurified ab175932 at a dilution of 1/50.



Flow Cytometry (Intracellular) - Anti-NRF1 antibody [EPR5554(N)] - ChIP Grade (ab175932)

NRF1 Flow Cytometry (Intracellular) staining using rabbit Anti-NRF1 antibody


Intracellular Flow Cytometry analysis of 293T cells labelling NRF1 with purified ab175932 at a dilution of 1/150 (red). Cells were fixed with 80% methanol. A FITC-conjugated goat anti-rabbit IgG (1/500) was used as the secondary antibody. Black - Isotype control, rabbit monoclonal IgG. Blue - Unlabelled control, cells without incubation with primary and secondary antibodies.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-NRF1 antibody [EPR5554(N)] - ChIP Grade (ab175932)

NRF1 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-NRF1 antibody

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human skeletal muscle tissue labeling NRF1 with unpurified ab175932 at a dilution of 1/50.
Perform heat mediated antigen retrieval before commencing with IHC staining protocol.




Immunocytochemistry/ Immunofluorescence - Anti-NRF1 antibody [EPR5554(N)] - ChIP Grade (ab175932)

NRF1 Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-NRF1 antibody

Immunocytochemistry/Immunofluorescence analysis of MCF-7 cells labelling NRF1 with purified ab175932 at a dilution of 1/100. Cells were fixed with 4% paraformaldehyde and permeabilized with 0.1% Triton X-100. ab150077, an Alexa Fluor® 488-conjugated goat anti-rabbit IgG (1/1000) was used as the secondary antibody. DAPI (blue) was used as the nuclear counterstain. ab7291, a mouse anti-tubulin (1/1000) and ab150120, an Alexa Fluor® 594-conjugated goat anti-mouse IgG (1/1000) were also used.
Control 1: primary antibody (1/100) and secondary antibody, ab150120, an Alexa Fluor® 594-conjugated goat anti-mouse IgG (1/1000).


Control 2: ab7291 (1/1000) and secondary antibody, ab150077, an Alexa Fluor® 488-conjugated goat anti-rabbit IgG (1/1000).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-NRF1 antibody [EPR5554(N)] - ChIP Grade (ab175932)

NRF1 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-NRF1 antibody


Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human gastric adenocarcinoma tissue labeling NRF1 with unpurified ab175932 at a dilution of 1/50.
Perform heat mediated antigen retrieval before commencing with IHC staining protocol.



Flow Cytometry (Intracellular) - Anti-NRF1 antibody [EPR5554(N)] - ChIP Grade (ab175932)

NRF1 Flow Cytometry (Intracellular) staining using rabbit Anti-NRF1 antibody


Intracellular flow cytometric analysis of permeabilized 293T cells labeling NRF1 with unpurified ab175932 at a dilution of 1/10 (red) compared to a negative control (rabbit IgG,green).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-NRF1 antibody [EPR5554(N)] - ChIP Grade (ab175932)


NRF1 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-NRF1 antibody

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human cevical carcinoma tissue labelling NRF1 with purified ab175932 at a dilution of 1/100. Heat mediated antigen retrieval was performed using EDTA buffer pH 9. ab97051, a HRP-conjugated goat anti-rabbit IgG (H+L) was used as the secondary antibody (1/500). Negative control using PBS instead of primary antibody. Counterstained with hematoxylin.




ChIP - Anti-NRF1 antibody [EPR5554(N)] - ChIP Grade (ab175932)

Chromatin was prepared from Hela cells according to the Abcam Dual-X-ChIP protocol. Cells were fixed with EGS for 30 minutes, then formaldehyde for 10 minutes.
The ChIP was performed with 25 µg of chromatin, 5 µg of ab175932 (red), and 20 µl of Protein A/G sepharose beads. 5 µg of rabbit normal IgG was added to the beads control (gray). The immunoprecipitated DNA was quantified by real time PCR (Sybr green approach).
Primers and probes are located in the first kb of the transcribed region.



ChIP-sequencing - Anti-NRF1 antibody [EPR5554(N)] - ChIP Grade (ab175932)


Chromatin was prepared from HeLa cells. Cells were fixed with 1% formaldehyde for 10 minutes. ChIP was performed with 10^7 HeLa cells and 8 µg of ab175932 [EPR5554(N)]. ChIP DNA was sequenced on the Illumina NovaSeq 6000 to a depth of 30 million reads. Additional screenshots of mapped reads can be downloaded here.



Immunoprecipitation - Anti-NRF1 antibody [EPR5554(N)] - ChIP Grade (ab175932)

ab175932 (unpurified) at a dilution of 1/10 immunoprecipitating NRF1 in 293T cell lysate.
All lanes:
Immunoprecipitation - Anti-NRF1 antibody [EPR5554(N)] - ChIP Grade (ab175932)

Predicted band size: 54 kDa



Immunoprecipitation - Anti-NRF1 antibody [EPR5554(N)] - ChIP Grade (ab175932)

ab175932 (purified) at a dilution of 1/50 immunoprecipitating NRF1 in 293T whole cell lysate.
Lane 1 (input): 293T whole cell lysate (10µg)

Lane 2 (+): ab175932 + 293T whole cell lysate.

Lane 3 (-): Rabbit monoclonal IgG (ab172730) instead of ab175932 in 293T whole cell lysate.

For western blotting, VeriBlot for IP Detection Reagent (HRP) (ab131366), was used for detection at 1/10,000 dilution.


Blocking buffer and concentration: 5% NFDM/TBST.

Diluting buffer and concentration: 5% NFDM /TBST.

All lanes:
Immunoprecipitation - Anti-NRF1 antibody [EPR5554(N)] - ChIP Grade (ab175932)


Predicted band size: 54 kDa

Observed band size: 68 kDa



ChIP - Anti-NRF1 antibody [EPR5554(N)] - ChIP Grade (ab175932)

Chromatin was prepared from NIH/3T3 treated with MG-132(2uM 16h) cells according to the Abcam Dual-X-ChIP protocol.Cells were fixed with EGS for 30 minutes, then formaldehyde for 10 minutes.
The ChIP was performed with 25 µg of chromatin, 5 µg of ab175932 (red), and 20 µl of Protein A/G sepharose beads. 5 µg of rabbit normal IgG was added to the beads control (gray). The immunoprecipitated DNA was quantified by real time PCR (Sybr green approach).
Primers and probes are located in the first kb of the transcribed region.



Western blot - Anti-NRF1 antibody [EPR5554(N)] - ChIP Grade (ab175932)

Blocking and dilution buffer: 5% NFDM/TBST
All lanes:
Western blot - Anti-NRF1 antibody [EPR5554(N)] - ChIP Grade (ab175932) at 1/10000 dilution


All lanes:
HEK293 whole cell lysate at 20 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/100000 dilution

Predicted band size: 54 kDa

Observed band size: 68 kDa



Western blot - Anti-NRF1 antibody [EPR5554(N)] - ChIP Grade (ab175932)

Blocking and dilution buffer: 5% NFDM/TBST
All lanes:
Western blot - Anti-NRF1 antibody [EPR5554(N)] - ChIP Grade (ab175932) at 1/10000 dilution


All lanes:
HeLa whole cell lysate at 20 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/100000 dilution


Predicted band size: 54 kDa

Observed band size: 68 kDa



ChIC/CUT&RUN sequencing - Anti-NRF1 antibody [EPR5554(N)] - ChIP Grade (ab175932)


CUT&RUN was performed using a ChIC/CUT&RUN pAG-MNAse, 2.5x10^5 HeLa cells, and 5µg of ab175932 [EPR5554(N)]. The resulting DNA was sequenced on the NovaSeq 6000 (PE150). Additional screenshots of mapped reads can be downloaded here.



ChIC/CUT&RUN sequencing - Anti-NRF1 antibody [EPR5554(N)] - ChIP Grade (ab175932)

ChIC/CUT&RUN was performed using a pAG-MNAse at a final concentration of 700 ng/mL, 2 x 10^5 HeLa (Human cervix adenocarcinoma epithelial cell line) cells and 5 µg of ab175932 [EPR5554(N)]. The resulting DNA was sequenced on the Illumina NovaSeq 6000 to a depth of 10 million reads. The negative IgG control ab172730 is also shown. The ChIP data was conducted on chromatin prepared from HeLa cells. Cells were fixed with 1% formaldehyde for 10 minutes. ChIP was

performed with 10^7 HeLa cells and 8 µg of ab175932. ChIP DNA was sequenced on the Illumina NovaSeq 6000 to a depth of 30 million reads. Additional screenshots of mapped reads can be found in the Protocol booklet in the Product Protocol section. The University of Geneva owns patents relevant to ChIC (Chromatin Immuno-Cleavage) methods.



Western blot - Anti-NRF1 antibody [EPR5554(N)] - ChIP Grade (ab175932)

Blocking and dilution buffer: 5% NFDM/TBST
All lanes:
Western blot - Anti-NRF1 antibody [EPR5554(N)] - ChIP Grade (ab175932) at 1/5000 dilution

Lane 1:
Mouse heart tissue lysate at 20 µg

Lane 2:
Mouse brain tissue lysate at 20 µg

Lane 3:
Rat heart tissue lysate at 20 µg


Lane 4:
Rat brain tissue lysate at 20 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/20000 dilution

Predicted band size: 54 kDa

Observed band size: 68 kDa



Western blot - Anti-NRF1 antibody [EPR5554(N)] - ChIP Grade (ab175932)

All lanes:
Western blot - Anti-NRF1 antibody [EPR5554(N)] - ChIP Grade (ab175932) at 1/1000 dilution

Lane 1:
MCF-7 cell lysate at 10 µg

Lane 2:
Hela cell lysate at 10 µg

Lane 3:
Human fetal heart tissue lysate at 10 µg

Lane 4:
293T cell lysate at 10 µg

Predicted band size: 54 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.