abcam

ab179519

Anti-Caspase-2L antibody [EPR16790]

Recombinant | RabMAb

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR16790
Purification technique	Affinity purification Protein A
Concentration	2.48 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

WB

Tested

Species	Mouse
Dilution info	1/2000
Notes	-

Species	Rat

Dilution info	1/2000
Notes	-

Species	Human
Dilution info	1/2000
Notes	-

ICC/IF

Tested

Species	Human
Dilution info	1/500
Notes	-

Expected

Species	Mouse, Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	1/500
Notes	ab172730 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.

Expected

Species	Mouse, Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

IHC-P

Tested

Species	Human
Dilution info	1/250
Notes	-

Expected

Species	Mouse, Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

Target data

See full target information Casp2

Function	Involved in the activation cascade of caspases responsible for apoptosis execution. Might function by either activating some proteins required for cell death or inactivating proteins necessary for cell survival (PubMed:7958843). Associates with PIDD1 and CRADD to form the PIDDosome, a complex that activates CASP2 and triggers apoptosis in response to genotoxic stress (By similarity).

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.


7 product images



Immunocytochemistry/ Immunofluorescence - Anti-Caspase-2L antibody [EPR16790] (ab179519)

Caspase-2L Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-Caspase-2L antibody


Immunofluorescent analysis of 4% paraformaldehyde-fixed 0.1% Triton X-100 permeabilized HeLa (Human epithelial cells from cervix adenocarcinoma) cells labeling Caspase-2L with ab179519 at 1/500 dilution followed by Goat anti-rabbit IgG (Alexa Fluor® 488) (ab150077) secondary antibody at 1/500 dilution (green). Confocal image showing nuclear and cytoplasmic staining on HeLa cells. The nuclear counter stain is DAPI (blue). Tubulin is detected with ab7291 (anti-Tubulin mouse mAb) at 1/1000 dilution and ab150120 (AlexaFluor®594 Goat anti-Mouse secondary) at 1/500 dilution (red).
The negative controls are as follows:
1. ab179519 at 1/500 dilution followed by ab150120 (AlexaFluor®594 Goat anti-Mouse secondary) at 1/500 dilution.
2. ab7291 (anti-Tubulin mouse mAb) at 1/1000 dilution followed by ab150077 (Alexa Fluor®488 Goat Anti-Rabbit IgG H&L) at 1/500 dilution.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Caspase-2L antibody [EPR16790] (ab179519)

Caspase-2L Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-Caspase-2L antibody


Immunohistochemical analysis of paraffin-embedded Human lung cancer tissue labeling Caspase-2L using ab179519 at 1/250 dilution. A Goat Anti-Rabbit IgG H&L (HRP) (ab97051) was used as secondary at 1/500 dilution. Counterstain: Hematoxylin.
Inset image: negative control obtained using PBS instead of ab179519.
Note: nuclear staining on the tumor cells of Human lung cancer was observed.
Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Caspase-2L antibody [EPR16790] (ab179519)

Caspase-2L Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-Caspase-2L antibody


Immunohistochemical analysis of paraffin-embedded Human testis tissue labeling Caspase-2L using ab179519 at 1/250 dilution. A Goat Anti-Rabbit IgG H&L (HRP) (ab97051) was used as secondary at 1/500 dilution. Counterstain: Hematoxylin.
Inset image: negative control obtained using PBS instead of ab179519.
Note: nuclear staining on the germ cells of the Human testis was observed.
Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.



Flow Cytometry (Intracellular) - Anti-Caspase-2L antibody [EPR16790] (ab179519)

Caspase-2L Flow Cytometry (Intracellular) staining using rabbit Anti-Caspase-2L antibody

Intracellular flow cytometric analysis of 2% paraformaldehyde-fixedJurkat (Human T cell leukemia cells from peripheral blood) cells labeling Caspase-2 with ab179519 at 1/500 dilution (red) compared with a rabbit monoclonal IgG isotype control (black) and a unlabelled control (cells without incubation with primary antibody and secondary antibody; blue). Goat anti rabbit IgG (FITC) at 1/150 dilution was used as the secondary antibody.



Western blot - Anti-Caspase-2L antibody [EPR16790] (ab179519)

Caspase-2L Western blot staining using rabbit Anti-Caspase-2L antibody

Blocking/Dilution buffer: 5% NFDM/TBST.
Observed MW: 48, 30, 14, 12 kDa

All lanes:
Western blot - Anti-Caspase-2L antibody [EPR16790] (ab179519) at 1/2000 dilution

Lane 1:
Jurkat (Human T cell leukemia cells from peripheral blood) whole cell lysates treated with 1uM staurosporine for 4 hours at 10 µg


Lane 2:
untreated Jurkat whole cell lysates at 10 µg

Secondary

All lanes:
Goat Anti-Rabbit IgG, (H+L),Peroxidase conjugated at 1/1000 dilution

Predicted band size: 51 kDa

Exposure time: 3min



Western blot - Anti-Caspase-2L antibody [EPR16790] (ab179519)

Caspase-2L Western blot staining of C6 (Rat glial tumor cells) whole cell lysate using rabbit Anti-Caspase-2L antibody

Blocking/Dilution buffer: 5% NFDM/TBST.
Observed MW: 48,12 kDa

All lanes:
Western blot - Anti-Caspase-2L antibody [EPR16790] (ab179519) at 1/10000 dilution


All lanes:
C6 (Rat glial tumor cells) whole cell lysate at 10 µg

Secondary

All lanes:
Goat anti-rabbit IgG, (H+L), peroxidase conjugated at 1/1000 dilution

Predicted band size: 51 kDa

Exposure time: 3min



Western blot - Anti-Caspase-2L antibody [EPR16790] (ab179519)

Caspase-2L Western blot staining of NIH/3T3 (Mouse embyro fibroblast cells) whole cell lysates using rabbit Anti-Caspase-2L antibody

Blocking/Dilution buffer: 5% NFDM/TBST.
Observed MW: 48,30,12 kDa

All lanes:
Western blot - Anti-Caspase-2L antibody [EPR16790] (ab179519) at 1/2000 dilution

All lanes:
NIH/3T3 (Mouse embyro fibroblast cells) whole cell lysates at 10 µg

Secondary

All lanes:
Goat Anti-Rabbit IgG, (H+L),Peroxidase conjugated at 1/1000 dilution

Predicted band size: 51 kDa

Exposure time: 3min

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.