abcam

ab181115

Anti-Raf1 antibody [EP4969] - N-terminal

Anti-Raf1 antibody [EP4969] - N-terminal (ab181115) is a rabbit monoclonal antibody detecting Raf1 in **Western Blot, Flow Cytometry (Intra), ICC/IF**. Suitable for **Human, Mouse, Rat**.

- KO validated for confirmed specificity
- Biophysical QC for unrivalled batch-batch consistency

Lab Essentials | Recombinant | RabMAb | KO Validated | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EP4969
Purification technique	Affinity purification Protein A
Concentration	1.129 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

WB

Tested

Species	Mouse

Dilution info	1/1000
Notes	-

Species	Rat
Dilution info	1/1000
Notes	-

Species	Human
Dilution info	1/1000
Notes	-

ICC/IF

Tested

Species	Human
Dilution info	1/100 - 1/250
Notes	-

Expected

Species	Mouse, Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	1/100
Notes	-

Expected

Species	Mouse, Rat
Dilution info	Use at an assay dependent concentration.

Notes	-

Target data

See full target information RAF1

Function	Serine/threonine-protein kinase that acts as a regulatory link between the membrane-associated Ras GTPases and the MAPK/ERK cascade, and this critical regulatory link functions as a switch determining cell fate decisions including proliferation, differentiation, apoptosis, survival and oncogenic transformation. RAF1 activation initiates a mitogen-activated protein kinase (MAPK) cascade that comprises a sequential phosphorylation of the dual-specific MAPK kinases (MAP2K1/MEK1 and MAP2K2/MEK2) and the extracellular signal-regulated kinases (MAPK3/ERK1 and MAPK1/ERK2). The phosphorylated form of RAF1 (on residues Ser-338 and Ser-339, by PAK1) phosphorylates BAD/Bcl2-antagonist of cell death at 'Ser-75'. Phosphorylates adenylyl cyclases: ADCY2, ADCY5 and ADCY6, resulting in their activation. Phosphorylates PPP1R12A resulting in inhibition of the phosphatase activity. Phosphorylates TNNT2/cardiac muscle troponin T. Can promote NF-kB activation and inhibit signal transducers involved in motility (ROCK2), apoptosis (MAP3K5/ASK1 and STK3/MST2), proliferation and angiogenesis (RB1). Can protect cells from apoptosis also by translocating to the mitochondria where it binds BCL2 and displaces BAD/Bcl2-antagonist of cell death. Regulates Rho signaling and migration, and is required for normal wound healing. Plays a role in the oncogenic transformation of epithelial cells via repression of the TJ protein, occludin (OCLN) by inducing the up-regulation of a transcriptional repressor SNAI2/SLUG, which induces down-regulation of OCLN. Restricts caspase activation in response to selected stimuli, notably Fas stimulation, pathogen-mediated macrophage apoptosis, and erythroid differentiation.

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

What is this antibody validated in?
Anti-Raf1 antibody [EP4969] - N-terminal (ab181115) is a rabbit recombinant monoclonal antibody and is validated for use in Western Blot (WB), Flow Cytometry (Intra), Immunocytochemistry/immunofluorescence (ICC/IF) in Human, Mouse, Rat samples.

What is the molecular weight of Raf1?
Anti-Raf1 [EP4969] - N-terminal (ab181115) specifically detects a band for Raf1 (UniProt: P04049) at a molecular weight of 73kDa.

Specificity confirmed
The specificity of Anti-Raf1 antibody [EP4969] - N-terminal (ab181115) has been confirmed by Western blot testing in RAF1 Knockout HAP1 cells.

Other related products
We have a range of other formats of antibody clone [EP4969] also available for your convenience: ab181115, Carrier free - ab236003, APC - ab310836, PE - ab310911, Alexa Fluor® 594 - ab311715, Alexa Fluor® 568 - ab312991, Alexa Fluor® 555 - ab313196

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Trial sizes available!
Test your antibody or perform pre-screening before committing to a larger quantity. Sold in 20µl. Discover our selection of trial-size antibodies.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	Raf1 also known as c-Raf or Raf-1 is a serine/threonine-protein kinase that plays an important role in cell division differentiation and survival. Its molecular weight is approximately 74 kDa. Raf1 is part of the MAPK/ERK signaling pathway and is ubiquitously expressed in many tissues including the brain heart and liver. As a component of the signaling cascade its activation translates extracellular signals into intracellular responses.
Biological function summary	Raf1 regulates important cellular processes by activating downstream kinases in response to external stimuli. Raf1 forms a complex with other proteins such as Ras facilitating its role as an essential component of the MAPK/ERK pathway. Upon activation by Ras Raf1 phosphorylates and activates MEK1 and MEK2 which in turn

	activate the ERK1 and ERK2. This signaling axis is involved in controlling gene expression and cellular proliferation.
Pathways	Raf1 is integrally involved in the MAPK/ERK pathway which is critical for transducing signals from growth factors and mitogens. It relates closely with proteins such as Ras MEK and ERK in this pathway. The pathway is important for regulating cellular responses to various stimuli and is particularly involved in processes such as cell cycle control and apoptosis.
Associated diseases and disorders	Raf1 has associations with certain types of cancer and Noonan syndrome. In various cancers mutations or dysregulation of Raf1 or related components such as Ras can lead to uncontrolled cellular proliferation. In Noonan syndrome Raf1 mutations result in anomalies in the Ras/MAPK pathway which can impact normal development. Raf1's interactions in these pathways indicate its relevance as a potential target for therapeutic intervention.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.


12 product images



Flow Cytometry (Intracellular) - Anti-Raf1 antibody [EP4969] - N-terminal (ab181115)

Raf1 Flow Cytometry (Intracellular) staining using rabbit Anti-Raf1 antibody


Intracellular Flow Cytometry analysis of K-562 (Human chronic myelogenous leukemia lymphoblast) cells labelling with unpurified ab181115 at 1/100 dilution (10.79µg/mL) (red). Cells were fixed with 4% paraformaldehyde . Goat anti rabbit IgG (Alexa Fluor® 488, ab150077) was used as the secondary antibody at 1/2000 dilution. Isotype control - 90% methanol . Unlabeled control - Rabbit monoclonal IgG (ab172730) / Black.



Immunocytochemistry/ Immunofluorescence - Anti-Raf1 antibody [EP4969] - N-terminal (ab181115)

Raf1 Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-Raf1 antibody


Immunofluorescent analysis of acetone-fixed K562 cells labeling Raf1 with unpurified ab181115 at 1/250 dilution, followed by Goat anti rabbit IgG (Alexa Fluor® 488) secondary antibody at 1/200 dilution. Counter stained with Dapi.



Immunocytochemistry/ Immunofluorescence - Anti-Raf1 antibody [EP4969] - N-terminal (ab181115)

Raf1 Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-Raf1 antibody

Immunocytochemistry/ Immunofluorescence analysis of HeLa (Human cervix adenocarcinoma epithelial cell) cells labeling Raf1 with purified ab181115 at 1/100 dilution (10 µg/ml). Cells were fixed in 4% Paraformaldehyde and permeabilized with 0.1% tritonX-100. Cells were counterstained with ab195889 Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor® 594) 1/200 (2.5 µg/ml). Goat anti rabbit IgG (Alexa Fluor® 488, ab150077) was used as the secondary antibody at 1/1000 (2 µg/ml) dilution. DAPI (blue) was used as nuclear counterstain. PBS instead of the primary antibody was used as the secondary antibody only control.



Western blot - Anti-Raf1 antibody [EP4969] - N-terminal (ab181115)

Raf1 Western blot staining using rabbit Anti-Raf1 antibody

Lanes 1- 2: Merged signal (red and green). Green - ab181115 observed at 73 kDa. Red - Anti-GAPDH antibody [6C5] - Loading Control (ab8245) observed at 37 kDa.
ab181115 was shown to react with Raf1 in wild-type HeLa cells in western blot. Loss of signal was observed when knockout cell line ab264978 (knockout cell lysate ab257126) was used. Wild-type HeLa and Raf1 knockout HeLa cell lysates were subjected to SDS-PAGE. Membrane was blocked for 1 hour at room temperature in 0.1% TBST with 3% non-fat dried milk. ab181115 and Anti-GAPDH antibody [6C5] - Loading Control (ab8245) overnight at 4°C at a 1 in 20000 dilution and a 1 in 20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye®800CW) preadsorbed (ab216773) and Goat anti-Mouse IgG H&L (IRDye®680RD) preadsorbed (ab216776) secondary antibodies at 1 in 20000 dilution for 1 hour at room temperature before imaging.

All lanes:
Western blot - Anti-Raf1 antibody [EP4969] - N-terminal (ab181115) at 1/20000 dilution

Lane 1:
Wild-type HeLa cell lysate at 20 µg


Lane 2:
RAF1 knockout HeLa cell lysate at 20 µg

Lane 2:
Western blot - Human RAF1 knockout HeLa cell line (ab264978)

Performed under reducing conditions.

Predicted band size: 73 kDa

Observed band size: 73 kDa



Western blot - Anti-Raf1 antibody [EP4969] - N-terminal (ab181115)


Raf1 Western blot staining using rabbit Anti-Raf1 antibody

Lanes 1: Wild-type HAP1 cell lysate (20 µg)
Lanes 2: Raf1 knockout HAP1 cell lysate (20 µg)
Lanes 1 and 2: Merged signal (red and green).
Green - Target observed at 74 kDa. Red - loading control, ab8226, observed at 42 kDa

This western blot image is a comparison between unpurified ab181115 and a competitor's top cited discontinued rabbit polyclonal antibody.

All lanes:
Western blot - Anti-Raf1 antibody [EP4969] - N-terminal (ab181115)

Predicted band size: 73 kDa



Western blot - Anti-Raf1 antibody [EP4969] - N-terminal (ab181115)

All lanes:
Western blot - Anti-Raf1 antibody [EP4969] - N-terminal (ab181115) at 1/1000 dilution

Lane 1:
HeLa cell lysate at 20 µg


Lane 2:
K562 cell lysate at 20 µg

Secondary

All lanes:
Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugate at 1/1000 dilution

Predicted band size: 73 kDa

Observed band size: 73 kDa



Western blot - Anti-Raf1 antibody [EP4969] - N-terminal (ab181115)

Raf1 Western blot staining using rabbit Anti-Raf1 antibody

Western blot: Anti-RAF1 antibody [EP4969] (ab181115) staining at 1/1000 dilution, shown in green; Mouse anti-GAPDH antibody [6C5] (ab8245) loading control staining at 1/20000 dilution, shown in red. In Western blot, ab181115 was shown to bind specifically to RAF1. A band was observed at 73 kDa in wild-type HCT 116 cell lysates with no signal observed at this size in RAF1 knockout cell line. To generate this image, wild-type and RAF1 knockout HCT 116 cell lysates were analysed. First, samples were run on an SDS-PAGE gel then transferred onto a nitrocellulose membrane. Membranes were blocked in 3 % milk in TBS-0.1 % Tween® 20 (TBS-T) before incubation with primary antibodies overnight at 4 °C. Blots were washed four times in TBS-T, incubated with secondary antibodies for 1 h at room temperature, washed again four times then imaged. Secondary antibodies used were Goat anti-Rabbit IgG H&L 800CW and Goat anti-Mouse IgG H&L 680RD at 1/20000 dilution.
All lanes:
Western blot - Anti-Raf1 antibody [EP4969] - N-terminal (ab181115) at 1/1000 dilution

Lane 1:
Wild-type HCT 116 cell lysate at 20 µg

Lane 2:
RAF1 knockout HCT 116 cell lysate at 20 µg


Secondary

All lanes:
Goat anti-Rabbit IgG H&L 800CW and Goat anti-Mouse IgG H&L 680RD at 1/20000 dilution

Performed under reducing conditions.

Predicted band size: 73 kDa

Observed band size: 73 kDa



Western blot - Anti-Raf1 antibody [EP4969] - N-terminal (ab181115)

ab181602 was used as GAPDH loading control. ab181115 was used to detect Raf1. ab150365 was used to detect Raf1 phospho S43.
All lanes:
Purified at 1/5000 dilution

Lane 1:
HEK-293 (Human embryonic kidney epithelial cell) whole cell lysates at 15 µg

Lane 2:
HEK-293 (Human embryonic kidney epithelial cell) treated with 100nM Phorbol-12-myristate-13-acetate for 1 hour whole cell lysates at 15 µg


Lane 3:
HEK-293 (Human embryonic kidney epithelial cell) treated with 100nM Phorbol-12-myristate-13-acetate for 1 hour whole cell lysates, then incubated with phosphatase at 15 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/20000 dilution

Predicted band size: 73 kDa

Observed band size: 73 kDa



Western blot - Anti-Raf1 antibody [EP4969] - N-terminal (ab181115)

Raf1 Western blot staining using rabbit Anti-Raf1 antibody

Lanes 1 and 2: Green signal from target – ab181115 observed at 74 kDa
Lanes 3 and 4: Red signal from loading control – ab8226 observed at 42 kDa
Lanes 5 and 6: Merged (red and green) signal
Unpurified ab181115 was shown to specifically react with Raf1 when Raf1 knockout samples were used. Wild-type and Raf1 knockout samples were subjected to SDS-PAGE. ab181115 and ab8226 (loading control to beta actin) were both diluted 1/1000 and incubated overnight at 4°C. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preadsorbed (ab216773) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preadsorbed (ab216776) secondary antibodies at 1/10000 dilution for 1 hour at room temperature before imaging.


Lanes 1 - 4:
Western blot - Anti-Raf1 antibody [EP4969] - N-terminal (ab181115) at 1/1000 dilution

Lanes 5 - 6:
Western blot - Anti-Raf1 antibody [EP4969] - N-terminal (ab181115)

Lanes 1, 3 and 5:
Wild-type HAP1 cell lysate at 20 µg

Lanes 2, 4 and 6:
Raf1 knockout HAP1 cell lysate at 20 µg

Predicted band size: 73 kDa



Western blot - Anti-Raf1 antibody [EP4969] - N-terminal (ab181115)

All lanes:
Western blot - Anti-Raf1 antibody [EP4969] - N-terminal (ab181115) at 1/1000 dilution

Lane 1:
HeLa (Human cervix adenocarcinoma epithelial cell) whole cell lysates at 20 µg


Lane 2:
K-562 (Human chronic myelogenous leukemia lymphoblast) whole cell lysates at 20 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/20000 dilution

Predicted band size: 73 kDa

Observed band size: 73 kDa



Western blot - Anti-Raf1 antibody [EP4969] - N-terminal (ab181115)

All lanes:
Western blot - Anti-Raf1 antibody [EP4969] - N-terminal (ab181115) at 1/1000 dilution


All lanes:
PC-12 (Rat adrenal gland pheochromocytoma) whole cell lysates at 20 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/20000 dilution

Predicted band size: 73 kDa

Observed band size: 73 kDa



Western blot - Anti-Raf1 antibody [EP4969] - N-terminal (ab181115)

All lanes:
Western blot - Anti-Raf1 antibody [EP4969] - N-terminal (ab181115) at 1/1000 dilution

Lane 1:
RAW 264.7(Mouse Abelson murine leukemia virus-induced tumor macrophage) whole cell lysates at 20 µg

Lane 2:
C2C12 (Mouse myoblasts myoblast) whole cell lysates at 20 µg

Lane 3:
NIH/3T3 (Mouse embryonic fibroblast) whole cell lysates at 20 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/20000 dilution

Predicted band size: 73 kDa

Observed band size: 73 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.