abcam

ab181213

Anti-GDI1 antibody [EPR13195]

Recombinant | RabMAb

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR13195
Purity	Tissue culture supernatant

Reactivity data

IP

Tested

Species	Human
Dilution info	1/30 - 1/50
Notes	-

Predicted

Species	Mouse, Rat

Dilution info	-
Notes	-

WB

Tested

Species	Human
Dilution info	1/1000 - 1/10000
Notes	-

Predicted

Species	Mouse, Rat
Dilution info	-
Notes	-

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	1/40
Notes	ab172730 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.

Predicted

Species	Mouse, Rat
Dilution info	-
Notes	-

Target data

See full target information GDI1

Function	Regulates the GDP/GTP exchange reaction of most Rab proteins by inhibiting the dissociation of GDP from them, and the subsequent binding of GTP to them.

Promotes the dissociation of GDP-bound Rab proteins from the membrane and inhibits their activation. Promotes the dissociation of RAB1A, RAB3A, RAB5A and RAB10 from membranes.

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.


3 product images



Flow Cytometry (Intracellular) - Anti-GDI1 antibody [EPR13195] (ab181213)

GDI1 Flow Cytometry (Intracellular) staining using rabbit Anti-GDI1 antibody

Intracellular flow cytometric analysis of HeLa cells (2% paraformaldehyde-fixed) labeling GDI1 with ab181213at 1/40 dilution (red), or a rabbit monoclonal IgG isotype control (green). Goat anti rabbit IgG (FITC) at 1/150 dilution was used as a secondary antibody.




Immunoprecipitation - Anti-GDI1 antibody [EPR13195] (ab181213)

Western blot analysis of immunoprecipitation pellet from Human fetal brain lysate (lane 1) or a negative control (lane 2) immunoprecipitated using ab181213 at 1/50 dilution.
Secondary: Anti-Rabbit IgG (HRP), specific to the non-reduced form of IgG at 1/1500 dilution.

All lanes:
Immunoprecipitation - Anti-GDI1 antibody [EPR13195] (ab181213)

Predicted band size: 51 kDa



Western blot - Anti-GDI1 antibody [EPR13195] (ab181213)

GDI1 Western blot staining using rabbit Anti-GDI1 antibody

All lanes:
Western blot - Anti-GDI1 antibody [EPR13195] (ab181213) at 1/10000 dilution

Lane 1:
U-87 MG (Human glioblastoma-astrocytoma epithelial cell line) cell lysate at 20 µg

Lane 2:
HeLa cell lysate at 20 µg

Lane 3:
SH-SY5Y cell lysate at 20 µg

Lane 4:
Human fetal brain tissue lysate at 20 µg

Secondary

All lanes:
Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/1000 dilution

Predicted band size: 51 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.