abcam

ab181225

Anti-E1 Ubiquitin Activating Enzyme 1/UBA1 antibody [EPR14204(B)]

Recombinant | RabMAb

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR14204(B)
Purification technique	Affinity purification Protein A
Concentration	0.095 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

WB

Tested

Species	Human
Dilution info	1/1000 - 1/10000
Notes	-

Species	Mouse
Dilution info	1/1000
Notes	-

Species	Rat
Dilution info	1/1000
Notes	-

ICC/IF

Tested

Species	Human
Dilution info	1/100 - 1/250
Notes	-

Expected

Species	Mouse, Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	1/10
Notes	ab172730 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.

Expected

Species	Mouse, Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

IHC-P

Tested

Species	Human
Dilution info	1/50 - 1/100
Notes	-

Expected

Species	Mouse, Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

Target data

See full target information UBA1

Function	Catalyzes the first step in ubiquitin conjugation to mark cellular proteins for degradation through the ubiquitin-proteasome system (PubMed:1447181, PubMed:1606621, PubMed:33108101). Activates ubiquitin by first adenylating its C-terminal glycine residue with ATP, and thereafter linking this residue to the side chain of a cysteine residue in E1, yielding a ubiquitin-E1 thioester and free AMP (PubMed:1447181). Essential for the formation of radiation-induced foci, timely DNA repair and for response to replication stress. Promotes the recruitment of TP53BP1 and BRCA1 at DNA damage sites (PubMed:22456334).

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	The E1 Ubiquitin Activating Enzyme 1 also known as UBA1 performs the initial step in the ubiquitination process. This enzyme activates ubiquitin a small protein that tags other proteins for degradation by the proteasome. UBA1 binds to ubiquitin in an ATP-dependent manner facilitating its transfer to E2 ubiquitin-conjugating enzymes. UBA1 has a molecular mass of approximately 117 kDa and shows expression in various tissues particularly in actively dividing cells.
Biological function summary	The enzyme plays an essential role in protein homeostasis by initiating the ubiquitin-proteasome system. UBA1 actively contributes to cellular regulation by targeting damaged or misfolded proteins for degradation thereby maintaining protein quality control. The enzyme is part of a complex alongside other proteins involved in the ubiquitination pathway which include E2 enzymes and E3 ubiquitin ligases that work together to achieve selective protein ubiquitination.
Pathways	E1 Ubiquitin Activating Enzyme 1 functions centrally in the ubiquitin-proteasome pathway a major mechanism governing protein turnover. It cooperates with other enzymes such as E2 and E3 to form a cascade that ensures substrate specificity and timely degradation. The enzyme also connects to the NF-kB signaling pathway where it regulates the stability of inhibitor kB proteins influencing immune responses and inflammation processes.
Associated diseases and disorders	Mutations or dysregulation of UBA1 contribute to spinal muscular atrophy (SMA) and X-linked spinal muscular atrophy (XL-SMA). These conditions involve motor neuron degeneration linked to the impaired ubiquitin-proteasome function. Additionally alterations in UBA1 activity can impact B-cell-activation through the NF-kB pathway possibly leading to autoimmune disorders. Understanding the interactions of UBA1 with associated proteins provides insight into potential therapeutic targets for treating these diseases.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended

We do not recommend this combination. It is not covered by our product promise.


We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

---


6 product images



Immunocytochemistry/ Immunofluorescence - Anti-E1 Ubiquitin Activating Enzyme 1/UBA1 antibody [EPR14204(B)] (ab181225)

E1 Ubiquitin Activating Enzyme 1/UBA1 Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-E1 Ubiquitin Activating Enzyme 1/UBA1 antibody

Immunofluorescent analysis of acetone-fixed HeLa cells labeling E1 Ubiquitin Activating Enzyme with ab181225 at 1/250 dilution followed by Goat anti rabbit IgG (Alexa Fluor® 555) secondary antibody at 1/200 dilution (red). Counter stained with Dapi (blue).




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-E1 Ubiquitin Activating Enzyme 1/UBA1 antibody [EPR14204(B)] (ab181225)

E1 Ubiquitin Activating Enzyme 1/UBA1 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-E1 Ubiquitin Activating Enzyme 1/UBA1 antibody

Immunohistochemical analysis of paraffin-embedded Human kidney tissue labeling E1 Ubiquitin Activating Enzyme with ab181225 at 1/100 dilution followed by prediluted HRP Polymer for Rabbit IgG. Counter stained with Hematoxylin.


Perform heat mediated antigen retrieval with EDTA buffer pH 9 before commencing with IHC staining protocol.



Flow Cytometry (Intracellular) - Anti-E1 Ubiquitin Activating Enzyme 1/UBA1 antibody [EPR14204(B)] (ab181225)

E1 Ubiquitin Activating Enzyme 1/UBA1 Flow Cytometry (Intracellular) staining using rabbit Anti-E1 Ubiquitin Activating Enzyme 1/UBA1 antibody

Intracellular flow cytometric analysis of2% paraformaldehyde-fixed K562 cells labeling E1 Ubiquitin Activating Enzyme with ab181225 at 1/10 dilution (red) compared to a Rabbit monoclonal IgG isotype control (green), followed byGoat anti rabbit IgG (FITC) secondary antibody at 1/150 dilution.



Western blot - Anti-E1 Ubiquitin Activating Enzyme 1/UBA1 antibody [EPR14204(B)] (ab181225)

E1 Ubiquitin Activating Enzyme 1/UBA1 Western blot staining using rabbit Anti-E1 Ubiquitin Activating Enzyme 1/UBA1 antibody

All lanes:
Western blot - Anti-E1 Ubiquitin Activating Enzyme 1/UBA1 antibody [EPR14204(B)] (ab181225) at 1/10000 dilution

Lane 1:
Human fetal brain tissue lysate at 20 µg

Lane 2:
K562 cell lysate at 20 µg

Lane 3:
HeLa cell lysate at 20 µg


Lane 4:
Jurkat cell lysate at 20 µg

Secondary

All lanes:
Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/1000 dilution

Predicted band size: 118 kDa

Observed band size: 118 kDa



Western blot - Anti-E1 Ubiquitin Activating Enzyme 1/UBA1 antibody [EPR14204(B)] (ab181225)

Diluting and blocking buffer: 5% NFDM/TBST 5% NFDM/TBST.

The lysates were freshly made and used for Western Blotting immediately to minimize protein degradation.

All lanes:

Western blot - Anti-E1 Ubiquitin Activating Enzyme 1/UBA1 antibody [EPR14204(B)] (ab181225) at 1/1000 dilution

Lane 1:
HeLa (human cervical adenocarcinoma epithelial cell) whole cell lysate at 20 µg

Lane 2:
293T (human embryonic kidney epithelial cell) whole cell lysate at 20 µg

Lane 3:
NIH/3T3 (mouse embryonic fibroblast) whole cell lysate at 20 µg

Lane 4:
PC-12 (rat adrenal gland pheochromocytoma cell) whole cell lysate at 20 µg


Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/20000 dilution

Predicted band size: 118 kDa, 117 kDa

Observed band size: 117 kDa, 118 kDa

Exposure time: 8s



Western blot - Anti-E1 Ubiquitin Activating Enzyme 1/UBA1 antibody [EPR14204(B)] (ab181225)

Blocking and diluting buffer: 5% NFDM/TBST
All lanes:
Western blot - Anti-E1 Ubiquitin Activating Enzyme 1/UBA1 antibody [EPR14204(B)] (ab181225) at 1/1000 dilution

Lane 1:
GST tagged Recombinant Human UBA7(Ube1L) protein (Full length, 166 KDa)

Lane 2:
GST tagged Recombinant Human UBA1 protein (Full length, 146 KDa)

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051)

Predicted band size: 118 kDa

Observed band size: 146 kDa, 166 kDa

Exposure time: 180s

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.