abcam

ab181247

Anti-IER3IP1 antibody [EPR13440]

Recombinant | RabMAb

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR13440
Purification technique	Affinity purification Protein A
Concentration	1.085 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

IP

Tested

Species	Human
Dilution info	1/50 - 1/70
Notes	-

Predicted

Species	Mouse, Rat
Dilution info	-
Notes	-

WB

Tested

Species	Human
Dilution info	1/1000
Notes	-

Predicted

Species	Mouse, Rat
Dilution info	-
Notes	-

ICC/IF

Tested

Species	Human
Dilution info	1/100 - 1/250
Notes	-

Predicted

Species	Mouse, Rat
Dilution info	-
Notes	-

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	1/210
Notes	ab172730 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.

Predicted

Species	Mouse, Rat
Dilution info	-
Notes	-

IHC-P

Tested

Species	Human
Dilution info	1/100
Notes	-

Predicted

Species	Mouse, Rat
Dilution info	-
Notes	-

Target data

See full target information IER3IP1

Function	Regulator of endoplasmic reticulum secretion that acts as a key determinant of brain size (PubMed:33122427). Required for secretion of extracellular matrix proteins (PubMed:33122427). Required for correct brain development by depositing sufficient extracellular matrix proteins for tissue integrity and the proliferation of neural progenitors (PubMed:33122427). Acts as a regulator of the unfolded protein response (UPR) (By similarity).

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	IER3IP1 also known as Immediate Early Response 3 Interacting Protein 1 is a small protein with a mass of approximately 11 kDa. It primarily localizes within the endoplasmic reticulum (ER) and shows expression in various tissues including the brain liver and pancreas. IER3IP1 plays a mechanical role in protein trafficking and the regulation of ER stress affecting cellular responses to environmental changes.
Biological function summary	IER3IP1 participates in processes related to cellular stress response and protein folding. It acts as a component involved in maintaining ER function possibly engaging in a protein complex to modulate its activity. This modulation is essential for controlling apoptosis and cell survival suggesting its involvement in critical cellular processes. The protein's interactions help sustain proper protein synthesis and folding within the ER.

Pathways	IER3IP1 interacts within the unfolded protein response pathway maintaining ER homeostasis. It engages with proteins like GRP78 and ATF6 which are significant in managing ER stress and protein folding pathways. These interactions demonstrate IER3IP1's importance in cellular mechanisms that protect cells from stress-induced damages.
Associated diseases and disorders	Mutations or misregulation of IER3IP1 correlate with microcephaly and epilepsy. Research indicates its connection with proteins involved in neurodevelopmental disorders possibly interacting with molecules like Reelin or Dystrophin. This relationship highlights the significance of IER3IP1 in maintaining neural cell health and suggests its potential as a target for therapeutic strategies.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.


6 product images



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-IER3IP1 antibody [EPR13440] (ab181247)

IER3IP1 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-IER3IP1 antibody


Immunohistochemical analysis of paraffin-embedded Human cardiac muscle tissue labeling IER3IP1 with ab181247 at 1/100 dilution followed by pre-diluted HRP-conjugated secondary antibody and counter-stained with Hematoxylin.
Perform heat mediated antigen retrieval with EDTA buffer pH 9 before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-IER3IP1 antibody [EPR13440] (ab181247)

IER3IP1 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-IER3IP1 antibody


Immunohistochemical analysis of paraffin-embedded Human kidney tissue labeling IER3IP1 with ab181247 at 1/100 dilution followed by pre-diluted HRP-conjugated secondary antibody and counter-stained with Hematoxylin.
Perform heat mediated antigen retrieval with EDTA buffer pH 9 before commencing with IHC staining protocol.



Immunocytochemistry/ Immunofluorescence - Anti-IER3IP1 antibody [EPR13440] (ab181247)

IER3IP1 Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-IER3IP1 antibody


Immunofluorescent analysis of HeLa cells (-20°C Acetone-fixed) labeling IER3IP1 with ab181247 at 1/250 dilution dollowed by Goat anti rabbit IgG (Alexa Fluor®555) at 1/200 dilution and DAPI staining (blue).



Flow Cytometry (Intracellular) - Anti-IER3IP1 antibody [EPR13440] (ab181247)

IER3IP1 Flow Cytometry (Intracellular) staining using rabbit Anti-IER3IP1 antibody


Intracellular flow cytometricl analysis of HeLa cells (paraformaldehyde-fixed, 2%)labeling IER3IP1 with ab181247 at 1/210 (red) compared to a negative control antibody (green),followed by Goat anti rabbit IgG (FITC) secondary at 1/150 dilution.



Immunoprecipitation - Anti-IER3IP1 antibody [EPR13440] (ab181247)

Western blot analysis of immunoprecipitation pellet from HeLa cell lysate immunoprecipitated using ab181247 at 1/50 dilution.
Secondary: Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugate at 1/1000.
All lanes:
Immunoprecipitation - Anti-IER3IP1 antibody [EPR13440] (ab181247)

Predicted band size: 9 kDa



Western blot - Anti-IER3IP1 antibody [EPR13440] (ab181247)

All lanes:
Western blot - Anti-IER3IP1 antibody [EPR13440] (ab181247) at 1/1000 dilution

Lane 1:
HepG2 lysate at 20 µg

Lane 2:
HeLa lysate at 20 µg

Lane 3:
Human fetal heart lysate at 20 µg

Secondary

All lanes:
Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/1000 dilution

Predicted band size: 9 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.