abcam

ab181859

Anti-DDAH1 antibody [EPR13921]

Recombinant | RabMAb | KO Validated

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR13921
Purification technique	Affinity purification Protein A
Concentration	0.163 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

IP

Tested

Species	Human
Dilution info	1/40
Notes	-

WB

Tested

Species	Human
Dilution info	1/1000 - 1/10000
Notes	-

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	1/30
Notes	ab172730 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.

IHC-P

Tested

Species	Human
Dilution info	1/50
Notes	Perform heat-mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Target data

See full target information DDAH1

Function	Hydrolyzes N(G),N(G)-dimethyl-L-arginine (ADMA) and N(G)-monomethyl-L-arginine (MMA) which act as inhibitors of NOS. Has therefore a role in the regulation of nitric oxide generation.

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks

Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	The DDAH1 protein also known as dimethylarginine dimethylaminohydrolase 1 plays an important role in regulating nitric oxide (NO) production by metabolizing asymmetric dimethylarginine (ADMA) an inhibitor of nitric oxide synthase. This enzyme has a molecular mass of approximately 31 kDa and is widely expressed in various tissues including liver kidney and brain. It impacts the cardiovascular system by controlling ADMA levels which influence endothelial function and vascular tone.
Biological function summary	The actions of DDAH1 contribute significantly to cardiovascular homeostasis. Increased levels of ADMA resulting from impaired DDAH1 activity can lead to decreased NO production impacting cardiovascular health. DDAH1 does not form part of a complex but acts individually to affect NO synthesis through ADMA degradation. Understanding the biological roles of DDAH1 helps to appreciate how NO-mediated processes are tightly regulated within the body and its impact on vascular health.
Pathways	DDAH1 participates actively in the nitric oxide synthesis pathway. It modulates the levels of endothelial nitric oxide synthase (eNOS) by regulating ADMA which serves as an eNOS inhibitor. Through this pathway DDAH1 ensures a balance between NO production and regulation supporting vascular homeostasis and blood pressure

	control. DDAH1 shares functional connections with proteins like eNOS which are central to maintaining vascular health and preventing endothelial dysfunction.
Associated diseases and disorders	Dysfunction of DDAH1 links closely to cardiovascular diseases and conditions like atherosclerosis. Elevated ADMA levels resulting from compromised DDAH1 activity contribute to endothelial dysfunction a precursor to atherosclerosis. The interconnected role with eNOS highlights DDAH1's importance in managing ADMA concentrations as imbalances can exacerbate disease progression. Investigating the influence of DDAH1 on these diseases offers insights into potential therapeutic targets for cardiovascular pathologies.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.


5 product images



Flow Cytometry (Intracellular) - Anti-DDAH1 antibody [EPR13921] (ab181859)

DDAH1 Flow Cytometry (Intracellular) staining using rabbit Anti-DDAH1 antibody


Intracellular flow cytometric analysis of 2% paraformaldehyde-fixed 293 cells labeling DDAH1 with ab181859 at 1/30 dilution (red), compared to a Rabbit monoclonal IgGisotype control (green), followed by Goat anti-Rabbit IgG (FITC) secondary antibody at 1/150 dilution.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-DDAH1 antibody [EPR13921] (ab181859)

DDAH1 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-DDAH1 antibody


Immunohistochemical analysis of paraffin-embedded Human hepatocellular carcinoma tissue labeling DDAH1 with ab181859 at 1/50 dilution, followed by prediluted HRP Polymer for Rabbit IgG. Counter stained with Hematoxylin.
Perform heat mediated antigen retrieval with EDTA buffer pH 9 before commencing with IHC staining protocol.



Immunoprecipitation - Anti-DDAH1 antibody [EPR13921] (ab181859)

Western blot analysis of 293 cell lysate immunoprecipitated using ab181859 at 1/40 dilution. Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugate secondary antibody was used at 1/1000 dilution.
All lanes:
Immunoprecipitation - Anti-DDAH1 antibody [EPR13921] (ab181859)

Predicted band size: 31 kDa



Western blot - Anti-DDAH1 antibody [EPR13921] (ab181859)

DDAH1 Western blot staining using rabbit Anti-DDAH1 antibody

Lanes 1 - 4: Merged signal (red and green). Green - ab181859 observed at 37 kDa. Red - loading control, ab7291 (Mouse anti-Alpha Tubulin [DM1A] observed at 55kDa.
ab181859 was shown to react with DDAH1 in wild-type A431 cells in Western blot. Loss of signal was observed when DDAH1 knockout sample was used. Wild-type A431 and DDAH1 knockout cell lysates were subjected to SDS-PAGE. Membranes were blocked in 3% Milk in TBS-T (0.1% Tween®) before incubation with ab181859 and ab7291 (Mouse anti-Alpha Tubulin [DM1A] overnight at 4°C at a 1 in 10000 dilution and a 1 in 20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preabsorbed (ab216773) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preabsorbed (ab216776) secondary antibodies at 1 in 20000 dilution for 1 hour at room temperature before imaging.

All lanes:
Western blot - Anti-DDAH1 antibody [EPR13921] (ab181859) at 1/10000 dilution

Lane 1:
Wild-type A431 cell lysate at 40 µg

Lane 2:
DDAH1 knockout A431 cell lysate at 40 µg

Lane 2:
Western blot - Human DDAH1 knockout A-431 cell line (ab261909)


Lane 3:
HUVEC (Human umbilical vein endothelial cell line) whole cell lysate at 40 µg

Lane 4:
Hep G2 (Human liver hepatocellular carcinoma cell line) whole cell lysate at 40 µg

Performed under reducing conditions.

Predicted band size: 31 kDa

Observed band size: 37 kDa



Western blot - Anti-DDAH1 antibody [EPR13921] (ab181859)

All lanes:
Western blot - Anti-DDAH1 antibody [EPR13921] (ab181859) at 1/5000 dilution

Lane 1:
293 cell lysate at 10 µg

Lane 2:
HepG2 cell lysate at 10 µg

Secondary

All lanes:
Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugate at 1/1000 dilution

Predicted band size: 31 kDa

Observed band size: 37 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.