abcam

ab183330

# Anti-Cytokeratin 17 antibody [SP95] - C-terminal

Recombinant | RabMAb

## Key facts

| | |
|---|---|
| **Isotype** | IgG |
| **Host species** | Rabbit |
| **Storage buffer** | pH: 7.6 Preservative: 0.1% Sodium azide Constituents: PBS, 1% BSA |
| **Form** | Liquid |
| **Clonality** | Monoclonal |
| **Immunogen** | The exact immunogen used to generate this antibody is proprietary information. |
| **Clone number** | SP95 |
| **Purification technique** | Affinity purification Protein A/G |
| **Concentration** | 0.015 mg/mL The concentration of this product may be batch-dependent<br>Batch concentration finder → |
| **Purification notes** | Purified from TCS by protein A/G. |

## Reactivity data

## IHC-P

### Tested

| | |
|---|---|
| **Species** | Human |
| **Dilution info** | 1/100 |
| **Notes** | - |

### Predicted

| | |
|---|---|
| **Species** | Mouse, Rat, Cow, Dog |
| **Dilution info** | - |
| **Notes** | - |

## Flow Cyt (Intra)

### Tested

| | |
|---|---|
| **Species** | Human |
| **Dilution info** | 1/20 - 1/100 |
| **Notes** | ab172730 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody. |

### Predicted

| | |
|---|---|
| **Species** | Mouse, Rat, Cow, Dog |
| **Dilution info** | - |
| **Notes** | - |

## WB

### Tested

| | |
|---|---|
| **Species** | Human |
| **Dilution info** | 1/100 |
| **Notes** | - |

### Predicted

| | |
|---|---|
| **Species** | Mouse, Rat, Cow, Dog |
| **Dilution info** | - |
| **Notes** | - |

## ICC/IF

### Tested

| | |
|---|---|
| **Species** | Human |
| **Dilution info** | 1/100 |
| **Notes** | - |

### Predicted

| | |
|---|---|
| **Species** | Mouse, Rat, Cow, Dog |
| **Dilution info** | - |
| **Notes** | - |

## Target data

See full target information KRT17

**Function**

Type I keratin involved in the formation and maintenance of various skin appendages, specifically in determining shape and orientation of hair (By similarity). Required for the correct growth of hair follicles, in particular for the persistence of the anagen (growth) state (By similarity). Modulates the function of TNF in the specific context of hair cycling. Regulates protein synthesis and epithelial cell growth through binding to the adapter protein SFN and by stimulating Akt/mTOR pathway (By similarity). Involved in tissue repair. May be a marker of basal cell differentiation in complex epithelia and therefore indicative of a certain type of epithelial 'stem cells'. Acts as a promoter of epithelial proliferation by acting a regulator of immune response in skin: promotes Th1/Th17-dominated immune environment contributing to the development of basaloid skin tumors (By similarity). May act as an autoantigen in the immunopathogenesis of psoriasis, with certain peptide regions being a major target for autoreactive T-cells and hence causing their proliferation.

## Storage

| | |
|---|---|
| **Shipped at conditions** | Conditional Ambient |
| **Appropriate short-term storage duration** | 1-2 weeks |
| **Appropriate short-term storage conditions** | +4°C |
| **Appropriate long-term storage conditions** | -20°C |

| | |
|---|---|
| **Aliquoting information** | Upon delivery aliquot |
| **Storage information** | Avoid freeze / thaw cycle |

## Notes

**What are the advantages of a recombinant monoclonal antibody?**
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

## Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

| | |
|---|---|
| **Activity summary** | Cytokeratin 17 often called K17 or keratin 17 is a type I intermediate filament protein with a molecular weight of approximately 46 kDa. It forms part of the cytoskeleton in epithelial cells and is notably present in the basal cells of epithelial tissues including the skin hair follicles and the lining of internal organs. Due to its specific expression pattern K17 serves as a useful biomarker in various research and clinical settings. |
| **Biological function summary** | The roles of K17 extend beyond mere structural support. This protein contributes to cellular processes such as proliferation tissue repair and the regulation of immune responses. K17 often forms a complex with its partner keratin 6 enabling structural stability and resilience in epithelial cells. Its presence influences the behavior of the cells in skin and other epithelial-lined organs. |
| **Pathways** | K17 participates in key pathways related to skin biology and inflammation notably the Wnt/β-catenin and NF-κB signaling pathways. These pathways connect K17 to cellular processes like growth differentiation and inflammation. In particular keratin 6 is another important protein in these cascades closely interacting with K17 in maintenance and repair processes within epithelial tissues. |
| **Associated diseases and disorders** | K17 has been implicated in psoriasis and certain types of cancer such as cervical and breast cancer. In psoriasis an inflammatory skin condition K17 interacts with pro-inflammatory cytokines contributing to the hyperproliferative state of the skin. Its overexpression serves as a marker in diagnosing and understanding the severity of these disorders highlighting its relationship with keratin 6 which is similarly upregulated in psoriatic lesions and certain tumors. These associations make K17 a significant focus for therapeutic targeting and diagnostic applications. |

## Product promise

**Tested**

We have tested this species and application combination and it works. It is covered by our product promise.

**Expected**

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

**Predicted**

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

**Not recommended**

We do not recommend this combination. It is not covered by our product promise.

---

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.


In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

---

## 17 product images

---




### Flow Cytometry (Intracellular) - Anti-Cytokeratin 17 antibody [SP95] - C-terminal (ab183330)

Cytokeratin 17 Flow Cytometry (Intracellular) staining using rabbit Anti-Cytokeratin 17 antibody


Flow cytometry overlay histogram showing left HeLa positive cells and right negative MCF7 stained with ab183330 (red line). The cells were fixed with 80% methanol (5 min) and then permeabilised with 0.1% PBS-Triton X-100 for 15 min. The cells were then incubated in 1x PBS containing 10% normal goat serum to block non-specific protein-protein interaction followed by the antibody (ab183330) (1x 106 in 100µl at 0.008µg/ml (1/262500)) for 30min at 22°C.

The secondary antibody Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed was incubated at 1/4000 for 30min at 22°C

Isotype control antibody (black line) was Recombinant Rabbit IgG, monoclonal [EPR25A] - Isotype Control used at the same concentration and conditions as the primary antibody. Unlabelled sample (blue line) was also used as a control.

Acquisition of >5000 events were collected using a 50 mW Blue laser (488nm) and 525/40 bandpass filter.


---



## Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Cytokeratin 17 antibody [SP95] - C-terminal (ab183330)

Cytokeratin 17 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-Cytokeratin 17 antibody


Immunohistochemical analysis of paraffin embedded Human bladder transitional cell carcinoma tissue labeling Cytokeratin 17 with ab183330 at 1/100.




## Immunocytochemistry/ Immunofluorescence - Anti-Cytokeratin 17 antibody [SP95] - C-terminal (ab183330)

Cytokeratin 17 Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-Cytokeratin 17 antibody


Immunocytochemistry/ Immunofluorescence analysis of HeLa (human cervix adenocarcinoma epithelial cell) cells labeling Cytokeratin 17 with purified ab183330 at 1/100 (1.6µg/ml). Cells were fixed in 4% paraformaldehyde and permeabilized with 0.1% Triton X-100. Cells were counterstained with ab195889 Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor® 594) 1/200 (2.5 µg/ml). Goat anti rabbit IgG (Alexa Fluor® 488, ab150077) was used as the secondary antibody at 1/1000 (2 µg/ml) dilution. DAPI was used as nuclear counterstain. PBS instead of the primary antibody was used as the secondary antibody only control.



## Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Cytokeratin 17 antibody [SP95] - C-terminal (ab183330)

Cytokeratin 17 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-Cytokeratin 17 antibody


Immunohistochemical analysis of paraffin embedded Human skin tissue labeling Cytokeratin 17 with ab183330 at 1/100.



## Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Cytokeratin 17 antibody [SP95] - C-terminal (ab183330)

Cytokeratin 17 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-Cytokeratin 17 antibody


Immunohistochemical analysis of paraffin embedded Human cervix adenocarcinoma tissue labeling Cytokeratin 17 with ab183330 at 1/100.



## Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Cytokeratin 17 antibody [SP95] - C-terminal (ab183330)

Cytokeratin 17 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-Cytokeratin 17 antibody


Immunohistochemical analysis of paraffin embedded Human tonsil tissue labeling Cytokeratin 17 with ab183330 at 1/100.



## Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Cytokeratin 17 antibody [SP95] - C-terminal (ab183330)

Cytokeratin 17 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-Cytokeratin 17 antibody


Immunohistochemical analysis of paraffin embedded Human prostate tissue labeling Cytokeratin 17 with ab183330 at 1/100.




## Flow Cytometry (Intracellular) - Anti-Cytokeratin 17 antibody [SP95] - C-terminal (ab183330)

Cytokeratin 17 Flow Cytometry (Intracellular) staining using rabbit Anti-Cytokeratin 17 antibody


Flow cytometry analysis of Hela (human cervix adenocarcinoma) labeling Cytokeratin 17 with purified ab183330 at 1/20 dilution (7.85 µg/ml) (red). Cells were fixed with 4% paraformaldehyde and permeabilised with 90% methanol. Goat anti rabbit IgG (Alexa Fluor® 488, ab150077) at 1/2000 dilution was used as a secondary antibody. Isotype control - Rabbit monoclonal IgG (ab172730) (Black). Unlableled control - Unlabelled cells (blue).



## Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Cytokeratin 17 antibody [SP95] - C-terminal (ab183330)

Cytokeratin 17 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-Cytokeratin 17 antibody


Immunohistochemical analysis of paraffin embedded Human thymus tissue labeling Cytokeratin 17 with ab183330 at 1/100.



## Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Cytokeratin 17 antibody [SP95] - C-terminal (ab183330)

Cytokeratin 17 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-Cytokeratin 17 antibody


Immunohistochemical analysis of paraffin embedded Human skin squamous cell carcinoma tissue labeling Cytokeratin 17 with ab183330 at 1/100.




## Flow Cytometry (Intracellular) - Anti-Cytokeratin 17 antibody [SP95] - C-terminal (ab183330)

Cytokeratin 17 Flow Cytometry (Intracellular) staining using rabbit Anti-Cytokeratin 17 antibody


Flow cytometric analysis of Cytokeratin HeLa cells using ab183330 (green) compare to negative control of rabbit IgG (blue).



## Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Cytokeratin 17 antibody [SP95] - C-terminal (ab183330)

Cytokeratin 17 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-Cytokeratin 17 antibody


Immunohistochemical analysis of paraffin embedded Human bladder tissue labeling Cytokeratin 17 with ab183330 at 1/100.



## Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Cytokeratin 17 antibody [SP95] - C-terminal (ab183330)

Cytokeratin 17 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-Cytokeratin 17 antibody


Immunohistochemical analysis of paraffin embedded Human prostate adenocarcinoma tissue labeling Cytokeratin 17 with ab183330 at 1/100.



## Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Cytokeratin 17 antibody [SP95] - C-terminal (ab183330)

Cytokeratin 17 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-Cytokeratin 17 antibody


Immunohistochemical analysis of paraffin embedded Human prostate carcinoma tissue labeling Cytokeratin 17 with ab183330 at 1/100.



## Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Cytokeratin 17 antibody [SP95] - C-terminal (ab183330)

Cytokeratin 17 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-Cytokeratin 17 antibody


Immunohistochemical analysis of paraffin embedded Human cervical adenocarcinoma tissue labeling Cytokeratin 17 with ab183330 at 1/100.



## Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Cytokeratin 17 antibody [SP95] - C-terminal (ab183330)

Cytokeratin 17 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-Cytokeratin 17 antibody

Immunohistochemical analysis of paraffin embedded Human cervix tissue labeling Cytokeratin 17 with ab183330 at 1/100.



## Western blot - Anti-Cytokeratin 17 antibody [SP95] - C-terminal (ab183330)

All lanes:
Western blot - Anti-Cytokeratin 17 antibody [SP95] - C-terminal (ab183330) at 1/25 dilution

All lanes:
HeLa cell lysate

Predicted band size: 48 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.