abcam

ab184568

Anti-LSM5 antibody [EPR10690]

Recombinant | RabMAb

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR10690
Purification technique	Affinity purification Protein A
Concentration	0.628 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

WB

Tested

Species	Human
Dilution info	1/1000 - 1/10000
Notes	-

Predicted

Species	Mouse, Rat
Dilution info	-
Notes	-

ICC/IF

Tested

Species	Human
Dilution info	1/500
Notes	-

Predicted

Species	Mouse, Rat
Dilution info	-
Notes	-

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	1/110
Notes	ab172730 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.

Predicted

Species	Mouse, Rat
Dilution info	-
Notes	-

IHC-P

Tested

Species	Human
Dilution info	1/50 - 1/100
Notes	-

Predicted

Species	Mouse, Rat
Dilution info	-
Notes	-

Target data

See full target information LSM5

Function	Plays a role in pre-mRNA splicing as component of the U4/U6-U5 tri-snRNP complex that is involved in spliceosome assembly, and as component of the precatalytic spliceosome (spliceosome B complex) (PubMed:28781166). The heptameric LSM2-8 complex binds specifically to the 3'-terminal U-tract of U6 snRNA (PubMed:10523320).

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	LSM5 also known as U6 snRNA-associated Sm-like protein LSm5 plays an important role in the mechanics of RNA processing. This protein has a mass of approximately 9.7 kDa. LSM5 is expressed in various tissues with notable concentrations found in the cytoplasm and nucleus. It functions as part of the larger Lsm protein family which is known for playing roles in mRNA degradation and splicing.
Biological function summary	LSM5 participates in the formation and stabilization of the U6 snRNA. As a member of the Lsm complex it supports RNA processing and stability. The Lsm complex typically consists of seven Lsm proteins including LSM5 forming a heptameric ring structure. This complex assists in the splicing of pre-mRNA a critical step in the regulation of gene expression. By binding to the 3’ end of U6 snRNA LSM5 facilitates the splicing mechanism necessary for proper mRNA maturation.
Pathways	LSM5 is actively involved in the RNA splicing pathway an important step for gene expression and mRNA maturation. One notable pathway it participates in is the spliceosome cycle closely associated with the processing of pre-mRNA into mature mRNA. In these pathways LSM5 interacts with other components such as U6 snRNP and helps in regulating spliceosomal dynamics. This highlights LSM5's role in maintaining the fidelity and efficiency of mRNA splicing.
Associated diseases and disorders	LSM5 links to certain conditions stemming from defective RNA processing. One example is spinal muscular atrophy a disorder that affects the motor neurons due to impaired splicing. LSM5 also associates with amyotrophic lateral sclerosis (ALS) which involves RNA metabolism disruption. In these diseases LSM5's interaction with proteins like SMN (survival of motor neuron) and TDP-43 can disrupt normal cellular functions and lead to disease progression.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended

We do not recommend this combination. It is not covered by our product promise.


We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

---


5 product images



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-LSM5 antibody [EPR10690] (ab184568)

LSM5 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-LSM5 antibody


Immunohistochemical analysis of paraffin embedded Human stomach tissue labeling LSM5 with ab184568 a 1/100 dilution. Counterstained with Hematoxylin.
Perform heat mediated antigen retrieval with EDTA buffer pH 9 before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-LSM5 antibody [EPR10690] (ab184568)

LSM5 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-LSM5 antibody


Immunohistochemical analysis of paraffin embedded Human astrocytoma tissue labeling LSM5 with ab184568 at a 1/100 dilution. Counterstained with Hematoxylin.
Perform heat mediated antigen retrieval with EDTA buffer pH 9 before commencing with IHC staining protocol.



Flow Cytometry (Intracellular) - Anti-LSM5 antibody [EPR10690] (ab184568)

LSM5 Flow Cytometry (Intracellular) staining using rabbit Anti-LSM5 antibody


Intracellular Flow Cytometry analysis of T47-Dcells using ab184568at a 1/110 dilution (red) or aRabbit monoclonal IgG (negative) (green). Goat anti rabbit IgG (FITC) secondary usedat a 1/150 dilution.



Immunocytochemistry/ Immunofluorescence - Anti-LSM5 antibody [EPR10690] (ab184568)

LSM5 Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-LSM5 antibody

Immunofluorescence analysis of 4% paraformaldehyde fixed T47-D cells labeling LSM5 with ab184568 at a 1/500 dilution. Goat anti rabbit IgG (Alexa Fluor®555) secondary used at a 1/200 diution. Conterstained with Dapi (right image).



Western blot - Anti-LSM5 antibody [EPR10690] (ab184568)

All lanes:
Western blot - Anti-LSM5 antibody [EPR10690] (ab184568) at 1/10000 dilution

All lanes:
T47-D lysate at 10 µg

Secondary

All lanes:
Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/1000 dilution

Predicted band size: 10 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.