abcam

ab186750

Anti-MBIP antibody [EPR13951]

Recombinant | RabMAb

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR13951
Purification technique	Affinity purification Protein A
Concentration	1.921 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

WB

Tested

Species	Human
Dilution info	1/1000 - 1/10000
Notes	-

Predicted

Species	Mouse, Rat
Dilution info	-
Notes	-

ICC/IF

Tested

Species	Human
Dilution info	1/500
Notes	-

Predicted

Species	Mouse, Rat
Dilution info	-
Notes	-

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	1/400
Notes	ab172730 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.

Predicted

Species	Mouse, Rat
Dilution info	-
Notes	-

IHC-P

Tested

Species	Human
Dilution info	1/100 - 1/250
Notes	-

Predicted

Species	Mouse, Rat
Dilution info	-
Notes	-

Target data

See full target information MBIP

Function	Functions as part of the ATAC complex, a complex with histone acetyltransferase activity on histones H3 and H4 (PubMed:19103755). Inhibits the MAP3K12 activity to induce the activation of the JNK/SAPK pathway (PubMed:10801814). Together with MOCS2B, inhibits the activity of stress kinase EIF2AK2/PKR; this may lead to effects such as suppression of JNK activation and subsequent stress-responsive transcription, or suppression of eIF2a phosphorylation to favor translation (PubMed:26705305).

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	MBIP also known as Mps-One Binder kinase activator-like protein 1 acts mechanically by being a component of certain cellular complexes involved in intracellular signaling. It has a molecular mass of approximately 25 kDa. Expression of MBIP occurs in many tissues but it is found at higher levels in brain and kidney tissues suggesting its significant role in these areas.
Biological function summary	MBIP regulates signaling pathways that control cell growth development and differentiation. As a part of the c-Jun N-terminal kinase (JNK) signalosome complex MBIP influences pathways that mediate responses to stress and inflammatory stimuli. The presence of MBIP in these complexes emphasizes its role in modulating the cellular response to environmental changes shedding light on how it contributes to maintaining cellular homeostasis.
Pathways	MBIP functions within the JNK signaling and the domino effect on MAP kinase pathways. In the JNK pathway MBIP interacts closely with proteins such as MKK4 and JNKs to ensure proper signal transduction and cellular outcomes. Its activity helps mediate cellular responses modulating processes like apoptosis and cell survival indicating its importance in maintaining cellular balance and health.
Associated diseases and disorders	MBIP shows relevance to conditions like cancer and neurodegenerative diseases. Aberrant MBIP expression or activity can disrupt the JNK pathway leading to uncontrolled cell proliferation in cancer or altered neuronal functioning in conditions like Alzheimer's disease. Connections between MBIP and other proteins such as JNKs and MKK4 highlight its influence on pathological developments in these disorders providing a critical link between cellular signaling and disease progression.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.


6 product images



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-MBIP antibody [EPR13951] (ab186750)

MBIP Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-MBIP antibody


Immunohistochemical analysis of paraffin-embedded Human clear cell carcinoma of kidney tissue labeling MBIP with ab186750 at 1/250 dilution. Secondary ab: ready to use HRP Polymer for Rabbit IgG. Counter stain: Hematoxylin.
Perform heat mediated antigen retrieval with EDTA buffer pH 9 before commencing with IHC staining protocol.



Immunocytochemistry/ Immunofluorescence - Anti-MBIP antibody [EPR13951] (ab186750)

MBIP Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-MBIP antibody


Immunofluorescence analysis of HeLa cells labeling MBIP with ab186750 at 1/500 dilution. Secondary antibody: Goat anti rabbit IgG (Alexa Fluor® 488). Fixative: 4% paraformaldehyde. Counter stain: Dapi (blue).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-MBIP antibody [EPR13951] (ab186750)

MBIP Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-MBIP antibody


Immunohistochemical analysis of paraffin-embedded Human cardiac muscle tissue labeling MBIP with ab186750 at 1/250 dilution. Secondary antibody: Ready to use HRP Polymer for Rabbit IgG. Counter stain: Hematoxylin.
Perform heat mediated antigen retrieval with EDTA buffer pH 9 before commencing with IHC staining protocol.



Flow Cytometry (Intracellular) - Anti-MBIP antibody [EPR13951] (ab186750)

MBIP Flow Cytometry (Intracellular) staining using rabbit Anti-MBIP antibody

Intracellular flow cytometric analysis of HeLa cell labeling MBIP usingab186750 at 1/400 dilution. Secondary antibody: Goat anti rabbit IgG (FITC) at 1/400 dilution. Fixative: 2% paraformaldehyde. Isotype control: Rabbit monoclonal IgG.



Western blot - Anti-MBIP antibody [EPR13951] (ab186750)

MBIP Western blot staining using rabbit Anti-MBIP antibody

Blocking buffer: 5% NFDM/TBST
All lanes:
Western blot - Anti-MBIP antibody [EPR13951] (ab186750) at 1/10000 dilution


Lane 1:
Hela cell lysate at 20 µg

Lane 2:
293 cell lysate at 20 µg

Secondary

All lanes:
Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/1000 dilution

Predicted band size: 39 kDa



Western blot - Anti-MBIP antibody [EPR13951] (ab186750)

MBIP Western blot staining using rabbit Anti-MBIP antibody

Blocking buffer: 5% NFDM/TBST
All lanes:
Western blot - Anti-MBIP antibody [EPR13951] (ab186750) at 1/10000 dilution

Lane 1:
Human fetal lung tissue lysate at 10 µg

Lane 2:
Human fetal heart tissue lysate at 10 µg

Secondary

All lanes:
Anti-Rabbit IgG (HRP), specific to the non-reduced form of IgG at 1/1000 dilution

Predicted band size: 39 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.