abcam

ab187075

Anti-Thrombomodulin antibody [EPR18217]

Recombinant | RabMAb | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR18217
Purification technique	Affinity purification Protein A
Concentration	0.555 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

Flow Cyt

Tested

Species	Mouse
Dilution info	1/60
Notes	-

Expected

Species	Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

WB

Tested

Species	Mouse
Dilution info	1/1000
Notes	-

Species	Rat
Dilution info	1/1000
Notes	-

IHC-P

Tested

Species	Mouse
Dilution info	1/2000
Notes	-

Species	Rat
Dilution info	1/2000
Notes	-

Target data

See full target information Thbd

Function Endothelial cell receptor that plays a critical role in regulating several physiological processes including hemostasis, coagulation, fibrinolysis, inflammation, and angiogenesis. Acts as a cofactor for thrombin activation of protein C/PROC on the surface of vascular endothelial cells leading to initiation of the activated protein C anticoagulant pathway. Also accelerates the activation of the plasma carboxypeptidase B2/CPB2, which catalyzes removal of C-terminal basic amino

acids from its substrates including kinins or anaphylatoxins leading to fibrinolysis inhibition (By similarity). Plays critical protective roles in changing the cleavage specificity of protease-activated receptor 1/PAR1, inhibiting endothelial cell permeability and inflammation (PubMed:38174561). Suppresses inflammation distinctly from its anticoagulant cofactor activity by sequestering HMGB1 thereby preventing it from engaging cellular receptors such as RAGE and contributing to the inflammatory response (By similarity).

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	Thrombomodulin also known as CD141 is a glycoprotein with a mass of approximately 74 kDa. It is an integral membrane protein expressed on the surface of endothelial cells. Thrombomodulin acts mechanically by binding

	thrombin an enzyme involved in blood coagulation to modulate its activity. This binding substantially alters thrombin's substrate specificity turning it into an anticoagulant enzyme that activates protein C instead of a procoagulant enzyme.
Biological function summary	Thrombomodulin regulates blood coagulation and maintains hemostatic balance. It forms a complex with thrombin on endothelial cells facilitating the conversion of protein C to activated protein C (APC). Activated protein C plays a pivotal role in controlling blood clot formation by proteolytically inactivating Factors Va and VIIIa. This function is important for preventing excessive clotting while ensuring proper wound healing processes.
Pathways	Thrombomodulin works chiefly within the anticoagulation pathway. It interacts directly with the thrombin-protein C pathway to achieve the activation of protein C. Additionally it interfaces with other pathways sharing mutual components such as endothelial cell proliferation and inflammation modulation. Related proteins include thrombin and protein C along with protein S which serves as a cofactor for APC's anticoagulant actions.
Associated diseases and disorders	Thrombomodulin dysregulation can lead to thrombosis and vascular inflammation. A deficiency or altered expression of thrombomodulin has been linked to thromboembolic disorders and atherosclerosis. In these conditions improper activation of protein C in the presence of decreased thrombomodulin function can exacerbate clot formation. Monitoring the levels of proteins like thrombin and protein C within these diseases can provide insights into endothelial dysfunction and potential therapeutic approaches.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

4 product images




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Thrombomodulin antibody [EPR18217] (ab187075)

Thrombomodulin Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-Thrombomodulin antibody

Immunohistochemical analysis of paraffin-embedded mouse lung tissue labeling Thrombomodulin with ab187075 at 1/2000 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) ready to use. Positive staining on endothelial cells of mouse lung (PMID: 23946288). Counter stained with Hematoxylin.
Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG H&L (HRP) ready to use.

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Thrombomodulin antibody [EPR18217] (ab187075)

Thrombomodulin Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-Thrombomodulin antibody

Immunohistochemical analysis of paraffin-embedded rat lung tissue labeling Thrombomodulin with ab187075 at 1/2000 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) ready to use. Positive staining on endothelial cells of rat lung (PMID: 23946288). Counter stained with Hematoxylin.
Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG H&L (HRP) ready to use.


Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.



Flow Cytometry - Anti-Thrombomodulin antibody [EPR18217] (ab187075)

Thrombomodulin Flow Cytometry staining using rabbit Anti-Thrombomodulin antibody

Flow cytometric analysis of bEND.3 (mouse brain endothelioma cell line) cell line labeling Thrombomodulin with ab187075 at 1/60 dilution (red) compared with a Rabbit IgG, monoclonal [EPR25A] - Isotype Control (ab172730) (black) and an unlabeled control (cells without incubation with primary antibody and secondary antibody) (blue). Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) (ab150077) at 1/2000 dilution was used as the secondary antibody.
Total viable cells were gated for the FC image.



Western blot - Anti-Thrombomodulin antibody [EPR18217] (ab187075)

Exposure time : Lane 1: 3 minutes; Lane 2: 15 seconds; Lane 3: 30 seconds; Lanes 4-5: 3 seconds.

Blocking/Dilution buffer: 5% NFDM/TBST.

The molecular weight observed is consistent with what has been described in the literature (PMID: 9619877).

All lanes:
Western blot - Anti-Thrombomodulin antibody [EPR18217] (ab187075) at 1/1000 dilution

Lane 1:
Rat heart lysate at 20 µg

Lane 2:
bEND.3 (mouse brain endothelioma cell line) whole cell lysate at 20 µg

Lane 3:
Mouse placenta lysate at 20 µg

Lane 4:
Rat lung lysate at 20 µg

Lane 5:
Mouse lung lysate at 20 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/100000 dilution

Developed using the ECL technique.

Predicted band size: 60 kDa

Observed band size: 105 kDa, 75 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.