abcam

ab187262

Anti-PAI1 antibody [EPR17795]

Recombinant | RabMAb | KO Validated | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR17795
Purification technique	Affinity purification Protein A
Concentration	0.638 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

IP

Tested

Species	Human
Dilution info	1/40
Notes	-

Not recommended

Species	Mouse, Rat
Dilution info	-
Notes	-

WB

Tested

Species	Human
Dilution info	1/1000
Notes	-

Not recommended

Species	Mouse, Rat
Dilution info	-
Notes	-

ICC/IF

Tested

Species	Human
Dilution info	1/500
Notes	-

Not recommended

Species	Mouse, Rat
Dilution info	-
Notes	-

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	-
Notes	-

Not recommended

Species	Mouse, Rat
Dilution info	-
Notes	-

IHC-P

Not recommended

Species	Human, Mouse, Rat
Dilution info	-
Notes	-

Target data

See full target information SERPINE1

Function	Serine protease inhibitor. Inhibits TMPRSS7 (PubMed:15853774). Is a primary inhibitor of tissue-type plasminogen activator (PLAT) and urokinase-type plasminogen activator (PLAU). As PLAT inhibitor, it is required for fibrinolysis down-regulation and is responsible for the controlled degradation of blood clots (PubMed:17912461, PubMed:8481516, PubMed:9207454, PubMed:21925150). As PLAU inhibitor, it is involved in the regulation of cell adhesion and spreading (PubMed:9175705). Acts as a regulator of cell migration, independently of its role as protease inhibitor (PubMed:15001579, PubMed:9168821). It is required for stimulation of keratinocyte migration during cutaneous injury repair (PubMed:18386027). It is involved in cellular and replicative senescence (PubMed:16862142). Plays a role in alveolar type 2 cells senescence in the lung (By similarity). Is involved in the regulation of cementogenic differentiation of periodontal ligament stem cells, and regulates odontoblast differentiation and dentin formation during odontogenesis (PubMed:25808697, PubMed:27046084).

Storage

Shipped at conditions	Conditional Ambient

Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.


9 product images



Immunocytochemistry/ Immunofluorescence - Anti-PAI1 antibody [EPR17795] (ab187262)

PAI1 Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-PAI1 antibody

Immunofluorescent analysis of 4% paraformaldehyde-fixed 0.1% Triton X-100 permeabilized HT1080 (Human fibrosarcoma cells) cells labeling PAI1 with ab187262 at 1/500 dilution followed by Goat anti-rabbit IgG (Alexa Fluor® 488) (ab150077) secondary antibody at 1/400 dilution (green). Cytoplasm staining on HT1080 cells was observed. The nuclear counter stain is DAPI (blue). Tubulin is detected with ab7291 (anti-Tubulin mouse mAb) at 1/500 dilution and ab150120 (AlexaFluor®594 Goat anti-Mouse secondary) at 1/500 dilution (red).
The negative controls are as follows:
-ve control 1: ab187262 at 1/500 dilution followed by ab150120 (AlexaFluor®594 Goat anti-Mouse secondary) at 1/500 dilution.
-ve control 2: ab7291 (anti-Tubulin mouse mAb) at 1/500 dilution followed by ab150077 (Alexa Fluor®488 Goat Anti-Rabbit IgG H&L) at 1/400 dilution.




Flow Cytometry (Intracellular) - Anti-PAI1 antibody [EPR17795] (ab187262)

PAI1 Flow Cytometry (Intracellular) staining using rabbit Anti-PAI1 antibody

ab187262 staining PAI1in the human cell line HepG2 (human hepatocellular carcinoma) by intracellular flow cytometry. Cells were fixed with 4% paraformaldehyde and the sample was incubated with the primary antibody at a dilution of 1/60. A goat anti rabbit IgG (Alexa Fluor® 488) at a dilution of 1/2000 was used as the secondary antibody.
Isoytype control: Rabbit monoclonal IgG (Black)

Unlabelled control: Cell without incubation with primary antibody and secondary antibody (Blue)




Immunocytochemistry/ Immunofluorescence - Anti-PAI1 antibody [EPR17795] (ab187262)

PAI1 Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-PAI1 antibody

Immunofluorescent analysis of 4% paraformaldehyde-fixed 0.1% Triton X-100 permeabilized HepG2 (Human liver hepatocellular carcinoma) cells labeling PAI1 with ab187262 at 1/500 dilution followed by Goat anti-rabbit IgG (Alexa Fluor® 488) (ab150077) secondary antibody at 1/400 dilution (green). Cytoplasm staining on HepG2 cells was observed. The nuclear counter stain is DAPI (blue). Tubulin is detected with ab7291 (anti-Tubulin mouse mAb) at 1/500 dilution and ab150120 (AlexaFluor®594 Goat anti-Mouse secondary) at 1/500 dilution (red).

The negative controls are as follows:
-ve control 1: ab187262 at 1/500 dilution followed by ab150120 (AlexaFluor®594 Goat anti-Mouse secondary) at 1/500 dilution.
-ve control 2: ab7291 (anti-Tubulin mouse mAb) at 1/500 dilution followed by ab150077 (Alexa Fluor®488 Goat Anti-Rabbit IgG H&L) at 1/400 dilution.

Immunoprecipitation - Anti-PAI1 antibody [EPR17795] (ab187262)




PAI1 was immunoprecipitated from 1mg of HepG2 (Human liver hepatocellular carcinoma) whole cell lysate with ab187262 at 1/40 dilution. Western blot was performed from the immunoprecipitate using ab187262 at 1/1000 dilution. VeriBlot for IP Detection Reagent (HRP) (ab131366), was used for detection at 1/1500 dilution.
Lane 1: HepG2 whole cell lysate 10 µg (Input). Lane 2: ab187262 IP in HepG2 whole cell lysate. Lane 3: Rabbit monoclonal IgG (ab172730) instead of ab187262 in HepG2 whole cell lysate.
Blocking and dilution buffer and concentration: 5% NFDM/TBST.

Exposure time: 30 seconds.

All lanes:
Immunoprecipitation - Anti-PAI1 antibody [EPR17795] (ab187262)

Predicted band size: 44 kDa, 45 kDa

Western blot - Anti-PAI1 antibody [EPR17795] (ab187262)



Blocking/Dilution buffer: 5% NFDM/TBST.
All lanes:
Western blot - Anti-PAI1 antibody [EPR17795] (ab187262) at 1/5000 dilution

All lanes:
HepG2 (Human liver hepatocellular carcinoma) whole cell lysate at 20 µg


Secondary

All lanes:
Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/1000 dilution

Predicted band size: 45 kDa

Observed band size: 45 kDa

Exposure time: 1min



Western blot - Anti-PAI1 antibody [EPR17795] (ab187262)

PAI1 Western blot staining using rabbit Anti-PAI1 antibody

Blocking/Dilution buffer: 5% NFDM/TBST.
All lanes:
Western blot - Anti-PAI1 antibody [EPR17795] (ab187262) at 1/1000 dilution

Lane 1:
Human fetal liver lysate at 10 µg

Lane 2:
Human fetal spleen lysate at 10 µg


Secondary

All lanes:
Anti-Rabbit IgG (HRP), specific to the non-reduced form of IgG at 1/1000 dilution

Predicted band size: 45 kDa

Observed band size: 45 kDa

Exposure time: 3min



Western blot - Anti-PAI1 antibody [EPR17795] (ab187262)

PAI1 Western blot staining using rabbit Anti-PAI1 antibody

Western blot: Anti-PAI1 antibody [EPR17795] ab187262 staining at 1/1000 dilution, shown in green; Mouse anti GAPDH ab8245 loading control staining at 1/20000 dilution, shown in magenta. A band was observed at 45 kDa in Wild-type U-87 MG cell lysates with no signal observed at this size in SERPINE1 knockout U-87 MG cell line. To generate this image, samples were run on an SDS-PAGE gel then transferred onto a nitrocellulose membrane. Membranes were blocked in 3 % Milk in TBS-0.1 % Tween® 20 (TBS-T) before incubation with primary antibodies overnight at 4 °C. Blots were washed four times in TBS-T, incubated with secondary antibodies for 1 h at room temperature, washed again four times then imaged. Secondary antibodies used were Goat anti-Rabbit 800CW and Goat anti-Mouse 680RD at 1/20,000 dilution.
All lanes:
Western blot - Anti-PAI1 antibody [EPR17795] (ab187262) at 1/1000 dilution

Lane 1:
Wild-type U-87 MG cell lysates at 20 µg

Lane 2:
Western blot - Human SERPINE1 knockout U-87 MG cell line (ab306851) at 20 µg

Lane 3:
A549 cell lysates at 20 µg

Lane 4:
HEK-293 cell lysates at 20 µg


Secondary

All lanes:
Goat anti-Rabbit 800CW & Goat anti-Mouse 680RD at 1/20000 dilution

Performed under reducing conditions.

Predicted band size: 45 kDa

Observed band size: 45 kDa



Western blot - Anti-PAI1 antibody [EPR17795] (ab187262)

Lanes 1 - 4: Merged signal (red and green). Green - ab187262 observed at 48 kDa. Red - loading control ab8245 (Mouse anti-GAPDH antibody [6C5]) observed at 37 kDa.
ab187262 was shown to react with PAI1 in wild-type A549 cells in Western blot. The band observed in the edited lysate lane above 45 kDa is likely to represent SERPINE1 with an insertion. This has not been investigated further. Wild-type A549 and SERPINE1 knockout cell lysates were subjected to SDS-PAGE. Membranes were blocked in 3 % milk in TBS-T (0.1 % Tween®) before incubation with ab187262 and ab8245 (Mouse anti-GAPDH antibody [6C5]) overnight at 4 °C at a 1 in 1000 dilution and a 1 in 20000 dilution respectively. Blots were incubated with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preabsorbed (ab216773) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preabsorbed (ab216776) secondary antibodies at 1 in 20000 dilution for 1 h at room temperature before imaging.

All lanes:
Western blot - Anti-PAI1 antibody [EPR17795] (ab187262) at 1/1000 dilution

Lane 1:
Wild-type A549 cell lysate at 20 µg

Lane 2:
SERPINE1 knockout A549 cell lysate at 20 µg


Lane 3:
HUVEC cell lysate at 20 µg

Lane 4:
HEK-293 cell lysate at 20 µg

Performed under reducing conditions.

Predicted band size: 45 kDa

Observed band size: 48 kDa



Western blot - Anti-PAI1 antibody [EPR17795] (ab187262)

Blocking/Dilution buffer: 5% NFDM/TBST.
All lanes:
Western blot - Anti-PAI1 antibody [EPR17795] (ab187262) at 1/5000 dilution

All lanes:
Human PAI1 full length protein at 0.01 µg

Secondary

All lanes:
Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/1000 dilution

Predicted band size: 45 kDa

Observed band size: 70 kDa

Exposure time: 30s

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.