abcam

ab188340

Anti-Histone H2B (mono methyl K5) antibody [EPR18092]

Recombinant | RabMAb | Advanced Validation

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR18092
Purification technique	Affinity purification Protein A
Concentration	1.04 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

WB

Tested

Species	Mouse
Dilution info	1/5000
Notes	-

Species	Human
Dilution info	1/5000
Notes	-

Expected

Species	Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

PepArr

Tested

Species	Human
Dilution info	-
Notes	-

Expected

Species	Mouse, Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

ICC/IF

Tested

Species	Human
Dilution info	1/2000
Notes	-

Expected

Species	Mouse, Rat
Dilution info	Use at an assay dependent concentration.

Notes	-

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	1/250
Notes	-

Expected

Species	Mouse, Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

IHC-P

Tested

Species	Mouse
Dilution info	1/2000
Notes	-

Species	Rat
Dilution info	1/2000
Notes	-

Species	Human
Dilution info	1/2000
Notes	-

Target data

See full target information H2BC21 meK5

Function	Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require

DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling. Has broad antibacterial activity. May contribute to the formation of the functional antimicrobial barrier of the colonic epithelium, and to the bactericidal activity of amniotic fluid.

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended

We do not recommend this combination. It is not covered by our product promise.


We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

---

8 product images




This image is courtesy of an anonymous customer review

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Histone H2B (mono methyl K5) antibody [EPR18092] (ab188340)

Immunohistochemistry analysis (Formalin/PFA-fixed paraffin-embedded sections) of formaldehyde fixed PBS-Tween permeabilized human colon tumor tissue. Stained with ab188340 at 1/2000 dilution. Secondary antibody used was HRP Agilent Detection System, Rabbit/Mouse. Blocking was done with 5% serum for 1 hour at 21°C. The sample was incubated with the primary antibody and TBST for 1 hour at 21°C. Antigen retrieval method was heat mediated Citrate

---




Immunocytochemistry/ Immunofluorescence - Anti-Histone H2B (mono methyl K5) antibody [EPR18092] (ab188340)

Histone H2B (mono methyl K5) Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-Histone H2B (mono methyl K5) antibody

Immunofluorescent analysis of 4% paraformaldehyde-fixed 0.1% Triton X-100 permeabilized HeLa (Human epithelial cells from cervix adenocarcinoma) cells labeling Histone H2B (mono methyl K5) with ab188340 at 1/2000 dilution followed by Goat anti-rabbit IgG (Alexa Fluor® 488) (ab150077) secondary antibody at 1/500 dilution (green). Confocal image showing nuclear staining on HeLa cell line. The nuclear counter stain is DAPI (blue). Tubulin is detected with ab7291 (anti-Tubulin mouse mAb) at 1/1000 dilution and ab150120 (AlexaFluor®594 Goat anti-Mouse secondary) at 1/500 dilution (red).
The negative controls are as follows:
-ve control 1: ab188340 at 1/2000 dilution followed by ab150120 (AlexaFluor®594 Goat anti-Mouse secondary) at 1/500 dilution.

-ve control 2: ab7291 (anti-Tubulin mouse mAb) at 1/1000 dilution followed by ab150077 (Alexa Fluor®488 Goat Anti-Rabbit IgG H&L) at 1/500 dilution.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Histone H2B (mono methyl K5) antibody [EPR18092] (ab188340)

Histone H2B (mono methyl K5) Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-Histone H2B (mono methyl K5) antibody

Immunohistochemical analysis of paraffin-embedded Human colon tissue labeling Histone H2B (mono methyl K5) with ab188340 at 1/2000 dilution followed by Goat Anti-Rabbit IgG H&L (HRP) (ab97051) secondary antibody at 1/500 dilution. Nucleus staining on Human colon tissue is observed. Counter stained with Hematoxylin.
Secondary antibody only control: Used PBS instead of primary antibody secondary antibody is Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/500 dilution.


Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.



Flow Cytometry (Intracellular) - Anti-Histone H2B (mono methyl K5) antibody [EPR18092] (ab188340)

Histone H2B (mono methyl K5) Flow Cytometry (Intracellular) staining using rabbit Anti-Histone H2B (mono methyl K5) antibody

Intracellular Flow Cytometry analysis of HeLa (human cervix adenocarcinoma) cells labelling Histone H2B (red) with purified ab188340 at a dilution of 1/250. The secondary antibody used was Alexa Fluorr® 488 goat-anti-rabbit IgG (1/2000). Cells were fixed with 4% paraformaldehyde and permeabilised with 90% methanol. Isotype control antibody was rabbit monoclonal IgG (black). The blue line shows cells without incubation with primary and secondary antibody.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Histone H2B (mono methyl K5) antibody [EPR18092] (ab188340)

Histone H2B (mono methyl K5) Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-Histone H2B (mono methyl K5) antibody

Immunohistochemical analysis of paraffin-embedded Mouse colon tissue labeling Histone H2B (mono methyl K5) with ab188340 at 1/2000 dilution followed by Goat Anti-Rabbit IgG H&L (HRP) (ab97051) secondary antibody at 1/500 dilution. Nucleus staining on mouse colon tissue is observed. Counter stained with Hematoxylin.
Secondary antibody only control: Used PBS instead of primary antibody secondary antibody is Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/500 dilution.

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Histone H2B (mono methyl K5) antibody [EPR18092] (ab188340)

Histone H2B (mono methyl K5) Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-Histone H2B (mono methyl K5) antibody

Immunohistochemical analysis of paraffin-embedded Rat colon tissue labeling Histone H2B (mono methyl K5) with ab188340 at 1/2000 dilution followed by Goat Anti-Rabbit IgG H&L (HRP) (ab97051) secondary antibody at 1/500 dilution. Nucleus staining on rat colon tissue is observed. Counter stained with Hematoxylin.
Secondary antibody only control: Used PBS instead of primary antibody secondary antibody is Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/500 dilution.

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.



Western blot - Anti-Histone H2B (mono methyl K5) antibody [EPR18092] (ab188340)

Histone H2B (mono methyl K5) Western blot staining using rabbit Anti-Histone H2B (mono methyl K5) antibody

Blocking/Dilution buffer: 5% NFDM/TBST.
All lanes:
Western blot - Anti-Histone H2B (mono methyl K5) antibody [EPR18092] (ab188340) at 1/5000 dilution

Lane 1:
NIH/3T3 (Mouse embyro fibroblast cells) whole cell lysate at 10 µg

Lane 2:
HeLa (Human epithelial cells from cervix adenocarcinoma) whole cell lysate at 10 µg


Secondary

All lanes:
Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/1000 dilution

Predicted band size: 14 kDa

Observed band size: 14 kDa

Exposure time: 15s



Peptide Array - Anti-Histone H2B (mono methyl K5) antibody [EPR18092] (ab188340)

ab188340 was tested in Peptide array against 501 different modified and unmodified histone peptides; each peptide is printed on the array at six concentrations (each in triplicate).
Circle area represents affinity between the antibody and a peptide: all antigen-containing peptides are displayed as red circles, all other peptides as blue circles. The affinity is calculated as area under curve when antibody binding values are plotted against the corresponding peptide concentration. Each circle area is normalized to the peptide with the strongest affinity.

The complete dataset, including full list of all peptides and information on the position of each peptide in the diagram, is available under the Product Protocol section.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.