abcam

ab192265

Anti-NDUFC2 antibody [EPR16499]

Recombinant | RabMAb

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR16499
Purification technique	Affinity purification Protein A
Concentration	0.192 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

WB

Tested

Species	Human
Dilution info	1/1000 - 1/2000
Notes	-

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	1/30
Notes	ab172730 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.

IHC-P

Tested

Species	Human
Dilution info	1/50 - 1/100
Notes	-

Target data

See full target information NDUFC2

Function	Accessory subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I), that is believed not to be involved in catalysis but required for the complex assembly. Complex I functions in the transfer of electrons from NADH to the respiratory chain. The immediate electron acceptor for the enzyme is believed to be ubiquinone.

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	NDUFC2 also known as NADH:ubiquinone oxidoreductase subunit C2 is a protein component of the mitochondrial complex I. It has a molecular mass of approximately 14.4 kDa. This protein is expressed within the mitochondria of various tissues with significant expression observed in high energy-demanding tissues such as the brain heart and muscles. NDUFC2 does not have a catalytic role but contributes to the structural stability of complex I.
Biological function summary	NDUFC2 plays a role as part of the mitochondrial respiratory chain complex I also known as NADH:ubiquinone oxidoreductase. This complex catalyzes the initial step of electron transport helping transfer electrons from NADH to ubiquinone. By contributing to this process NDUFC2 supports ATP production through oxidative phosphorylation which is essential for cellular energy metabolism. Being part of such a complex structure NDUFC2 interacts closely with other subunits of complex I.
Pathways	NDUFC2 involves itself in the oxidative phosphorylation and the citric acid cycle. These pathways are vital for energy conversion and metabolic processes. Within oxidative phosphorylation NDUFC2 relates to proteins such as NDUFB10 and NDUFS1 which are additional components of complex I. This protein along with its complex counterparts ensures efficient electron transport and ATP synthesis.
Associated diseases and disorders	NDUFC2 associates with mitochondrial disorders that stem from complex I deficiencies. Such diseases include Leigh syndrome a severe neurological disorder and mitochondrial encephalomyopathy. Interactions with other complex I proteins like NDUFS3 and NDUFV1 highlight the role of NDUFC2 in these conditions. Mutations or deficiencies in NDUFC2 can impact cellular energy metabolism contributing to the pathophysiology of these disorders.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.


3 product images



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-NDUFC2 antibody [EPR16499] (ab192265)

NDUFC2 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-NDUFC2 antibody


Immunohistochemical analysis of paraffin-embedded Human kidney tissue labeling NDUFC2 with ab192265 at 1/100 dilution followed by prediluted HRP Polymer for Rabbit/Mouse IgG. Counter stained with Hematoxylin.
Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.



Flow Cytometry (Intracellular) - Anti-NDUFC2 antibody [EPR16499] (ab192265)

NDUFC2 Flow Cytometry (Intracellular) staining using rabbit Anti-NDUFC2 antibody

Intracellular flow cytometric analysis of 2% paraformaldehyde-fixed A549 cells labeling NDUFC2 with ab192265 at 1/30 dilution (pink) compared with a rabbit monoclonal IgG isotype control (green). Goat anti rabbit IgG (FITC) at 1/150 dilution was used as the secondary antibody.



Western blot - Anti-NDUFC2 antibody [EPR16499] (ab192265)

Blocking/Dilution buffer: 5% NFDM /TBST.
All lanes:
Western blot - Anti-NDUFC2 antibody [EPR16499] (ab192265) at 1/1000 dilution

Lane 1:
HeLa cell lysate at 20 µg

Lane 2:
HepG2 cell lysate at 20 µg

Lane 3:
A549 cell lysate at 20 µg

Secondary

All lanes:
Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/1000 dilution

Predicted band size: 14 kDa

Observed band size: 14 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.